



The Tennessee Wildlife Resources Agency has conducted annual turkey hunts on the Oak Ridge Reservation since 1997. Harvested turkeys are screened before release to evaluate potential radiation doses to consumers. Since 1977, less than 0.34 percent of harvested turkeys have been retained because of potential radiation contamination.

6

Oak Ridge Reservation Environmental Monitoring Program

ORR environmental surveillance is conducted to comply with DOE requirements to protect the public and the environment against undue risks associated with DOE activities. These requirements are established in DOE Order 458.1, *Radiation Protection of the Public and the Environment* (DOE 2020), and related guidance is provided in *Environmental Radiological Effluent Monitoring and Environmental Surveillance* (DOE 2015). The objective of the ORR environmental surveillance program is to characterize environmental conditions in areas outside the ORR facility boundaries, both on and off ORR.

6.1. Meteorological Monitoring

Eight meteorological towers provide data on meteorological conditions and on the transport and diffusion qualities of the atmosphere on ORR. Data collected at the towers are used in routine dispersion modeling to predict impacts from facility operations and as input to emergency response atmospheric models, which are used for simulated and actual accidental releases from a facility. Data from the towers are also used to support various research and engineering projects. Additionally, ORNL and Y-12 operate three wind profilers on ORR to better characterize upper-level winds (winds higher than 60 m above ground level).

6.1.1. Data Collection and Analysis

The eight meteorological towers on ORR are described in Table 6.1 and depicted in Figure 6.1. In this document, ORR towers are designated by “MT” followed by a numeral. Other commonly used names for these towers are provided in Table 6.1. Meteorological data are collected at different heights above the ground (1, 2, 3, 10, 15, 30, 33, 35, and 60 m) to assess the vertical structure of the atmosphere, particularly with respect to wind shear and stability. Stable boundary

layers and significant wind shear zones associated with the local ridge-and-valley terrain and the Great Valley of Eastern Tennessee (see Appendix B) can significantly affect the movement of a plume after a facility release (Bowen et al. 2000). Data are collected at 10 or 15 m at most towers, but the wind measurement height is 25 m for MT11 and 20 m for MT13. Data are collected at some towers at 30, 33, 35, and 60 m.

Temperature, relative humidity, and precipitation are measured at most sites at 2 m, but wind speed and wind direction typically are not. Atmospheric stability (a measure of the vertical mixing properties of the atmosphere) is measured at most towers; however, measurements involving vertical temperature profiles (i.e., measurements made by the solar radiation delta-T method) limit accurate determination of nighttime stability to the 60 m towers. The solar radiation delta-T method is a stability calculation that involves the

temperature difference between 15 and 30 m heights, 15 m wind speeds, and the solar radiation value. Stability is also calculated for most sites using the sigma phi method, which relies heavily on the measurement of the standard deviation of vertical wind speed using 3D sonic wind monitors. Barometric pressure is measured at one or more of the towers at each ORR site (MT2, MT4, MT6, MT9, MT12, and MT13). Precipitation is measured at MT6 and MT9 at the Y-12 Complex; at MT13 at ETP; and at MT2, MT3, MT4, and MT12 at ORNL. Solar radiation is measured at MT6 and MT9 at the Y-12 Complex and at MT2, MT4, and MT12 at ORNL. Instrument calibrations are managed by UT-Battelle, LLC, and are performed every 6 months by qualified instrument technicians. Additionally, Holian Environmental audits the Y-12-owned towers (MT6, MT9, and MT11) every 3 months.

Table 6.1. ORR meteorological towers


Tower	Alternate tower names	Location (latitude, longitude)	Altitude above MSL (m)	Measurement heights (m)
ETTP				
MT13	J, YEOC	35.93043N, -84.39346W	237	20
ORNL				
MT2	D, 1047	35.92559N, -84.32379W	261	1, 2, 15, 35, 60
MT3	B, 6555	35.93273N, -84.30254W	256	15, 30
MT4	A, 7571	35.92185N, -84.30470W	266	1, 3, 15, 30
MT12	F	35.95285N, -84.30314W	354	10
Y-12 Complex				
MT6	W, West	35.98058N, -84.27358W	326	2, 10, 30, 60
MT9	Y, PSS Tower	35.98745N, -84.25363W	290	2, 15, 33
MT11	S, South Tower	35.98190N, -84.25504W	352	25

^a Tower "C" before May 2014.

Acronyms:

ETTP = East Tennessee Technology Park
 MSL = mean sea level
 ORNL = Oak Ridge National Laboratory
 ORR = Oak Ridge Reservation

PSS = plant shift superintendent
 Y-12 Complex = Y-12 National Security Complex
 YEOC = Y-12 Complex Emergency Operations Center

**Acronyms:**

CTF = Central Training Facility

ETTP = East Tennessee Technology Park

LIDAR = light detection and ranging

ORNL = Oak Ridge National Laboratory

ORR = Oak Ridge Reservation

SODAR = sonic detection and ranging

Y-12 = Y-12 National Security Complex

Figure 6.1. The ORR meteorological monitoring network, including light and sonic detection and ranging devices and the Central Training Facility sensor

Sonic detection and ranging (SODAR) devices have been installed at the east end of the Y-12 Complex (Pine Ridge). The SODAR devices use acoustic waves to estimate wind direction, wind speed, and turbulence at altitudes higher than the reach of meteorological towers (40–800 m above ground level). Although SODAR measurements are somewhat less accurate than measurements made on the meteorological towers, the SODAR devices provide useful information regarding stability, upper-air winds, and mixing depth. Mixing depth is the thickness of the air layer adjacent to the

ground over which an emitted or entrained inert nonbuoyant tracer could be mixed by turbulence within 1 h. Similar to SODAR but slightly more accurate, light detection and ranging (LIDAR) is a remote sensing technology that uses lasers to measure meteorological conditions.

Meteorological data are collected in real time from the meteorological towers at 2 min, 15 min, and 1 h average intervals for emergency response purposes and for dispersion modeling at the ORNL and Y-12 Complex Emergency Operations Centers.

Annual dose estimates are calculated using the archived hourly data. Data quality is checked continuously against predetermined data constraints, and out-of-range parameters are marked as invalid and excluded from compliance modeling. Appropriate substitution data are identified when possible. Quality assurance records of missing and erroneous data are routinely kept for the eight ORR towers.

6.1.2. Results

Prevailing winds generally flow up-valley from the southwest and west-southwest or down-valley from the northeast and east-northeast, patterns that typically result from channeling effects produced by the parallel ridges flanking the ORR sites. Winds in the valleys tend to follow the ridge axes, limiting cross-ridge flow within local valley bottoms. These conditions dominate over most of ORR, but flow variation is greater at ETTP, which is located within a less constrained open valley bottom.

On ORR, low wind speeds dominate near the valley surfaces largely because of the decelerating influence of nearby ridges and mountains. Wind acceleration is sometimes observed at ridgetop level, particularly when flow is not parallel to the ridges (see Appendix B).

The atmosphere over ORR is often characterized by stable conditions at night and for a few hours after sunrise. These conditions, when coupled with low wind speeds and channeling effects in the valleys, result in poor dilution of emissions from the facilities. However, high roughness values (caused by terrain and obstructions such as trees and buildings) may significantly mitigate these factors by increasing turbulence (atmospheric mixing). These features are

captured in dispersion model data input and are reflected in modeling studies conducted for each facility.

Precipitation data from MT2 have previously been used in streamflow modeling and in certain research efforts. The data indicate the variability of regional precipitation: the high winter rainfall resulting from frontal systems and the uneven but occasionally intense summer rainfall associated with frequent air mass thunderstorms. The total precipitation at ORNL during 2024 (1,227.9 mm, or 48.34 in.) was about 15 percent below the long-term 1991–2020 average of 1,417.8 mm (55.80 in.). The average annual wind data recovery rates (a measure of acceptable data) during 2024 were greater than 96 percent for MT3, MT4, and MT12. Tower MT2 was down a portion of the year because of maintenance and updates but still recorded a recovery rate greater than 70 percent in 2024. Missing data at tower MT2 were corrected through profiling and substitution along with consideration of ambient meteorological measurements and synoptic weather by on-site meteorologists.

6.2. Ambient Air Monitoring

In addition to exhaust stack monitoring conducted at ORR installations (see Chapters 3, 4, and 5), ambient air monitoring is performed to measure radiological parameters directly in the ambient air adjacent to the facilities (Figure 6.2). Ambient air monitoring provides a means to verify that contributions of fugitive and diffuse sources are insignificant, serves as a check on dose-modeling calculations, and would enable the determination of contaminant levels at monitoring locations in the event of an emergency.



Figure 6.2. ORR ambient air station


6.2.1. Data Collection and Analysis

Ambient air monitoring conducted by individual site programs is discussed in Chapters 3, 4, and 5. The ORR ambient air monitoring program complements the individual site programs and enables the impacts of ORR operations to be assessed on an integrated basis.

The objectives of the ORR ambient air monitoring program are to perform surveillance of airborne radionuclides at the reservation perimeter and to collect reference data from a location not affected by activities on ORR. The perimeter air monitoring network was established in the early 1990s and was modified in 2016 in response to changes in DOE activities and operations since the 1990s. The stations monitored in 2024 are shown in Figure 6.3. Reference samples are collected at Station 52 (Fort Loudoun Dam). Sampling was conducted at each ORR station during 2024 to quantify levels of alpha-, beta-, and gamma-emitting radionuclides.

Atmospheric dispersion modeling was used to select appropriate sampling locations likely to be affected most by releases from the Oak Ridge facilities. Therefore, in the event of a release, no residence or business near ORR should receive a radiation dose greater than doses calculated at the sampled locations.

The sampling system at each ambient air monitoring station consists of two separate instruments. Particulates are captured by a high-volume air sampler equipped with a glass-fiber filter. The filters are collected weekly, composited quarterly, and then submitted to an analytical laboratory to quantify gross alpha and gross beta activity and to determine the concentrations of specific isotopes of interest on ORR. The second instrument is designed to collect tritiated water vapor. The sampler consists of a prefilter followed by an adsorbent trap that contains indicating silica gel. The samples are collected weekly or biweekly, composited quarterly, and then submitted to an analytical laboratory for ^3H analysis.

**Notes:**

1. Reference samples are collected at Station 52 (Fort Loudoun Dam).
2. Station 7 is an ORNL site-specific monitoring location and is not part of the ORR perimeter network.

Acronyms:

ETTP = East Tennessee Technology Park
ORNL = Oak Ridge National Laboratory

ORR = Oak Ridge Reservation
Y-12 = Y-12 National Security Complex

Figure 6.3. Locations of ORR perimeter air monitoring stations

6.2.2. Results

Data from the ORR ambient air network are analyzed to assess the impact of DOE operations on the local air quality. Each measured radionuclide concentration (Table 6.2) is compared with derived concentration standards (DCSs) for air established by DOE as guidelines for controlling exposure to members of the public (DOE 2021a). All radionuclide concentrations measured at the ORR ambient air stations during 2024 were less than 1 percent of applicable DCSs.

6.3. External Gamma Radiation Monitoring

Members of the public could hypothetically be exposed directly to gamma radiation from radionuclides released into the environment, from previously released radionuclides deposited on soil and vegetation or in sediments, from radiation-generating facilities (especially high-energy accelerators), and from the storage of radioactive materials (DOE 2021b). Continuous direct radiation levels are monitored at locations around ORR to complement the sample data collected as part of the ORR ambient air monitoring program (see Section 6.2).

Table 6.2. Radionuclide concentrations at ORR perimeter air monitoring stations sampled quarterly, 2024

Station	Average concentration (pCi/mL) ^a (Number detects/n)									
	⁷ Be	²¹⁴ Bi	³ H	⁴⁰ K	²¹⁰ Pb	²¹² Pb	²¹⁴ Pb	⁹⁹ Tc	²⁰⁸ Tl	^{233/234} U ^{235/236} U ²³⁸ U
01 ^b	3.02E-08 (4/4)		5.38E-06 (3/4)	2.13E-10 (1/4)	1.58E-08 (3/4)					4.06E-11 2.37E-12 3.59E-11 (4/4) (2/4) (4/4)
02	2.68E-08 (4/4)	6.62E-11 (1/4)	6.36E-06 (4/4)	2.31E-10 (0/4)	1.83E-08 (4/4)	3.42E-11 (1/4)				3.31E-11 2.24E-12 3.89E-11 (4/4) (2/4) (4/4)
03	2.92E-08 (4/4)	9.01E-11 (1/4)	1.44E-05 (3/4)	9.32E-10 (2/4)	1.59E-08 (3/4)	3.80E-11 (1/4)	4.86E-11 (1/4)		2.44E-11 (1/4)	3.98E-11 4.11E-12 3.61E-11 (4/4) (2/4) (4/4)
09	2.81E-08 (4/4)	1.03E-10 (1/4)	1.77E-05 (4/4)	1.64E-09 (3/4)	2.22E-08 (4/4)		9.59E-11 (1/4)			4.10E-11 3.35E-12 3.49E-11 (4/4) (3/4) (4/4)
11	2.85E-08 (4/4)		5.83E-06 (3/4)	1.06E-09 (2/4)	2.23E-08 (4/4)					3.47E-11 2.20E-12 3.61E-11 (4/4) (2/4) (4/4)
35	2.73E-08 (4/4)		5.61E-06 (3/4)	7.22E-10 (1/4)	4.99E-09 (1/4)			1.62E-09 (1/4)	1.50E-11 (1/4)	4.49E-11 2.40E-12 3.33E-11 (4/4) (1/4) (4/4)
37	2.91E-08 (4/4)		2.66E-06 (3/4)	9.21E-10 (2/4)	1.78E-08 (4/4)					3.42E-11 3.23E-12 3.94E-11 (4/4) (1/4) (4/4)
40	2.74E-08 (4/4)		2.70E-06 (3/4)	1.60E-09 (4/4)	1.99E-08 (3/4)					4.36E-11 3.15E-12 3.71E-11 (4/4) (3/4) (4/4)
46	3.07E-08 (4/4)		3.27E-06 (3/4)	1.16E-09 (2/4)	1.94E-08 (3/4)					3.50E-11 2.65E-12 3.66E-11 (4/4) (2/4) (4/4)
49	2.82E-08 (4/4)		1.89E-06 (2/4)	1.57E-09 (3/4)	1.88E-08 (4/4)					3.86E-11 2.06E-12 3.75E-11 (4/4) (1/4) (4/4)
52 ^c	2.61E-08 (4/4)		1.46E-06 (1/4)	1.18E-09 (3/4)	1.08E-08 (2/4)			-4.36E-10 (0/4)		3.67E-11 2.67E-12 3.63E-11 (4/4) (2/4) (4/4)


^a 1 pCi = 3.7E-02 Bq.^b In one quarter of 2024, ⁶⁰Co was also detected at Station 01. The average concentration was 5.49E-11 pCi/mL.^c Station 52 is the reference location.

6.3.1. Data Collection and Analysis

External gamma exposure rates are continuously recorded every minute by dual-range Geiger-Müller tube detectors colocated with ORR ambient air stations 2, 3, 9, 11, 40, 46, 49, and 52 (see Section 6.2). The data are downloaded weekly and are averaged for the entire year. Figure 6.4 shows locations that were monitored during 2024; Table 6.3 summarizes the data for each station.

6.3.2. Results

The mean exposure rate for the reservation network in 2024 was 9.3 µR/h, and the mean rate at the reference location (Fort Loudoun Dam) was 8.4 µR/h. Background direct radiation exposure rates have been collected at the Fort Loudoun Dam (Station 52) reference location for many years. From 2014 through 2023, the exposure rates at the reference location ranged from 8.0 to 11.4 µR/h and averaged 9.2 µR/h.



Note:

Reference samples are collected at Station 52 (Fort Loudoun Dam).

Acronyms:

ETTP = East Tennessee Technology Park

ORNL = Oak Ridge National Laboratory

ORR = Oak Ridge Reservation

Y-12 = Y-12 National Security Complex

Figure 6.4. External gamma radiation monitoring locations on ORR

Table 6.3. External gamma exposure rate averages for ORR, 2024

Air station number	Number of data points (daily)	Measurement ($\mu\text{R}/\text{h}$) ^a		
		Min	Max	Mean
02	359	7.6	9.8	8.4
03	359	8.0	10.2	8.8
09	366	8.0	16	9.0
11	360	8.5	11.3	9.8
40	366	8.3	10.7	9.5
46	362	9.1	11.5	10.2
49	365	8.3	11.1	9.3
52	366	7.6	9.8	8.4

^a To convert microrentgens per hour ($\mu\text{R}/\text{h}$) to milliroentgens per year, multiply by 8.760.

6.4. Surface Water Monitoring

The ORR surface water monitoring program consists of sample collection and analysis from four locations on the Clinch River, including public water intakes (Figure 6.5). The program is conducted in conjunction with site-specific surface water monitoring activities to enable an assessment of the impacts of past and current DOE operations on the quality of local surface water.


6.4.1. Data Collection and Analysis

Grab samples are collected quarterly at all four locations and are analyzed for general water quality parameters, screened for radioactivity, and analyzed for mercury and specific radionuclides when appropriate. Table 6.4 lists the locations and associated sampling frequencies and parameters.

In 2022, a more sensitive analytical method for determining mercury concentrations in surface

water samples was adopted. The new method can detect concentrations near 0.2 ng/L, whereas the detection limit for the previous method is about 67 ng/L. As expected, the ability to detect mercury at much lower levels resulted in detections in 8 of the 12 surface water samples collected for mercury analyses in 2024, while in the past, with the less sensitive method, mercury was rarely detected. At the sampling locations, the Clinch River is classified by the State of Tennessee for multiple uses, including recreation and domestic water supply (TDEC 2024a). These two designated uses have numeric Tennessee water quality criteria (WQCs) related to protection of human health. The WQCs are used as references where applicable (TDEC 2024b). The Tennessee WQCs do not include criteria for radionuclides. Four percent of the DOE DCS is used as the criterion for radionuclide comparison (DOE 2021a).

3/12/2021 - EPSDM004/Hf9

**Acronyms:**

CRK = Clinch River kilometer

ETTP = East Tennessee Technology Park

ORNL = Oak Ridge National Laboratory

ORR = Oak Ridge Reservation

Y-12 = Y-12 National Security Complex

Figure 6.5. ORR surface water surveillance sampling locations**6.4.2. Results**

In 2024, as has been the case since 2009, no statistical differences were found in the concentrations of routinely monitored radionuclides in surface water samples collected from the Clinch River upstream and downstream of DOE inputs. No radionuclides were detected above 4 percent of the respective DCSs.

Mercury was detected in 8 of the 12 samples collected in 2024, including one sample from the location upstream of DOE inputs (Clinch River

kilometer [CRK] 66). Results from one sample collected from CRK 32 were below the method detection level. The annual mean mercury concentration at the location downstream of DOE inputs (CRK 16) was statistically significantly higher than the mean concentration at the location upstream of DOE inputs (CRK 66). However, detected mercury results at all locations were well below the Tennessee mercury WQC for recreational use (51 ng/L).

Table 6.4. ORR surface water sampling locations, frequencies, and parameters, 2024

Location ^a	Description	Frequency	Parameters
CRK 16	Clinch River downstream from all DOE ORR inputs	Quarterly	Mercury, gross alpha, gross beta, gamma scan, ³ H, field measurements ^b
CRK 32	Clinch River downstream from ORNL	Quarterly	Mercury, gross alpha, gross beta, gamma scan, total radioactive strontium, ³ H, field measurements ^b
CRK 58	Water supply intake for Knox County	Quarterly	Gross alpha, gross beta, gamma scan, ³ H, field measurements ^b
CRK 66	Melton Hill Reservoir above City of Oak Ridge water intake	Quarterly	Mercury, gross alpha, gross beta, gamma scan, total radioactive strontium, ³ H, field measurements ^b

^a Locations indicate the water body and distances upstream of the confluence of the Clinch and Tennessee Rivers (e.g., CRK 16 is 16 km upstream from the confluence of the Clinch River with the Tennessee River in the Watts Bar Reservoir).

^b Field measurements consist of dissolved oxygen, pH, and temperature.

Acronyms:

CRK = Clinch River kilometer

DOE = US Department of Energy

ORNL = Oak Ridge National Laboratory

ORR = Oak Ridge Reservation

6.5. Groundwater Monitoring

Work continued in 2024 to implement key recommendations from the *Groundwater Strategy for the U.S. Department of Energy Oak Ridge Reservation* (DOE 2013), which was agreed to in 2014 by DOE, EPA, and the Tennessee Department of Environment and Conservation (TDEC). Work performed during 2024 under the ORR Groundwater Program included preparation of a report on the installation of and data obtained during the first year of sampling from three multizone exit pathway groundwater monitoring wells in west Bethel Valley adjacent to the Clinch River (DOE 2024a). Work continued on site-scale groundwater flow models for ETTP.

6.5.1. Off-Site Groundwater Assessment

During fiscal year (FY) 2024, the Oak Ridge Office of Environmental Management continued to collect and analyze samples from the off-site groundwater monitoring well array west of the Clinch River adjacent to Melton Valley. In addition, exit pathway groundwater monitoring in Melton Valley is conducted as part of the Oak Ridge Office of Environmental Management program, including sampling at six multiport monitoring wells in

western Melton Valley (wells 4537, 4538, 4539, 4540, 4541, and 4542). The results of this monitoring are summarized in the *2025 Remediation Effectiveness Report* (DOE 2025).

DOE completed an off-site groundwater assessment project and issued a final report in October 2017 (DOE 2017). The project was a cooperative effort among the parties to the ORR Federal Facility Agreement to investigate off-site groundwater quality and potential movement. To follow up on work from the off-site groundwater assessment, DOE conducts annual sampling and analysis of groundwater from several off-site residential wells and springs. Off-site groundwater data are summarized in the *2025 Remediation Effectiveness Report* (DOE 2025).

6.5.2. Regional and Site-Scale Flow Model

During FY 2017, DOE completed a project to construct and calibrate a regional-scale groundwater flow model that encompasses ORR and adjacent areas. The regional model provides a framework to support creation of smaller, site-scale groundwater flow models for use in planning and monitoring the effectiveness of future cleanup decisions and actions. During FY 2024, DOE continued developing a groundwater flow and

solute transport model for the ETP site to support Comprehensive Environmental Response, Compensation, and Liability Act (CERCLA 1980) groundwater decision-making and conducted groundwater flow modeling for the Molten Salt Reactor Experiment site to support the development of a CERCLA decision for components of that facility not included in a previous CERCLA decision.

6.6. Food

Food sources are analyzed to evaluate potential radiation doses to consumers of local food crops, fish, and harvested game and to monitor trends in environmental contamination and possible long-term accumulation of radionuclides. Samples of hay, vegetables, milk, fish, deer, Canada geese, and turkeys are usually collected every year from areas that could be affected by activities on the reservation and from off-site reference locations. Milk was not collected in 2024 because no dairies were found in potential ORR deposition areas. Surveys are conducted annually to determine whether any dairies are operating in areas of interest.

The wildlife administrative release limits associated with deer, turkeys, and geese harvested on ORR are conservative and were established based on the “as low as reasonably achievable” principle to ensure that doses to consumers are managed at levels well below regulatory dose thresholds. This concept is not a dose limit but rather a philosophy that has the objective of maintaining exposures to workers, members of the public, and the environment below regulatory limits and as low as can be reasonably achieved. The administrative release limit of 5 pCi/g ^{137}Cs is based on the assumption that one person consumes all of the meat from a maximum-weight deer, goose, or turkey. This limit ensures that members of the public who harvest wildlife on the reservation will not receive significant radionuclide doses from that consumption pathway. In addition, a conservative administrative limit of 1.5 times background for gross beta activity has been established. This threshold is near the detection

limit for field measurements of $^{89/90}\text{Sr}$ in deer leg bone.

6.6.1. Hay

Eating beef and drinking milk obtained from cattle that eat hay are potential radiation exposure pathways to humans. Hay is sampled annually to characterize possible doses from this pathway.

6.6.1.1. Data Collection and Analysis

Hay from the eastern edge of ORR, which historically has been cut for off-site use, has been sampled since 2015 but was not harvested there in 2024. Therefore, a sample was collected from a location southwest of ORNL when hay was cut for off-site use there in December 2024. Analyses for gross alpha, gross beta, gamma emitters, and uranium isotopes were performed.

6.6.1.2. Results

No gamma-emitting radionuclides were detected above the minimum detectable activity except for the naturally occurring radionuclide ^{40}K . Uranium isotopes were not detected above minimum detectable activities in any of the samples. Analytical results for hay samples are provided in Table 6.5.

6.6.2. Vegetables

Contaminants may reach vegetation by deposition of airborne materials, uptake from soil, and deposition of materials contained in irrigation water. As available, food crops are sampled annually from garden locations that have the potential to be affected by airborne releases from ORR to evaluate possible radiation doses to consumers. Vegetables are also sampled from a reference location for comparison. If available, crops that represent broad-leaf systems (e.g., lettuce, turnip greens), root-plant-vegetable systems (e.g., tomatoes), and root-system vegetables (e.g., turnips, potatoes) are obtained from each location and analyzed for radionuclides. Vegetable availability varies greatly from year to year.

Table 6.5. Concentrations of radionuclides detected in hay,^a December 2024 (pCi/kg)^b

Radionuclide	Result
Gross alpha	c
Gross beta	11,900
⁷ Be	c
⁴⁰ K	14,000
²³⁴ U	c
²³⁵ U	c
²³⁸ U	c

^a Detected radionuclides are those at or above minimum detectable activity.

^b 1 pCi = 3.7×10^{-2} Bq.

^c Value was less than or equal to minimum detectable activity.

6.6.2.1. Data Collection and Analysis

In 2024, tomatoes were purchased from farms near ORR and from a reference location, and a mixture of kale, turnip greens, and cabbages was collected to represent broad-leaf systems. No root vegetables were available in 2024. The locations were chosen based on availability and the likelihood of being affected by routine releases from the Oak Ridge facilities. All vegetable samples were analyzed for gross alpha, gross beta, gamma emitters, and uranium isotopes.

6.6.2.2. Results

Analytical results for vegetable samples are provided in Table 6.6. No gamma-emitting radionuclides were detected above the minimum detectable activity except for the naturally occurring radionuclides ⁴⁰K and ⁷Be.

6.6.3. Milk

Milk is a potentially significant exposure pathway to humans for some radionuclides deposited from airborne emissions because of the relatively large surface area on which a cow can graze daily, the rapid transfer of milk from producer to consumer, and the importance of milk in the diet. Since 2016, no dairies in potential ORR deposition areas have been located, and no milk samples have been collected. Surveys to identify dairies in potential

deposition areas are conducted each year, and milk sampling will resume when dairy operations in appropriate areas are located.

6.6.4. Fish

Members of the public could be exposed to contaminants originating from DOE ORR activities by consuming fish caught in area waters. This potential exposure pathway is monitored annually by collecting fish from three locations on the Clinch River and by analyzing edible flesh for specific contaminants. The locations are as follows (Figure 6.6):

- Clinch River upstream from all DOE ORR inputs (CRK 70)
- Clinch River downstream from ORNL (CRK 32)
- Clinch River downstream from all DOE ORR inputs (CRK 16)

6.6.4.1. Data Collection and Analysis

Sunfish (*Lepomis macrochirus*, *L. auritus*, and *Ambloplites rupestris*) and catfish (*Ictalurus punctatus*) are collected from each of the three locations to represent both top-feeding and bottom-feeding predator species. In 2024, a composite sample of each of those species at each location was analyzed for selected metals; polychlorinated biphenyls (PCBs); ³H; gross alpha; gross beta; gamma-emitting radionuclides; total radioactive strontium; and isotopes of americium, neptunium, plutonium, and thorium. To accurately estimate exposure levels to consumers, only edible portions of the fish were submitted for analysis. Once every 5 years, additional radiological analyses are performed to confirm the dose calculations (Chapter 7). In 2024, additional radionuclides analyzed in fish samples included uranium isotopes and ⁹⁹Tc. Results are presented in Table 6.7.

TDEC issues advisories on consumption of certain fish species caught in specified Tennessee waters. The advisories apply to fish that could contain potentially hazardous contaminants. TDEC has issued a “do not consume” advisory for catfish in

the entire Melton Hill Reservoir, not just in areas that could be affected by ORR activities, because of PCB contamination. Similarly, TDEC has issued a precautionary advisory for catfish in the Clinch River arm of Watts Bar Reservoir because of PCB contamination (TDEC 2023). TDEC also issues

precautionary advisories for consumption of fish when mercury levels exceed 0.3 ppm (Denton 2007). As of 2023, the three locations on the Clinch River where ORR fish are collected do not have mercury “do not consume” advisories.

Table 6.6. Concentrations of radionuclides detected in tomatoes and leafy greens, 2024 (pCi/kg)^a

Location	Gross alpha	Gross beta	⁷ Be	⁴⁰ K	²³⁴ U	²³⁵ U	²³⁸ U
Leafy greens^b							
North of Y-12	c	2,060	314	2,450	c	c	0.784
East of ORNL	c	4,590	c	5,750	c	c	c
West of ETPP	c	4,210	c	5,740	c	c	c
South of ORNL	c	4,340	c	4,950	c	c	c
Reference location	c	11,400	c	14,100	16.3	c	c
Tomatoes							
North of Y-12	c	1,960	c	3,090	c	c	c
East of ORNL	c	2,040	c	2,250	c	c	c
West of ETPP	c	1,760	c	2,630	c	c	c
South of ORNL	c	1,820	c	2,280	c	c	c
Reference location	c	2,340	c	2,760	c	c	c

^a Detected radionuclides are those at or above minimum detectable activity. 1 pCi = 3.7 × 10⁻² Bq.

^b Cabbage was collected at the North of Y-12 and West of ETPP locations, and a mixture of kale and turnip greens was collected at the East of ORNL, South of ORNL, and reference locations.


^c Value was less than or equal to minimum detectable activity.

Acronyms:

ETPP = East Tennessee Technology Park

ORNL = Oak Ridge National Laboratory

Y-12 = Y-12 National Security Complex



Acronyms:

CRK = Clinch River kilometer

ETTP = East Tennessee Technology Park

ORNL = Oak Ridge National Laboratory

ORR = Oak Ridge Reservation

Y-12 = Y-12 National Security Complex

Figure 6.6. Fish sampling locations for the ORR surveillance program

Table 6.7. Tissue concentrations in catfish and sunfish for detected PCBs and radionuclides, 2024^a

	CRK 16 Downstream		CRK 32		CRK 70 Upstream	
	Catfish	Sunfish	Catfish	Sunfish	Catfish	Sunfish
Metals (mg/kg)						
Mercury	c	c	c	c	B0.195 ^b	c
PCBs (µg/kg)						
Aroclor 1248	28.7	c	c	c	c	c
Aroclor 1254	87.1	J9.11 ^d	J17 ^d	c	33.2	c
Aroclor 1260	70.5	J9.5 ^d	22.7	c	44.6	c
Radionuclides (pCi/g)						
Beta activity	3.43	3.16	2.61	3.52	2.58	3.49
⁴⁰ K	2.99	3.32	3.33	3.41	3.5	4.22

^a Only parameters that were detected for at least one species are listed in the table.

^b “B” indicates that the analyte was detected in the associated method blank.

^c Value was less than or equal to minimum detectable activity.

^d “J” indicates that the result is an estimated value.

Acronyms:

CRK = Clinch River kilometer

PCB = polychlorinated biphenyl

6.6.4.2. Results

PCBs, specifically Aroclors 1260 and 1254, were detected in sunfish at CRK 16 and in catfish at CRK 32 in 2024. Mercury was detected in catfish at CRK 70 upstream of DOE inputs; the lab reported that mercury was detected in the associated method blank, indicating this result may be biased high. Mercury was not detected above the minimum detectable level at any other location in 2024. These results are consistent with the TDEC advisories. Detected PCBs, mercury, and radionuclide concentrations are shown in Table 6.7.

6.6.5. White-Tailed Deer

In 2024, three quota deer hunts were conducted on ORR: November 2 and 3, November 9 and 10, and December 14 and 15. Each hunt was limited to 450 shotgun/muzzleloader permittees and 600 archery permittees. UT-Battelle staff; Tennessee Wildlife Resources Agency personnel; and student members of the Wildlife Society, University of Tennessee chapter, performed most of the necessary operations at the checking station.

6.6.5.1. Data Collection and Analysis

In 2024, approximately 23,000 acres were available to deer hunters in the Oak Ridge Wildlife Management Area, which encompasses some properties not owned by DOE, including Haw Ridge Park (City of Oak Ridge) and the University of Tennessee Arboretum.

6.6.5.2. Results

The total ORR deer harvest in 2024 was 192, of which 114 (~59.4 percent) were bucks and 78 (~40.6 percent) were does. The heaviest buck weighed 185 lb, the heaviest doe weighed 121 lb, and the greatest number of antler points was 11. No deer were retained for exceeding the administrative release limit of 1.5 times background for beta activity in bone (~20 pCi/g ^{89/90}Sr) or for exceeding 5 pCi/g ¹³⁷Cs in edible tissue.

Compared with the deer harvests from 2020 to 2023, the 2024 total harvest of 192 was more consistent with prepandemic numbers. In 2019, the total harvest was 221 deer (~56.6 percent

bucks and ~43.4 percent does), the heaviest buck was 181 lb, the heaviest doe was 112 lb, and greatest number of antler points was 13.

Since 1985, 14,070 deer have been harvested from the Oak Ridge Wildlife Management Area, of which 220 (~2 percent) have been retained because of potential radiological contamination. The heaviest buck ever harvested weighed 218 lb (1998), and the heaviest doe ever harvested weighed 139 lb (1985). The average weight of all harvested deer is approximately 87 lb. (All weights are field-dressed weights.) The oldest deer harvested was a doe estimated to have been 12 years old (1989); the average age of all harvested deer is approximately 2 years. See the ORR hunt information website [here](#) for more information.

6.6.6. Waterfowl

The consumption of waterfowl is a potential pathway for exposing members of the public to radionuclides released from ORR operations. During the fall statewide season in 2024, Canada goose hunting was allowed on the Three Bend Area of ORR (shorelines only, excluding Gallaher Bend) 0.5 h before sunrise until noon on September 5, 10, 11, 17, and 18 and on October 8, 9, 15, and 16. Hunting was allowed for wood duck and teal on September 9 and 10.

6.6.6.1. Data Collection and Analysis

Canada geese are rounded up each summer for noninvasive gross radiological surveys to characterize concentrations of gamma-emitting radionuclides accumulated by waterfowl that feed and live on ORR.

6.6.6.2. Results

During the June 13, 2024, roundup on ORR, 22 geese (16 adults; 6 juveniles) were captured. All 22 captured geese were subjected to live whole-body gamma scans. Gamma scan results showed that all were well below the administrative release limit of 5 pCi/g ^{137}Cs .

6.6.7. Wild Turkey

Two wild turkey quota hunts were conducted April 13–14 and April 20–21. None of the 37 (34 adults; 3 juveniles) total turkeys harvested was retained for potential radiological contamination.

Since 1997, 1,007 turkeys have been harvested on spring turkey hunts. Twelve additional turkeys have been harvested since 2012 by archery hunters during fall deer hunts. The largest turkey ever harvested on ORR weighed 25.7 lb (harvested in 2009). Of all turkeys harvested, only three (less than 0.34 percent) have been retained because of potential radiological contamination: one in 1997, one in 2001, and one in 2005. Additional information is available on the ORR hunt website [here](#).

6.7. Habitat Quality Improvement

Maintaining ecosystems; protecting natural areas; and ensuring functioning of support infrastructure, such as power and communications rights-of-way, roadways, and waterways, through active management are important not only in natural areas but in developed areas as well. ORR conservation needs include control of invasive, non-native plants and animals; restoration of pollinator habitats; forest restoration and conservation; and creation and management of mitigation areas. DOE created the *Conservation Action Plan* (DOE 2021c) to be updated annually; this plan includes climate adaptation and resilience research, fish and wildlife habitat conservation and restoration, and invasive plant and animal management, among other projects.

6.7.1. Invasive Plant Management

Invasive, non-native plant species are among the greatest ecological threats to the United States and around the world. Invasive plants can threaten forests, wetlands, cultural resources, and other resources by increasing the risk of fire and storm damage and by encroaching onto roads, railroads,

power structures, waterways, and agricultural sites. To address these threats, the Federal Noxious Weed Act (1974) was amended and incorporated into the Federal Plant Protection Act (2000). This act mandates federal agencies to develop and coordinate management programs to control invasive plants on lands under their respective jurisdictions and to adequately fund integrated pest management plans. The ORNL Natural Resources Management Program has maintained an invasive plant management plan for ORR since 2004. Details of federal and state laws and regulations driving this plan can be found in technical manuscripts ORNL/TM-2004/98 (Parr et al. 2004), ORNL/TM-2004/98/R1 (Quarles et al. 2011), and ORNL/TM-2004/98/R2 (McCracken and Giffen 2017).

ORNL/TM-2001/113, *Assessment of Nonnative Invasive Plants in the DOE Oak Ridge National Environmental Research Park* (Drake et al. 2002), details the results of extensive survey efforts. These and subsequent surveys have been performed to identify invasive plant problems on ORR. Data from the surveys are used to develop control plans identifying which invasive species to target and in which locations.

More than 1,100 species of plants are found on ORR, and of these, approximately 179 plant species are non-native. Fifty-nine aggressive non-native (invasive) plant species have been identified on ORR, but control efforts have been primarily focused on the subset of 12 species shown in Table 6.8. These target species have been found across ORR in disturbed areas; on power line and gas line rights-of-way; throughout riparian buffer zones; and along state highways, railroad lines, and remote-access fire roads. They have invaded natural areas to varying degrees, causing vast ecological harm in plant and animal communities. In concert with efforts to control the 12 highly invasive species, other invasive plant species are also targeted for control using *Early Detection and Rapid Response* (DOI 2020). In 2023 and 2024, two new incursions of aggressively invasive plant species, cogon grass (*Imperata cylindrical*) and coltsfoot (*Tussilago farfara*), were identified and treated on ORR. The locations

where these incursions occurred were marked for future surveys and treatments.

Table 6.8. Twelve most problematic invasive plants on ORR

Common name	Scientific name
Japanese grass, Nepal grass	<i>Microstegium vimineum</i>
Japanese honeysuckle	<i>Lonicera japonica</i>
Chinese privet	<i>Ligustrum sinense</i>
Kudzu	<i>Pueraria montana</i>
Multiflora rose	<i>Rosa multiflora</i>
Tree of heaven	<i>Ailanthus altissima</i>
Autumn olive	<i>Elaeagnus umbellata</i>
Oriental bittersweet	<i>Celastrus orbiculatus</i>
Princess tree	<i>Paulownia tomentosa</i>
Winter creeper	<i>Euonymus hederaceus</i>
Bradford/Callery pear	<i>Pyrus calleryana</i>
Mimosa	<i>Albizia julibrissin</i>

The 32,800-acre DOE ORR consists mostly of undeveloped land, including forests, wetlands, open waterways, riparian buffer zones, and several hundred acres of grassland communities and fallow fields. Three major developed facilities lie within ORR boundaries—ORNL, the Y-12 Complex, and ETPP. Surrounding these developed facilities and interspersed throughout ORR are safety and security areas, utility corridors, access roads, research and education areas, cultural and historic preservation sites, waste burial grounds, monitoring sites, fire roads, emergency evacuation corridors, new facility construction and laydown areas, and public use areas. This multiplicity of land uses presents challenges for effectively preventing and managing invasive species.

Numerous DOE contractors have responsibilities for managing portions of ORR land, as do other federal and state agencies such as the Tennessee Valley Authority and the Tennessee Wildlife Resources Agency. A portion of the ORR Natural Resources Management Team's annual funding is designated for creation and implementation of an invasive plant management plan, which focuses mainly on control efforts in natural areas and

reference areas. However, efforts also include controlling specific invasive plant incursions into locations within and surrounding the campuses of developed facilities on ORR. The *Invasive Plant Management Plan for the Oak Ridge Reservation* (Parr et al. 2004) and its two subsequent revisions (Quarles et al. 2011; McCracken and Giffen 2017) explain options for addressing the problem of invasive plants on ORR and discuss selection of appropriate control measures. Areas selected for invasive plant control tend to cover several acres or are spread out across portions of ORR. Use of select herbicides is the most cost-effective treatment method in most cases, and the invasive plants present determine which herbicides will be most effective without harming surrounding native plant and animal habitats. Controlled burns are used in combination with herbicide applications to clear native grasslands, forest understory, kudzu infestations, and other select vegetation. Clearing these areas allows native plant regeneration and facilitates herbicide spot treatments in subsequent years. ORR fire protection and management programs are discussed in Section 6.8.

Invasive plant control on ORR has been conducted annually since 2003. Table 6.9 indicates the extent of annual invasive plant treatments.

Table 6.9. Annual invasive plant control on ORR, 2003–2024

Year	Area treated
2003	98 acres
2004	136 acres
2005	125 acres
2006	254 acres
2007	236 acres
2008	427 acres
2009	526 acres
2010	884 acres
2011	806 acres
2012	615 acres
2013	329 acres
2014	950 acres
2015	629 acres
2016	952 acres
2017	542 acres + 47 road miles
2018	507 acres + 53 road miles
2019	450 acres + 57 road miles
2020	400 acres + 65 road miles
2021	400 acres + 51 road miles
2022	266 acres + 77 road miles
2023	260 acres + 84 road miles
2024	348 acres + 75 road miles

Invasive plant management activities were completed in 2024 at the following ORR locations:

- Portions of First Creek and White Oak Creek riparian buffer zones, including the White Oak Lake weir
- Bethel Valley Road and Old Bethel Valley Road
- Haw Ridge former 3rd Street steam line and utility right-of-way access roads
- Fire protection roads across ORR
- Bethel Valley Road native grasslands
- Three Bend Wildland and Conservation Area fields, field edges, and roads, including the Gallaher Bend kudzu area
- Tower Shielding Facility roadsides and forest edges
- Pine Ridge and Chestnut Ridge overlooking the Y-12 campus (kudzu control)
- Midway Turnpike
- Gum Branch Road, McNew Hollow Road, Old County Road, and Watson Road
- Filled coal ash pond and slough kudzu area
- Wheat Church vista and trails
- Black Oak Ridge Conservation Easement roads, North Boundary Greenway, Raby Road, Gheen Road, North Boundary Road, and Poplar Creek Road
- McKinney Ridge kudzu and Blair Quarry kudzu areas
- Black Oak Ridge kudzu areas, including Black Oak Ridge Conservation Easement east and west kudzu areas
- Melton Valley Drive kudzu area
- Portions of the ETP Perimeter Road kudzu area
- Other roads and areas

6.7.2. Wetlands

Wetland determinations and delineations are conducted to facilitate compliance with TDEC and

US Army Corps of Engineers wetland protection requirements. In 2024, two determinations and delineations were completed on the ORNL campus to ensure project activities did not adversely affect wetlands. One was included in an Aquatic Resource Alteration Permit. A third wetland determination and delineation was completed off-site at the Black Oak Ridge Conservation Area to help facilitate and support ORNL research.

6.8. Fire Protection Management and Planning

Wildland fire management is an important part of DOE's overall management of ORR. A comprehensive wildfire management program has been established and implemented for the entire ORR. The *Oak Ridge Reservation Wildland Fire Management Plan* (WFMP) (DOE 2024b) assigns responsibilities for wildland fire management and is reviewed every 3 years and revised as needed. The *Oak Ridge Reservation Wildland Fire Implementation Plan* (DOE 2024c) contains details on program implementation. The WFMP was prepared to satisfy the requirements of DOE Order 420.1C, Change 3, *Facility Safety* (DOE 2019); DOE-STD-1066-2023, *Fire Protection* (DOE 2023); and relevant portions of Chapters 19 through 23 in National Fire Protection Association 1140, *Standard for Wildland Fire Protection* (NFPA 2022).

The WFMP outlines the overall goals and strategies necessary to manage, plan for, and respond to fire in the wildland areas of ORR and to reduce the risk of wildland fire to personnel and facilities on ORR and to the public. The WFMP is reviewed at least annually.

The WFMP applies to all DOE employees, contractors, and subcontractors working on ORR and to all DOE ORR tenant activities. The DOE ORR federal manager is responsible for ORR wildland fire management activities.

The primary goal of the WFMP is to lower the overall risk of wildland fire on ORR by conducting fire prevention activities and actions to reduce the spread of a fire should one start. Another goal of

the WFMP is to contain wildfires that do start to the ORR unit of origin by conducting suppression activities.

The WFMP is implemented by multiple organizations, including non-DOE entities such as the City of Oak Ridge and the Tennessee Department of Agriculture Division of Forestry. Memorandums of understanding that ensure collaboration among organizations are maintained for each organization that provides firefighting support on ORR.

DOE actions associated with wildland fire management include the following:


- Controlling ignition sources in the wildland areas, particularly on days when fire danger is forecasted
- Managing wildfire fuels in and near developed areas
- Developing and implementing controlled burning plans authorized by the DOE ORR federal manager
- Preparing and updating wildland fire preplans that include maps of fuel types, topographic features, roads, cultural resources, sensitive natural resources, contamination areas, and potential hazards
- Developing stakeholder involvement plans in support of the wildland fire program
- Reviewing current data to determine the potential for wildland fire, including indications of wildland fire risk
- Preparing a wildland fire risk report, including a wildland fire hazard severity analysis based on the *Standard for Wildland Fire Protection* (NFPA 2022)
- Maintaining a wildland fire road grid to support fire detection, containment, and suppression
- Conducting tabletop wildland fire exercises at least once every 3 years and full-scale exercises at least once every 5 years

- Incorporating wildland fire mitigation and response activities and procedures into the ORR land-use planning process

The DOE roads and grounds contractor is responsible for establishing and maintaining the wildland fire roads, many of which delineate wildland management units (Figure 6.7), and for maintaining barricades that control access to ORR secondary roads. The management contractors at ORNL and Y-12 are responsible for providing personnel and equipment for initial response to wildland fire events and for establishing incident command. The City of Oak Ridge at Heritage Center has entered into a mutual aid agreement with DOE to provide assistance for wildland fire activities. The Tennessee Division of Forestry has entered into a memorandum of understanding to provide trained personnel and heavy equipment, including fire plows, when requested to assist with wildland fires on ORR.

Because ORR is a large (32,258.54 acres), mostly forested property with access restrictions, it is a challenge for site emergency personnel to maintain familiarity with all remote areas and back roads and to recognize and assess concerns associated with those areas quickly. Wildland management unit prefire plans are designed to aid responders who may or may not be familiar with an area.

The prefire plans are concise documents for each of the 28 ORR wildland management units (Figure 6.7) that summarize access issues, assets, and hazard concerns. Each plan includes the wildland management unit's name and identification number, its general location within ORR, and its boundaries and size. Important information and hazard descriptions are listed early in the document, followed by guidance on tactics, access, vegetation and fuels, water sources, topographic considerations, and hazard controls. Plan maps depict access points, utilities, hazards, research areas, fuel types, water sources, urban interface areas, and sensitive resources. Prefire plans are reviewed on a 3-year cycle and are updated as significant changes occur.

**Acronyms:**

ETTP = East Tennessee Technology Park

ORNL = Oak Ridge National Laboratory

ORR = Oak Ridge Reservation

Y-12 = Y-12 National Security Complex

Figure 6.7. Wildland management units on ORR

Copies of the plans are kept in responder vehicles for immediate reference during remote events and are available to site fire departments and emergency operations centers, shift superintendent offices, and appropriate management staff. The plans are easily updated, stored, and shared electronically. They are meant to enable quick decisions but not to dictate tactics. The ORR forester is the point of contact for plan distribution.

The 2016 Great Smoky Mountains wildfires, also known as the Gatlinburg wildfires, demonstrated that large fires, although more frequent in western states, can occur on or near ORR. Issues related to

wildland and urban interface areas are a growing concern. These areas may be characterized by relatively high housing density and increasing recreational use by the public. DOE has prioritized interface areas and has conducted controlled wildfire fuel reduction burns to limit the spread of fire to and from the community. The presence of dense pine forests increases community vulnerability to potential high-intensity wildfires. Actions to protect these areas include thinning or replacing dense pine growth, mechanical treatments to thin younger pine proactively, and mulching heavy logging slash and insect-damaged timber to interrupt fuel beds.

During 2024, collaborative efforts were undertaken to reduce wildfire fuel hazards near developed areas at the ORR border. DOE partnered with the Tennessee Division of Forestry and one of its incident management teams to safely remove accumulations of woody storm debris and standing hazard trees and install additional firebreaks. The 2-week collaboration involved more than 50 state forestry personnel, 15 pieces of heavy equipment, and 40 support vehicles. The Tennessee Division of Forestry installed over 15 miles of firebreaks, removed 48 truckloads of wood debris, and performed 38 acres of fuel reduction. This effort also enabled Tennessee Division of Forestry staff to practice emergency preparedness and gain familiarity with ORR, where the division is a mutual-aid firefighting partner. These actions supplemented an ongoing series of controlled burns to proactively reduce fuel loads in the nearby forest.

6.9. Quality Assurance

UT-Battelle performs the activities associated with administration, sampling, data management, and reporting for ORR environmental surveillance programs. Project scope is established by a task team whose members represent DOE, UT-Battelle, Consolidated Nuclear Security LLC, and UCOR. UT-Battelle integrates quality assurance, environmental, and safety considerations into every aspect of ORR environmental monitoring. (See Chapter 5, Section 5.7, for a detailed discussion of UT-Battelle quality assurance program elements for environmental monitoring and surveillance activities.)

6.10. References

- Bowen et al. 2000. Bowen, B. M., J. A. Baars, and G. L. Stone, "Nocturnal Wind Shear and Its Potential Impact on Pollutant Transport." *Journal of Applied Meteorology* 39(3), 437–45.
- CERCLA 1980. *Comprehensive Environmental Response, Compensation, and Liability Act*. Title 42 US Code Chapter 103.
- Denton 2007. Denton, G. M., *Mercury Levels in Tennessee Fish*. Tennessee Department of Environment and Conservation, Nashville, Tennessee.
<https://www.tn.gov/content/dam/tn/environment/water/documents/fishmercurylevels.pdf>.
- DOE 2013. *Groundwater Strategy for the U.S. Department of Energy Oak Ridge Reservation, Oak Ridge, Tennessee*. Volume 1. Main Text. DOE/OR/01-2628/V1&D1, US Department of Energy Office of Environmental Management, Washington, DC. (PDF version available through online catalog at <http://doeic.science.energy.gov/>.)
- DOE 2015. *Environmental Radiological Effluent Monitoring and Environmental Surveillance*. DOE-HDBK-1216-2015, US Department of Energy, Washington, DC. Retrieved May 18, 2021, from <https://www.standards.doe.gov/standards-documents/1200/1216-bhdbk-2015>.
- DOE 2017. *Offsite Groundwater Assessment Remedial Site Evaluation, Oak Ridge, Tennessee*. DOE/OR/01-2715&D2_R, US Department of Energy Office of Environmental Management, Washington, DC.
- DOE 2019. *Facility Safety*. DOE Order 420.1C, US Department of Energy, Washington, DC. Retrieved June 15, 2021, from <https://www.directives.doe.gov/directives-documents/400-series/0420.1-BOrder-C-chg3-ltdchg>.
- DOE 2020. *Radiation Protection of the Public and the Environment*. DOE Order 458.1, US Department of Energy, Washington, DC.
- DOE 2021a. *Derived Concentration Technical Standard*. DOE-STD-1196-2021, US Department of Energy, Washington, DC. Retrieved March 28, 2022, from <https://www.standards.doe.gov/standards-documents/1100/1196-astd-2021/@@images/file>.

- DOE 2021b. *Environmental Monitoring Plan for the Oak Ridge Reservation, Calendar Year 2021*. DOE/ORO-228/R12, US Department of Energy Office of Science, Washington, DC. Retrieved April 11, 2021, from https://doeic.science.energy.gov/ASER/ORR_EMP_CY2021_Final.pdf.
- DOE 2021c. *U.S. Department of Energy Conservation Action Plan: Report to the White House Council on Environmental Quality*. US Department of Energy, Washington, DC.
- DOE 2023. *Fire Protection*. DOE-STD-1066-2023, US Department of Energy, Washington, DC. <https://www.standards.doe.gov/standards-documents/1000/1066-astd-2023/@images/file>.
- DOE 2024a. *Melton Valley/Bethel Valley Exit Pathway Phase 1 Remedial Investigation Completion Report for the U.S. Department of Energy Oak Ridge Site Oak Ridge, Tennessee*. DOE/OR/01-2953&D2, US Department of Energy, Oak Ridge Office of Environmental Management, Oak Ridge, Tennessee.
- DOE 2024b. *Oak Ridge Reservation Wildland Fire Management Plan*. US Department of Energy Office of Science, Washington, DC.
- DOE 2024c. *Oak Ridge Reservation Wildland Fire Implementation Plan*. US Department of Energy Office of Science, Washington, DC.
- DOE 2025. *2025 Remediation Effectiveness Report for the U.S. Department of Energy Oak Ridge Site Oak Ridge, Tennessee: Data and Evaluations*. DOE/OR/01-2989&D1, Prepared by the Water Resources Restoration Program, UCOR for the US Department of Energy Office of Environmental Management, Oak Ridge, Tennessee.
- DOI 2020. *Early Detection and Rapid Response*. National Invasive Species Council, US Department of the Interior, Washington, DC. Retrieved April 20, 2021, from <https://www.doi.gov/invasivespecies/early-detection-and-rapid-response>.
- Drake et al. 2002. Drake, S. J., J. F. Weltzin, and P. D. Parr, *Assessment of Nonnative Invasive Plants in the DOE Oak Ridge National Environmental Research Park*. ORNL/TM-2001/113, Oak Ridge National Laboratory, Oak Ridge, Tennessee.
- Federal Noxious Weed Act of 1974*. Public Law 93-629 (7 U.S.C. 2801 et seq.; 88 Stat. 2148).
- Federal Plant Protection Act of 2000*. Public Law 106-224, Title IV (7 U.S.C. 7701 et seq). <https://www.govinfo.gov/content/pkg/STATUTE-114/pdf/STATUTE-114-Pg358.pdf#page=1>.
- McCracken and Giffen 2017. McCracken, M. K., and N. R. Giffen, *Invasive Plant Management Plan for the Oak Ridge Reservation*. ORNL/TM-2004/98/R2, Oak Ridge National Laboratory, Oak Ridge, Tennessee.
- NFPA 2022. *Standard for Wildland Fire Protection*. NFPA 1140, National Fire Protection Association, Quincy, Massachusetts. <https://www.nfpa.org/codes-and-standards/all-codes-and-standards/list-of-codes-and-standards/detail?code=1140>.
- Parr et al. 2004. Parr, P. D., M. G. Ryon, H. D. Quarles, N. R. Giffen, M. S. Salk, and J. W. Webb, *Invasive Plant Management Plan for the Oak Ridge Reservation*. ORNL/TM-2004/98, Oak Ridge National Laboratory, Oak Ridge, Tennessee.
- Quarles et al. 2011. Quarles, H. D., M. S. Salk, P. D. Parr, M. G. Ryon, and N. R. Giffen, *Invasive Plant Management Plan for the Oak Ridge Reservation*. ORNL/TM-2004/98/R1, Oak Ridge National Laboratory, Oak Ridge, Tennessee.
- TDEC 2023. *Posted Streams, Rivers, and Reservoirs in Tennessee*. Tennessee Department of Environment and Conservation, Division of Water Resources, Nashville, Tennessee. https://www.tn.gov/content/dam/tn/environment/water/watershed-planning/wr_wq_fish-advisories.pdf.

TDEC 2024a. "Use Classifications for Surface Waters." *Rules of the Tennessee Department of Environment and Conservation*. Chapter 0400-40-04. Tennessee Department of Environment and Conservation, Bureau of Environment, Division of Water Supply.
<https://publications.tnsosfiles.com/rules/0400/0400-40/0400-40-04.20240317.pdf>.

Rules of the Tennessee Department of Environment and Conservation. Chapter 0400-40-03. Tennessee Department of Environment and Conservation, Bureau of Environment, Division of Water Supply.
<https://publications.tnsosfiles.com/rules/0400/0400-40/0400-40-03.20240317.pdf>.