

Appendix F: Chemicals

Appendix

F. Chemicals

This appendix presents basic information about chemicals. The information is intended as a basis for understanding the dose or relative toxicity assessment associated with possible releases from the Oak Ridge Reservation (ORR), and is not a comprehensive discussion of chemicals and their effects on the environment and biological systems.

F.1 Perspective on Chemicals

The lives of modern humans have been greatly improved by the development of chemicals such as pharmaceuticals, building materials, housewares, pesticides, and industrial chemicals. Through the use of chemicals we can increase food production, cure diseases, build more efficient houses, and send people to the moon. At the same time, we must be cautious to ensure that our own existence is not endangered by uncontrolled and over-expanded use of chemicals (Chan et al. 1982).

Just as all humans are exposed to radiation in their normal daily routines, humans are also exposed to chemicals. Some potentially hazardous chemicals exist in the natural environment. In many areas of the country, soils contain naturally elevated concentrations of metals such as selenium, arsenic, or molybdenum, which may be hazardous to humans or animals. Even some of the foods we eat contain natural toxins. Aflatoxins are found in chili peppers, corn, millet, peanuts, rice, sorghum, sunflower seeds, tree nuts, and wheat. Cyanide is found in apple seeds. However, exposure to many more hazardous chemicals result from the direct or indirect actions of humans. Building materials used in the construction of homes may contain chemicals such as formaldehyde (in some insulation materials), asbestos (formerly used in insulations and ceiling tiles), and lead (formerly used in paints and gasoline). Some chemicals are present as a result of the application of pesticides and fertilizers to soil. Other chemicals may have been transported long distances through the atmosphere from industrial sources before being deposited on soil or water.

F.2 Pathways of Chemicals from Oak Ridge Reservation to the Public

“Pathways” refers to the route or way in which a person can come into contact with a chemical substance. Chemicals released to the air may remain suspended for long periods, or they may be rapidly deposited on plants, soil, and water. Chemicals may also be released as liquid wastes, called “effluents,” which can enter streams and rivers.

People are exposed to chemicals by inhalation (breathing air), ingestion (eating exposed plants and animals or drinking water), or direct contact (touching the soil or swimming in water). For example, fish that live in a river that receives effluents may take in some of the chemicals present in the water. People eating the fish and drinking water from the river would then be exposed to the chemicals. The public is not normally exposed to chemicals on ORR because access to the reservation is limited. However, chemicals released as a result of ORR operations can move through the environment to off-site locations, resulting in potential exposure of the public.

F.3 Definitions

F.3.1 Toxicity

Chemicals have varying types of effects. Chemical health effects are divided into two broad categories: adverse or systemic effects (noncarcinogens) and cancer (carcinogens). Sometimes a chemical can have both noncarcinogenic and carcinogenic effects. The toxic effect can be acute (a short-term, severe health effect) or chronic (a longer term, persistent health effect). Noncarcinogenic toxicity is often evident in a shorter length of time than a carcinogenic effect. The potential health effects of noncarcinogens range from skin irritation to death (or mortality). Carcinogens cause or increase the incidence of malignant neoplasms or cancers.

Toxicity refers to an adverse effect of a chemical on human health. Every day we ingest chemicals in food, water, and sometimes medications. Even those chemicals typically considered toxic are usually nontoxic or harmless below a certain concentration.

Concentration limits or advisories are set by government agencies for some chemicals that are known or thought to have adverse effects on human health. These concentration limits can be used to calculate chemical doses that would not harm even those individuals who are particularly sensitive to the chemical.

F.3.2 Dose Terms for Noncarcinogens

F.3.2.1 Reference Dose

A reference dose is an estimate of a daily exposure level for the human population, including sensitive subpopulations. These reference doses are likely to be without appreciable risk of deleterious effects during a lifetime. Units are expressed as milligrams of chemical per kilogram of an adult's body weight per day (mg/kg-day). Values for reference doses are derived from doses of chemicals that resulted in no adverse effect, or the lowest dose that showed an adverse effect on humans or laboratory animals.

Uncertainty factors are typically used in deriving reference doses. Uncertainty adjustments may be made if animal toxicity data are extrapolated to humans, to account for human sensitivity; extrapolated from subchronic to chronic no-observed-adverse-effect levels; extrapolated from lowest-observed-adverse-effect levels to no-observed-adverse-effect levels; and to account for database deficiencies. The use of uncertainty factors in deriving reference doses is thought to protect sensitive human populations. The US Environmental Protection Agency (EPA) maintains the Integrated Risk Information System (IRIS) database, which contains verified reference doses and up-to-date health risk and EPA regulatory information for numerous chemicals.

F.3.2.2 Primary Maximum Contaminant Levels

For chemicals for which reference doses are not available in IRIS, Tennessee Water Quality Criteria, which reflect maximum contaminant levels expressed in milligrams of chemical per liter of drinking water, are converted to reference dose values by multiplying by 2 L (the average daily adult water intake) and dividing by 80 kg (the reference adult body weight). The result is a "derived" reference dose expressed in milligrams per kilogram per day (mg/kg-day).

F.3.3 Dose Term for Carcinogens

F.3.3.1 Slope Factor

A slope factor is a plausible upper-bound estimate of the probability of a response per unit intake of a chemical during a lifetime. The slope factor is used to estimate an upper-bound probability of an individual developing cancer as a result of a lifetime exposure to a particular level of a chemical. Units are expressed as risk per dose (mg/kg-day).

The slope factor converts the estimated daily intake averaged over a lifetime exposure to the incremental risk of an individual developing cancer. Because it is unknown for most chemicals whether a threshold (a dose below which no adverse effect occurs) exists for carcinogens, units for carcinogens are set in terms of risk factors. Acceptable risk levels for carcinogens range from 10^{-4} (risk of developing cancer over a human lifetime is 1 in 10,000) to 10^{-6} (risk of developing cancer over a human lifetime is 1 in 1,000,000). In other words, a certain chemical concentration in food or water could cause a risk of one additional cancer for every 10,000 (10^{-4}) to 1,000,000 (10^{-6}) exposed persons, respectively.

F.4 Measuring Chemicals

Environmental samples are collected in areas surrounding ORR and are analyzed for those chemical constituents most likely to be released from ORR. Typically, chemical concentrations in liquids are expressed in terms of milligrams or micrograms of chemical per liter of water; concentrations in solids (soil and fish tissue) are expressed in terms of milligrams or micrograms of chemical per gram or kilogram of sample material.

The instruments used to measure chemical concentrations are sensitive; however, there are limits below which they cannot detect chemicals of interest. Concentrations detected below the reported analytical detection limits of the instruments are recorded by the laboratory as estimated values, which have a greater uncertainty than concentrations detected above the detection limits of the instruments. Health effect calculations that use these estimated values are indicated by the less-than symbol (<), which indicates that the value for a parameter was not quantifiable at the analytical detection limit.

F.5 Risk Assessment Methodology

F.5.1 Exposure Assessment

To evaluate an individual's exposure by way of a specific exposure pathway, the intake amount of the chemical must be determined. For example, chemical exposure by drinking water and eating fish from the Clinch River is assessed in the following manner: Clinch River surface water and fish samples are analyzed to estimate chemical contaminant concentrations. It is assumed that individuals drink about 2 L (0.5 gal) of water per day directly from the river, which amounts to 730 L (193 gal) per year, and that they eat 0.07 kg (roughly 0.2 lb) of fish per day from the river (27 kg or 60 lb per year). Estimated daily intakes or estimated doses to the public are calculated by multiplying measured (statistically significant) concentrations in water by 2 L, or those in fish by 0.07 kg. This intake is first multiplied by the exposure duration (26 years) and exposure frequency (350 days/year) and then divided by an averaging time (26 years for noncarcinogens and 70 years for carcinogens) and 80 kg body weight. These assumptions are conservative, and in many cases they result in higher estimated intakes and doses than an individual would actually receive.

F.5.2 Dose Estimate

When the contaminant oral daily intake has been estimated, the dose is determined. For chemicals, the dose to humans is measured as milligrams per kilogram-day (mg/kg-day). In this case, the “kilogram” refers to the body weight of an adult. When a chemical dose is calculated, the length of time an individual is exposed to a certain concentration is important. To assess off-site doses, it is assumed that the exposure duration occurs over 30 years. Such exposures are called “chronic” in contrast to short-term exposures, which are called “acute.”

F.5.3 Calculation Method

Current risk assessment methodologies use the term “hazard quotient” to evaluate noncarcinogenic health effects. Because intakes are calculated in milligrams per kilogram per day in the hazard quotient methodology, they are expressed in terms of dose. Hazard quotient values of less than 1 indicate an unlikely potential for adverse health effects, whereas hazard quotient values greater than 1 indicate a concern for adverse health effects or the need for further study.

To evaluate carcinogenic risk, slope factors are used instead of reference doses.

To estimate the risk of inducing cancers from ingestion of water and fish, the estimated dose or intake (I) is multiplied by the slope factor (risk per mg/kg-day). As mentioned earlier, acceptable risk levels for carcinogens range from 10^{-4} (risk of developing cancer over a human lifetime is 1 in 10,000) to 10^{-6} (risk of developing cancer over a human lifetime is 1 in 1,000,000).

F.6 References

- Chan, P.K., G.P. O’Hara, and A.W. Hayes, 1982. “Principles and Methods for Acute and Subchronic Toxicity.” *Principles and Methods of Toxicology*. Raven Press, New York.
- Memorandum: Human Health Evaluation Manual, Supplemental Guidance: Update of Standard Default Exposure Factors, OSWER Directive 9200.1-120, U.S Environmental Protection Agency, February 6, 2014.
- TDEC 2008. “General Water Quality Criteria.” Chapter 1200-4-3 in *Rules of Tennessee Department of Environment and Conservation, Tennessee Water Quality Control Board, Division of Water Pollution Control*. June.