

Appendix E: Radiation

Appendix

E. Radiation

This appendix presents basic information about radiation. The information is intended as a basis for understanding the potential doses associated with releases of radionuclides from the Oak Ridge Reservation (ORR), not as a comprehensive discussion of radiation and its effects on the environment and on biological systems.

Radiation comes from natural and human sources. People are constantly exposed to naturally occurring radiation. For example, cosmic radiation, radon in air, potassium in food and water, and uranium, thorium, and radium in the earth's crust are all sources of radiation. The following discussion describes important aspects of radiation and its types, sources, and pathways; radiation measurement; and dose information.

E.1 Atoms and Isotopes

All matter is made up of atoms. An atom is “a unit of matter consisting of a single nucleus surrounded by a number of electrons equal to the number of protons in the nucleus” (Alter 1986). The number of protons in the nucleus determines an element's atomic number or chemical identity. With the exception of hydrogen, the nucleus of each type of atom also contains at least one neutron. Unlike protons, the neutrons may vary in number among atoms of the same element. The number of neutrons and protons determines the atomic weight. Atoms of the same element that have different numbers of neutrons are called isotopes. In other words, isotopes have the same chemical properties but different atomic weights (see Figure E.1).




Figure E.1. The hydrogen atom and its isotopes

For example, the element uranium has 92 protons. All isotopes of uranium, therefore, have 92 protons. However, each uranium isotope has a different number of neutrons:

- uranium-238 has 92 protons and 146 neutrons
- uranium-235 has 92 protons and 143 neutrons
- uranium-234 has 92 protons and 142 neutrons

Some isotopes are stable, or nonradioactive; some are radioactive. Radioactive isotopes are called radionuclides or radioisotopes. In an attempt to become stable, radionuclides emit rays or particles. This

emission of rays and particles is known as radioactive decay. Each radioisotope has a radioactive half-life, which is the average time required for half of a specified number of atoms to decay. Half-lives can be very short (fractions of a second) or very long (millions of years), depending on the isotope. Table E.1 shows the half-lives of selected radionuclides.

Table E.1. Selected radionuclide half-lives

Radionuclide	Symbol	Half-life (in years unless otherwise noted)	Radionuclide	Symbol	Half-life (in years unless otherwise noted)
Americium-241	²⁴¹ Am	432.2	Plutonium-238	²³⁸ Pu	87.74
Americium-243	²⁴³ Am	7.37E+3	Plutonium-239	²³⁹ Pu	2.411E+4
Argon-41	⁴¹ Ar	1.827 hours	Plutonium-240	²⁴⁰ Pu	6.564E+3
Beryllium-7	⁷ Be	53.22 days	Potassium-40	⁴⁰ K	1.251E+9
Californium-252	²⁵² Cf	2.645	Radium-226	²²⁶ Ra	1.6E+3
Carbon-11	¹¹ C	20.39 minutes	Radium-228	²²⁸ Ra	5.75
Carbon-14	¹⁴ C	5.70E+3	Ruthenium-103	¹⁰³ Ru	39.26 days
Cerium-141	¹⁴¹ Ce	32.508 days	Samarium-153	¹⁵³ Sm	46.5 hours
Cerium-144	¹⁴⁴ Ce	284.91 days	Strontium-89	⁸⁹ Sr	50.53 days
Cesium-134	¹³⁴ Cs	2.0648	Strontium-90	⁹⁰ Sr	28.79
Cesium-137	¹³⁷ Cs	30.167	Techneium-99	⁹⁹ Tc	2.111E+5
Cesium-138	¹³⁸ Cs	32.41 minutes	Thorium-228	²²⁸ Th	1.9116
Cobalt-58	⁵⁸ Co	70.86 days	Thorium-230	²³⁰ Th	7.538E+4
Cobalt-60	⁶⁰ Co	5.271	Thorium-232	²³² Th	1.405E+10
Curium-242	²⁴² Cm	162.8 days	Thorium-234	²³⁴ Th	24.1 days
Curium-244	²⁴⁴ Cm	18.1	Tritium	³ H	12.32
Iodine-129	¹²⁹ I	157E+7	Uranium-234	²³⁴ U	2.455E+5
Iodine-131	¹³¹ I	8.02 days	Uranium-235	²³⁵ U	7.04E+8
Krypton-85	⁸⁵ Kr	10.756	Uranium-236	²³⁶ U	2.342E+7
Krypton-88	⁸⁸ Kr	2.84 hours	Uranium-238	²³⁸ U	4.468E+9
Lead-212	²¹² Pb	10.64 hours	Xenon-133	¹³³ Xe	5.243 days
Manganese-54	⁵⁴ Mn	312.12 days	Xenon-135	¹³⁵ Xe	9.14 hours
Neptunium-237	²³⁷ Np	2.144E+6	Yttrium-90	⁹⁰ Y	64.1 hours
Niobium-95	⁹⁵ Nb	34.991 days	Zirconium-95	⁹⁵ Zr	64.032 days

Source: ICRP 2008

E.2 Radiation

Radiation, or radiant energy, is energy in the form of waves or particles moving through space. Visible light, heat, radio waves, and alpha particles are examples of radiation. When people feel warmth from sunlight, they are actually absorbing the radiant energy emitted by the sun.

Electromagnetic radiation is radiation in the form of electromagnetic waves. Examples include gamma rays, ultraviolet light, and radio waves. Particulate radiation is radiation in the form of particles. Examples include alpha and beta particles. Radiation also is characterized as ionizing or nonionizing because of the way in which it interacts with matter.

E.2.1 Ionizing Radiation

Normally an atom has an equal number of protons and electrons; however, atoms can lose or gain electrons in a process known as ionization. Some forms of radiation (called ionizing radiation) can ionize atoms by knocking electrons off atoms. Examples of ionizing radiation include alpha and beta particles and gamma and x-rays.

Ionizing radiation is capable of changing the chemical state of matter and subsequently causing biological damage. By this mechanism, it is potentially harmful to human health.

E.2.2 Nonionizing Radiation

Nonionizing radiation is described as a series of energy waves composed of oscillating electric and magnetic fields traveling at the speed of light. Nonionizing radiation includes the spectrum of ultraviolet, visible light, infrared, microwave, radio frequency, and extremely low frequency. Lasers commonly operate in the ultraviolet, visible, and infrared frequencies. Microwave radiation is absorbed near the skin, while radio frequency radiation may be absorbed throughout the body. At high enough intensities, both will damage tissue through heating. Excessive visible radiation can damage the eyes and skin (Department of Labor, OSHA *Safety and Health Topics*). However, in the discussion that follows, the term “radiation” is used to describe ionizing radiation.

E.3 Measuring Ionizing Radiation

To determine the possible effects of radiation on the health of the environment and the public, the radiation must be measured. More precisely, its potential to cause damage must be ascertained.

E.3.1 Activity

To determine radiation in the environment, the rate of radioactive decay or activity is measured. The rate of decay varies widely among various radioisotopes. For that reason, 1 gram of a radioactive substance may contain the same amount of activity as several tons of another material. This activity is expressed in a unit of measure known as a curie (Ci). More specifically, 1 Ci equals 3.7×10^{10} (37,000,000,000) atomic disintegrations per second (dps). In the International System of Units, 1 dps equals 1 becquerel (Bq).

E.3.2 Absorbed Dose

The total amount of energy absorbed per unit mass of the exposed material as a result of exposure to radiation is expressed in a unit of measure known as a rad. The effect of the absorbed energy (the biological damage that occurs) is important, not the actual amount. In the International System of Units, 100 rad equals 1 gray (Gy).

E.3.3 Effective Dose

The measure of potential biological damage to the body caused by exposure to and subsequent absorption of radiation is expressed in a unit of measure known as a rem. For radiation protection purposes, 1 rem of any type of radiation has the same damaging effect. Because a rem represents a fairly large dose, the measure is usually expressed as millirem (mrem), which is 1/1000 of a rem. In the International System of Units, 1 sievert (Sv) equals 100 rem; 1 millisievert (mSv) equals 100 mrem. The effective dose (ED) is the weighted sum of equivalent dose over specified tissues or organs. The ED is based on tissue-weighting factors for 12 specific tissues or organs plus a weight factor for the remaining organs and

tissues. In addition, the ED is based on the recent lung model, gastrointestinal absorption fractions, and biokinetic models used for selected elements. Specific types of EDs are defined as follows:

- Committed ED – the weighted sum of the committed ED in specified tissues in the human body during the 50-year period following intake
- Collective ED – the product of the mean ED for a population and the number of persons in the population

E.4 Radiation Exposure Pathways

People can be exposed to radionuclides in the environment through a number of routes, as shown in Figure E.2. Potential routes for internal and external exposure are referred to as pathways. For example, radionuclides in air could be inhaled directly or could fall on grass in a pasture. If the grass were then consumed by cows, it would be possible for the radionuclides to impact the cow's milk, and people drinking the milk would be exposed to this radiation. Similarly, radionuclides in water could be ingested by fish, and fishermen or other consumers could then ingest the radionuclides in the fish tissue. People swimming in the water also would be exposed. Exposure to ionizing radiation varies significantly with geographic location, diet, drinking water source, and building construction.



Figure E.2. Examples of radiation pathways

E.5 Radiation Sources and Doses

Basically, radioactive decay, or activity, generates radiant energy. People absorb some of the energy to which they are exposed, either from external or internal radiation. The effect of this absorbed energy is responsible for an individual's dose. Whether radiation is natural or human-made, it has the same effect on people.

There are five broad categories for radiation exposure to the US population (NCRP 2009):

- exposure to ubiquitous background radiation, including radon in homes
- exposure to patients from medical procedures
- exposure from consumer products or activities involving radiation sources
- exposure from industrial, security, medical, educational, and research radiation sources
- exposure to workers that results from their occupations

Figure E.3 gives the percent contributions of various sources of exposure to the total collective dose for the US population in 2006. As shown, the major sources are radon and thoron (37 percent), computed tomography (24 percent), and nuclear medicine (12 percent) (NCRP 2009). Consumer, occupational, and industrial sources contribute about 2 percent to the total US collective dose.




Figure E.3. All exposure categories for collective effective dose for 2006 (NCRP 2009)

E.5.1 Background Radiation

Naturally occurring radiation is the major source of radiation in the environment. Sources of background radiation exposure include the following:

- external exposure from space or cosmic radiation
- external exposure from terrestrial radiation
- internal exposure from inhalation of radon, thoron, and their progeny
- internal exposure from radionuclides in the body

E.5.1.1 External Exposures

Space or Cosmic Radiation

Energetically charged particles from outer space continuously hit the earth's atmosphere. These particles and the secondary particles and photons they create are called cosmic radiation. Because the atmosphere provides some shielding against cosmic radiation, the intensity of this radiation increases with altitude above sea level. For example, a person in Denver is exposed to more cosmic radiation than a person in New Orleans.

The average annual effective dose to people in the United States from cosmic radiation is about 33 mrem, or 0.33 mSv (NCRP 2009). Effective dose rates from cosmic radiation depend on geomagnetic latitude and elevation above sea level.

Terrestrial Radiation

Terrestrial radiation refers to radiation emitted from radioactive materials in the earth's rocks, soils, and minerals. Radon (Rn), radon progeny (the relatively short-lived decay products from the decay of the radon isotope ^{222}Rn), potassium (^{40}K), isotopes of thorium (Th), and isotopes of uranium (U) are the elements responsible for most terrestrial radiation.

The average annual dose from terrestrial gamma radiation is about 21 mrem (0.21 mSv) in the United States, but it varies geographically across the country (NCRP 2009). Typical reported values are about 23 mrem (0.23 mSv) on the Atlantic and Gulf coastal plains, about 90 mrem (0.9 mSv) on the eastern slopes of the Rocky Mountains, and about 46 mrem (0.46 mSv) elsewhere (EPA 2014).

E.5.1.2 Internal Exposures

Radionuclides in the environment enter the body with the air people breathe and the foods they eat. They also can enter through an open wound. Natural radionuclides that can be inhaled and ingested include isotopes of uranium and its progeny, especially radon (^{222}Rn) and its progeny, thoron (^{220}Rn) and its progeny, potassium (^{40}K), rubidium (^{87}Rb), and carbon (^{14}C). Radionuclides contained in the body are dominated by ^{40}K and polonium (^{210}Po); others include ^{87}Rb and ^{14}C (NCRP 1987).

Radon and Thoron and Decay Products

The major contributors to the annual effective dose from background radiation sources are radon and thoron and their short-lived decay products. As shown in Figure E.3, 37 percent of the dose from all exposure categories is from radon and thoron and their decay products, which contribute an average dose of about 228 mrem (2.28 mSv) per year (NCRP 2009). Radon is an inert gas and a small fraction is retained in the body; however, the dose to the lung comes from the short-lived radon decay products. Radon levels vary widely across the United States. Elevated levels are most commonly found in the Appalachians, the upper Midwest, and the Rocky Mountain states (NCRP 2009).

Other Internal Radiation Sources

Other sources of internal radiation include ^{40}K and ^{232}Th and the ^{238}U series. The primary source of ^{40}K in body tissues is food, primarily fruits and vegetables. The sources of radionuclides from ^{232}Th and ^{238}U series are food and water (NCRP 2009). The average dose from these other internal radionuclides is about 29 mrem (0.29 mSv) per year. This dose is attributed predominantly to the naturally occurring radioactive isotope of potassium, ^{40}K .

E.5.2 Human-Made Radiation

In addition to background radiation, there are human-made sources of radiation to which most people are exposed. Examples include consumer products, medical sources, industrial by-products, and fallout from atmospheric atomic bomb tests. No atmospheric testing of atomic weapons has occurred since 1980 (NCRP 1987).

E.5.2.1 Consumer Products

Some consumer products are sources of radiation. The radiation in these products, such as smoke detectors, radioluminous products (e.g., self-illuminating exit signs in commercial buildings), and airport x-ray baggage inspection systems, is essential to the performance of the device. In other products, such as tobacco products and building materials, the radiation occurs incidentally to the product's function (NCRP 1987, NCRP 2009).

The US average annual dose to an individual from consumer products and activities is about 13 mrem (0.13 mSv), ranging between 0.1 and 40 mrem (0.001 and 0.4 mSv). Cigarette smoking accounts for about 35 percent of this dose. Other important sources are building materials (27 percent), commercial air travel (26 percent), mining and agriculture (6 percent), miscellaneous consumer-oriented products (3 percent), combustion of fossil fuels (2 percent), highway and road construction materials (0.6 percent), and glass and ceramics (less than 0.003 percent). Television and video, sewage sludge and ash, and self-illuminating signs all contribute negligible doses (NCRP 2009).

E.5.2.2 Medical Sources

Radiation is an important tool of diagnostic medicine and treatment, which are the main sources of exposure to the public from human-made radiation. Exposure is deliberate and is directly beneficial to the patients exposed. In general, medical exposures from diagnostic or therapeutic x-rays result from beams directed to specific areas of the body. Thus, not all body organs are uniformly irradiated. Nuclear medicine examinations and treatments involve the internal administration of radioactive compounds, or radiopharmaceuticals, by injection, inhalation, consumption, or insertion. Even then, radionuclides are not distributed uniformly throughout the body. Radiation and radioactive materials also are used in preparing medical instruments, including sterilizing heat-sensitive products such as plastic heart valves.

Nuclear medicine examinations, which involve internal administration of radiopharmaceuticals, generally account for the largest portion of dose from human-made sources. However, the radionuclides used for specific tests are not uniformly distributed throughout the body. In these cases the concept of ED, which relates the significance of exposures of organs or body parts to the effect on the entire body, is useful in making comparisons. The average annual ED from medical examinations is roughly 300 mrem (3 mSv), including 147 mrem (1.47 mSv) from computed tomography scans, 77 mrem (0.77 mSv) from nuclear medicine procedures, 43 mrem (0.43 mSv) from interventional fluoroscopy, and 33 mrem (0.33 mSv) from conventional radiography and fluoroscopy (NCRP 2009). Not everyone receives such exams each year.

E.5.2.3 Other Sources

Other sources of radiation include emissions of radioactive materials from nuclear facilities such as uranium mines, fuel-processing plants, and nuclear power plants; transportation of radioactive materials; and emissions from mineral-extraction facilities. The dose to the general public from nuclear fuel cycle facilities, such as uranium mines, mills, fuel-processing plants, nuclear power plants, and transportation routes, has been estimated at less than 1 mrem (0.01 mSv) per year (NCRP 1987).

Small doses to individuals occur because of radioactive fallout from atmospheric atomic bomb tests, emissions of radioactive materials from nuclear facilities, emissions from certain mineral extraction facilities, and transportation of radioactive materials. The combination of these sources contributes less than 1 mrem (0.01 mSv) per year to an individual's average dose (NCRP 1987).

E.6 References

- Alter, H. 1986. *A Glossary of Terms in Nuclear Science and Technology*. American Nuclear Society, La Grange Park, Illinois.
- Department of Labor. *OSHA Safety and Health Topics, Non Ionizing Radiation*. Last accessed July 6, 2018.
- EPA. *Calculate Your Radiation Dose*. US Environmental Protection Agency. Last accessed July 6, 2018.
- ICRP 2008. *Nuclear Decay Data for Dosimetric Calculations*. ICRP Publication 107. *Annals of the ICRP* 38(3). International Commission on Radiological Protection.
- NCRP 1987. *Ionizing Radiation Exposure of the Population of the United States*. NCRP Report No. 93. National Council on Radiation Protection and Measurements, Washington, DC.
- NCRP 2009. *Ionizing Radiation Exposure of the Population of the United States*. NCRP Report No. 160, National Council on Radiation Protection and Measurements, Bethesda, Maryland.