
## **Bear Creek Valley Mercury Sources Remedial Site Evaluation** for the U.S. Department of Energy Oak Ridge Site Oak Ridge, Tennessee



DOCUMENT IS UNCLASSIFIED, NON-SENSITIVE APPROVED FOR PUBLIC RELEASE

Name/Organization: Dave Lannom

Date: 06/24/2025 eIRO#: 8183




#### DOE/OR/01-2977&D2

# Bear Creek Valley Mercury Sources Remedial Site Evaluation for the U.S. Department of Energy Oak Ridge Site Oak Ridge, Tennessee

Date Issued—August 2025

Prepared for the U.S. Department of Energy Office of Environmental Management

United Cleanup Oak Ridge LLC under contract 89303322DEM000067



### **CONTENTS**

FIG	URES		v
TAI	BLES .		vii
ACl	RONY	MS	ix
EXI	ECUTI	VE SUMMARY	ES-1
1.	INTR	ODUCTION	1-1
	1.1	REMEDIAL SITE EVALUATION OBJECTIVE	1-1
	1.2	BACKGROUND	1-1
		1.2.1 Site Description	1-1
		1.2.2 Summary of Potential Mercury Source Areas	1-8
		1.2.3 Summary of Historical Bear Creek Valley Mercury Data	1-9
	1.3	REPORT CONTENT	1-11
2.		EMENTATION OF THE BEAR CREEK VALLEY MERCURY SOURCES	
	REM	EDIAL SITE EVALUATION SAMPLING AND ANALYSIS PLAN	
	2.1	SCOPE OF WORK	2-1
		2.1.1 Channel Sediment Sampling	2-6
		2.1.2 Creek Bank Soil Sampling	2-6
		2.1.3 Floodplain Soil Sampling	2-7
		2.1.4 Surface Water Sampling	2-7
		2.1.5 Speciation of Mercury via Sequential Extraction	2-7
	2.2	CHANGES FROM THE BEAR CREEK VALLEY MERCURY SOURCES	
		REMEDIAL SITE EVALUATION SAMPLING AND ANALYSIS PLAN	2-8
3.		R CREEK VALLEY MERCURY SOURCES REMEDIAL SITE EVALUATION	2.1
		JLTS	
	3.1	SURFACE WATER DATA	
	3.2	FLOODPLAIN SOIL, CREEK BANK SOIL, AND CHANNEL SEDIMENT DATA	
		3.2.1 Soil and Sediment Particle Size	
		3.2.2 Soil and Sediment Chemical Characteristics	3-13
	3.3	BEAR CREEK VALLEY SOIL AND SEDIMENT MERCURY SEQUENTIAL EXTRACTIONS	2 25
	2.4	3.3.1 Analyte Correlation Evaluations	3-41
	3.4	BEAR CREEK VALLEY MERCURY COMPARISON TO LOWER EAST FORK POPLAR CREEK	3_41
4.	WAT	ER RESOURCES RESTORATION PROGRAM BEAR CREEK MONITORING	5-41
••		JLTS	4-1
	4.1	SURFACE WATER DATA	4-1
	4.2	BIOLOGICAL DATA	4-7
5.	CON	CLUSIONS AND RECOMMENDATIONS	
	5.1	CONCLUSIONS	5-1
	5.2	RECOMMENDATIONS	5-2

6.	REFERENCES	6-1
AP	PENDIX A. D2 VERSION OF THE BEAR CREEK VALLEY MERCURY SOURCES REMEDI	AL
	SITE EVALUATION SAMPLING AND ANALYSIS PLAN, OAK RIDGE,	
	TENNESSEE (DOE/OR/01-2958&D2) THAT INCLUDES TENNESSEE	
	DEPARTMENT OF ENVIRONMENT AND CONSERVATION AND U.S.	
	ENVIRONMENTAL PROTECTION AGENCY COMMENTS ON THE D1	
	VERSION OF THE DOCUMENT	.A-1
AP	PENDIX B. PHOTOGRAPHS OF TRANSECT LOCATIONS	B-1
AP	PENDIX C. ANALYTICAL DATA SUMMARY TABLES BY MEDIA	.C-1
AP	PENDIX D. PARTICLE SIZE ANALYSIS	.D-1
AP	PENDIX E. BEAR CREEK VALLEY REMEDIAL SITE EVALUATION ANALYTE	
	CORRELATIONS	.E-1
AP	PENDIX F. WATER RESOURCES RESTORATION PROGRAM SURFACE WATER TOTAL	
	MERCURY AND METHYLMERCURY LONGITUDINAL DATA PLOTS FOR	
	BEAR CREEK	. F-1
AP]	PENDIX G. BEAR CREEK FIELD OBSERVATIONS	.G-1

## **FIGURES**

Figure 1.1. BCV within the Oak Ridge Site.	1-2
Figure 1.2. BCV Watershed and historic monitoring locations.	1-3
Figure 1.3. BCV hydrogeologic CSM for Upper Bear Creek	1-6
Figure 1.4. Surface water data for mercury and methylmercury in Bear Creek, NT-3, and Creek, 2011–2022.	
Figure 2.1. BCV transect locations.	2-3
Figure 2.2. BCV RSE transect conceptual diagram.	2-5
Figure 3.1. Total and dissolved mercury and methylmercury for the initial surface water	r sampling
event, upstream to downstream <sup>a</sup>	
Figure 3.2. Total and dissolved mercury and methylmercury for the April 2024 surface sampling event, upstream to downstream	
Figure 3.3. Total and dissolved methylmercury to mercury ratios for the April 2024 surf sampling event, upstream to downstream	
Figure 3.4. Concentration graphs for selected analytes in Bear Creek (upstream to down and Hinds Creek reference site surface water, original sampling event dat	,
Figure 3.5. Bear Creek floodplain soil, creek bank soil, and channel sediment average pasize distributions.	
Figure 3.6. BCT (upstream to downstream) and Hinds Creek floodplain soil chemical day	ata3-21
Figure 3.7. BCT (upstream to downstream) and Hinds Creek upper creek bank soil cher	nical data 3-22
Figure 3.8. BCT (upstream to downstream) and Hinds Creek lower creek bank soil cher	nical data 3-23
Figure 3.9. BCT (upstream to downstream) channel sediment chemical data	3-24
Figure 3.10. Bear Creek channel sediment and April 2024 surface water percent methyl shown upstream to downstream	
Figure 3.11. Results of Bear Creek floodplain soil sequential mercury extraction	
Figure 3.12. Results of Bear Creek bank soil sequential mercury extraction.	
Figure 3.13. Results of Bear Creek channel sediment sequential mercury extraction	
Figure 3.14. Sequential mercury extraction results at location BCT5.	3-32
Figure 3.15. Sequential mercury extraction results at location BCT6.	
Figure 3.16. Sequential mercury extraction results at location BCT7.	
Figure 3.17. Sequential mercury extraction results at location BCT8.	
Figure 3.18. Sequential mercury extraction results at location BCT9.	3-36
Figure 3.19. Sequential mercury extraction results at location BCT11.	3-37
Figure 3.20. Sequential mercury extraction results at location BCT12B	
Figure 3.21. Sequential mercury extraction results at location BCT14.	
Figure 3.22. Sequential mercury extraction results at the Hinds Creek reference site (HC	
Figure 4.1. Summary of WRRP surface water total mercury data, 2011–2023	4-1
Figure 4.2. Summary of WRRP surface water methylmercury data, 2011–2023	4-2
Figure 4.3. Flow rate versus total mercury and methylmercury concentrations at BCK 1:  Bear Creek NT-3	2.34 and4-3

Figure 4.4. Flow rate versus total mercury and methylmercury concentrations at BCK 11.54A and Bear Creek NT-8	4-4
Figure 4.5. Flow rate versus total mercury and methylmercury concentrations at spring SS-5 and BCK 9.2.	4-:
Figure 4.6. Daily total mercury flux rate for surface water flow measurement stations	4-0
Figure 4.7. Bear Creek Zone 3 surface water total mercury flux proportions measured at various monitoring stations	4-8
Figure 4.8. Bear Creek Zone 3 estimated total methylmercury daily flux rates.	4-9
Figure 4.9. Bear Creek Zone 3 estimated total methylmercury flux proportions measured at various monitoring stations.	4-10
Figure 4.10. Average concentrations of mercury in Bear Creek fish	4-11

## **TABLES**

Table 1.1. BCV cap and soil cover areas	1-9
Table 1.2. Routine surface water and biota sampling in Bear Creek	1-10
Table 2.1. List of transect locations in BCV	2-1
Table 2.2. Sampling summary by location	2-2
Table 2.3. Summary of field and laboratory measurements	2-6
Table 3.1. Sample location site characteristics	3-1
Table 3.2. Bear Creek mercury sources RSE transects surface water field parameters from the initial sampling event	3-2
Table 3.3. Bear Creek mercury sources RSE transects surface water field parameters, April 2024	3-3
Table 3.4. Summary of selected analytes in surface water at BCT locations collected from the initial sampling event	3-5
Table 3.5. Surface water total and dissolved mercury and methylmercury, April 2024 sampling event	3-8
Table 3.6. Bear Creek and Hinds Creek soil and sediment particle size data	3-14
Table 3.7. Summary of selected analyte concentrations in BCTs floodplain soil, upper and lower creek bank soil, and channel sediment samples	3-19
Table 3.8. Mercury sequential extraction method fractions and descriptions	3-26
Table 3.9. Results of mercury sequential extraction analyses from floodplain soil, creek bank soil, and channel sediment at eight BCT locations and at the Hinds Creek reference site	3-27
Table 3.10. Comparison of channel sediment sequential mercury extraction steps F3, F4, and F5 averages in Bear Creek and LEFPC	3-41

#### **ACRONYMS**

amsl above mean sea level

AWQC ambient water quality criteria

BCK Bear Creek kilometer
BCT Bear Creek transect
BCV Bear Creek Valley
BYBY Boneyard/Burnyard
COC contaminant of concern

CR Clinch River

CSM conceptual site model
DOC dissolved organic carbon
DOE U.S. Department of Energy
EFPC East Fork Poplar Creek

EMDF Environmental Management Disposal Facility

EPA U.S. Environmental Protection Agency

FFA Federal Facility Agreement

FY fiscal year

HCDA Hazardous Chemical Disposal Area

HCK Hinds Creek kilometer

LEFPC Lower East Fork Poplar Creek

NT North Tributary

OREIS Oak Ridge Environmental Information System

ORNL Oak Ridge National Laboratory

ORR Oak Ridge Reservation

OU operable unit ppb parts per billion QA quality assurance

redox oxidation-reduction potential RER Remediation Effectiveness Report

RI remedial investigation
ROD Record of Decision
RSE remedial site evaluation
SAP Sampling and Analysis Plan

TDEC Tennessee Department of Environment and Conservation

TOC total organic carbon

UEFPC Upper East Fork Poplar Creek

WRRP Water Resources Restoration Program Y-12 Y-12 National Security Complex

#### **EXECUTIVE SUMMARY**

This report presents results of the Bear Creek Valley (BCV) mercury sources remedial site evaluation (RSE) conducted in fiscal year 2024. The objective of the BCV mercury sources RSE is to evaluate potential sources of mercury and methylmercury within the BCV Watershed, as included in the mercury-management approach for Bear Creek in the *Record of Decision for Comprehensive Environmental Response, Compensation, and Liability Act Oak Ridge Reservation Waste Disposal at the Environmental Management Disposal Facility, Oak Ridge, Tennessee* (DOE/OR/01-2794&D2/R2), to determine if active remediation is warranted. The U.S. Department of Energy collected and evaluated the BCV mercury sources RSE data, including channel sediment, creek bank and floodplain soil, and surface water. All sampling was conducted as planned between December 2023 and April 2024, per the *Bear Creek Valley Mercury Sources Remedial Site Evaluation Sampling and Analysis Plan, Oak Ridge, Tennessee* (DOE/OR/01-2958&D2), with the exception of a few changes due to field conditions and stream morphology. This BCV mercury sources RSE did not identify a source of mercury that significantly contributes to Bear Creek mercury contamination or mercury bioaccumulation in fish, or that would warrant active remediation. In addition, all Bear Creek surface water samples collected in this evaluation had mercury concentrations less than the 51-ng/L ambient water quality criteria level.

The results of this evaluation add significant data to the Bear Creek mercury conceptual model in several ways. Data did not identify one or more specific principal source(s) of mercury to Bear Creek that would indicate an early Comprehensive Environmental Response, Compensation, and Liability Act of 1980 (CERCLA) action for mercury management/reduction. The distribution of mercury in floodplain and creek bank soils and channel sediment shows the highest concentrations in BCV headwater areas upstream of the confluence with North Tributary-3. The vicinity of the Reeves Road crossing of Bear Creek, where beaver ponds have been prevalent for several years, stands out as an inflection point in the valley-wide mercury concentration gradient. A key finding of the evaluation comes from the data provided through the mercury sequential extraction analyses that document a majority of the mercury in BCV floodplain soil, creek bank soil, and channel sediment is associated with organic components of the media. This result contrasts significantly from conditions in East Fork Poplar Creek, where mercury is predominantly associated with more strongly bound fractions of the mercury sequential extraction series. The mercury association with organic components of Bear Creek media may play a key role in the apparent ease of methylation and subsequent bioaccumulation in aquatic biota.

Although this BCV mercury sources RSE determines no remedial investigations or actions are required at this time, enhanced monitoring of instream mercury and methylmercury concentrations and flux rates and further development of the mercury bioaccumulation conceptual model continue under the mercury remediation technology development program. Recommendations for future technology development program investigations to enhance the understanding of the mercury bioaccumulation conceptual model are provided. While no remedial action is required at this time, the scientific investigations proposed may indicate an opportunity to reduce mercury in Bear Creek in the future, and CERCLA actions will be taken at that time as appropriate.

#### 1. INTRODUCTION

This report presents results of the Bear Creek Valley (BCV) mercury sources remedial site evaluation (RSE) conducted in fiscal year (FY) 2024. Introductory information about the BCV mercury sources RSE objective, site description, summary of potential mercury source areas, summary of historical BCV monitoring results, and report content is provided below.

#### 1.1 REMEDIAL SITE EVALUATION OBJECTIVE

The objective of the BCV mercury sources RSE is to evaluate potential sources of mercury and methylmercury within the BCV Watershed, as included in the mercury-management approach for Bear Creek in the *Record of Decision for Comprehensive Environmental Response, Compensation, and Liability Act Oak Ridge Reservation Waste Disposal at the Environmental Management Disposal Facility, Oak Ridge, Tennessee* (DOE/OR/01-2794&D2/R2; Environmental Management Disposal Facility [EMDF] Record of Decision [ROD]), to determine if active remediation is warranted. The BCV Watershed is located in the north-central portion of the Oak Ridge Reservation (ORR) west of the Y-12 National Security Complex (Y-12) (Figure 1.1). Y-12 began operations in the 1940s as part of the Manhattan Project for the purpose of enriching uranium for the first atomic bombs. Since that time, the Y-12 missions have changed, and in the 1950s, new processes for separating lithium used large amounts of mercury. Although process functions were performed adjacent to BCV in the Y-12 Main Plant area, wastes from operations at Y-12 were disposed in pits, trenches, and burial grounds in the 2800-acre BCV Watershed.

The Bear Creek Valley Mercury Sources Remedial Site Evaluation Sampling and Analysis Plan, Oak Ridge, Tennessee (DOE/OR/01-2958&D2; BCV Mercury Sources RSE Sampling and Analysis Plan [SAP]) identified the locations, media, and sampling methodology to support the RSE objectives. Impacts of source areas and hydrology on mercury concentrations are assessed in channel sediment, creek bank and floodplain soils, and surface water at multiple sampling transects throughout the length of the stream. The data quality objectives for the BCV mercury sources RSE and the BCV Mercury Sources RSE SAP were presented and discussed with the Project Team, comprised of representatives of the Tennessee Department of Environment and Conservation (TDEC), the U.S. Environmental Protection Agency (EPA), the U.S. Department of Energy (DOE), and the contractor (United Cleanup Oak Ridge LLC), on June 29, 2023. Results of the RSE are combined with biota data from fall 2023 to evaluate potential mercury source areas in BCV. The BCV Mercury Sources RSE SAP is included as Appendix A of this document.

#### 1.2 BACKGROUND

#### 1.2.1 Site Description

The BCV Watershed is located at the western end of Y-12 in the north-central portion of the ORR west of the Upper East Fork Poplar Creek (UEFPC) Watershed (Figure 1.1). BCV contains closed and active waste disposal facilities. The boundary between the BCV Watershed and the UEFPC Watershed is defined by a surface water divide between eastward-flowing East Fork Poplar Creek (EFPC) and westward-flowing Bear Creek (Figure 1.2). The integration point for Bear Creek is at Bear Creek kilometer (BCK) 9.2, where more than 99% of the available water from the eastern portion of BCV passes through this location either as surface water or groundwater. As illustrated in Figure 1.2, the BCV Administrative Watershed is subdivided into three zones based on end use. The subareas of BCV investigated under the BCV Mercury Sources RSE SAP represent geographic areas located at or downstream from potential DOE on-site source areas. Based on the EMDF ROD, the end use for Zone 1 and Zone 2 will be revised to restricted recreational and controlled industrial, respectively, which will be codified in an upcoming non-significant change to the *Record of Decision for the Phase I Activities in Bear Creek Valley at the Oak Ridge Y-12 Plant, Oak Ridge, Tennessee* (DOE/OR/01-1750&D4; BCV Phase I ROD). The following sections briefly summarize the BCV geography, geology, and hydrogeologic conceptual model.




Figure 1.1. BCV within the Oak Ridge Site.




Figure 1.2. BCV Watershed and historic monitoring locations.

#### 1.2.1.1 Geography

The ORR is approximately 32,500 acres and is located within and adjacent to the corporate limits of the City of Oak Ridge, Tennessee, in Roane and Anderson Counties. The ORR is bounded to the east, south, and west by the Clinch River (CR) and on the north by the City of Oak Ridge. The area west of the ORR and the CR is comprised of rural, residential, and industrial park areas.

The entire BCV Watershed is approximately 4800 acres and encompasses two principal drainage areas: (1) the catchment area of Bear Creek and its tributaries between the western end of Y-12 to approximately 0.25 miles west of State Route 95 (enclosed between Pine Ridge and Chestnut Ridge), also described as the BCV Administrative Watershed; and (2) an area between the Bear Creek water gap in Pine Ridge and the confluence with EFPC (Figure 1.2). The BCV Administrative Watershed is approximately 2800 acres and lies within the north-central portion of the ORR, approximately 0.75 miles south of the industrial/residential portion of the City of Oak Ridge (Figure 1.2).

Bear Creek is wholly contained on the ORR and originates in the eastern portion of the watershed near the former S-3 Site. It then flows west along strike of BCV to State Route 95 and thereafter flows north and empties into EFPC 12.87 km (8 miles) downstream. The average gradient of the creek is 30 ft/mile, dropping from nearly 1000 ft above mean sea level (amsl) at its headwaters to 760 ft amsl at its confluence with EFPC. A series of tributaries draining the south flank of Pine Ridge and springs along the southern edge of BCV are key contributors to the flow in Bear Creek (Figure 1.2).

#### 1.2.1.2 BCV Watershed hydrogeological conceptual model

A full description of the hydrogeologic conceptual model for the BCV Watershed may be found in the Report on the Remedial Investigation for Bear Creek Valley at the Oak Ridge Y-12 Plant, Oak Ridge, Tennessee (DOE/OR/01-1455&D2; BCV Remedial Investigation [RI]). Groundwater flow and contaminant transport within the bedrock geologic units are important factors in the migration of contaminants from the watershed. Bedrock geologic units strike northeast-southwest, parallel to the axis of the valley, with predominantly shale formations underlying the central and northern portions of the valley and most major waste disposal units (Figure 1.3). In the northern portion of the valley, shallow groundwater and storm flow through unconsolidated material and weathered bedrock are important components of groundwater recharge and contaminant migration from source units to Bear Creek and its tributaries. Waste and contamination at the sources in BCV are situated in the subsurface, and shallow groundwater is the principal mechanism and pathway for release of contaminants. After release, most contaminants travel via short pathways in shallow groundwater (i.e., storm flow zone) to be discharged into tributaries to Bear Creek. Some contaminants, in particular those from the S-3 Ponds, remain entrained in groundwater and discharge directly into the underlying stratum.

The southern portion of the valley is underlain by the Maynardville Limestone, a 200-ft-thick limestone formation, containing a well-developed karst network created by dissolution and enlargement of fractures and joints. The underlying geology results in an asymmetric topographic cross-section of the valley, with the lowest elevations on the south side coincident with the Maynardville Limestone. Groundwater flow in the Maynardville Limestone occurs in both shallow and deep karst features, and corresponding flow rates and volumes are much higher than in the shale-dominated formation underlying the central and north portions of the valley. Large, individual springs or groups of springs mark the locations of discharge from both the shallow and deep karst flow systems into the surface water system. In addition, surface water flow in Bear Creek is highly connected with groundwater in the underlying Maynardville Limestone through karst features and losing and gaining reaches of the creek. Because of these characteristics, the interconnected Bear Creek channel and underlying karst system act as the principal hydraulic drain for the valley and are part of the carbonate aquifer system in the BCV conceptual site model (CSM).




Figure 1.3. BCV hydrogeologic CSM for Upper Bear Creek.

By comparison, the shale-dominated formations in the central and northern portions of the valley function as aquitards. Because of the karst conditions in the Maynardville Limestone, during late summer into autumn, reaches of Bear Creek (much of the reach from North Tributary [NT]-4 to NT-8) are prone to becoming dry (Figure 1.3). During the drought season, these reaches of Bear Creek are not supportive of fish, although small populations may persist in scattered pools and tributary streams.

The climate of the Oak Ridge area and its surroundings may be broadly classified as humid subtropical. The term "humid" indicates the region receives an overall surplus of precipitation compared to the levels of evaporation and transpiration throughout the year, while the "subtropical" designation indicates the region experiences a wide range of seasonal temperatures. Average annual precipitation in Oak Ridge is 56.3 in. Wet deposition is a major source of mercury entering aquatic and terrestrial ecosystems, where it can cause significant ecological and human health risks. The wet deposition of mercury is, therefore, measured globally by the Mercury Deposition Network. The total mercury wet deposition in the southeastern United States is relatively high (Xiaotian et al. 2002); the wet deposition in East Tennessee in 2022 was approximately 14  $\mu$ g/m²; however, site-specific data for Bear Creek are not available. Surface water total mercury concentrations have historically been relatively low, but mercury concentrations in fish collected in Bear Creek are relatively high, occasionally exceeding the EPA-recommended ambient water quality criterion (AWQC) for mercury (0.3  $\mu$ g/g in fish) until recent years (ORNL/TM-2023/3069).

Recently, the Oak Ridge National Laboratory (ORNL) conducted special studies of Bear Creek to better understand the biotic and abiotic factors contributing to mercury concentrations in fish in Bear Creek. These field studies focused on gaining an understanding of the processes controlling mercury methylation and bioaccumulation, with beaver dams and periphyton being key areas of interest. Studies included understanding the role of beaver dams in mercury dynamics in Bear Creek (2017–2018), evaluating the effects of fine-grained sediment deposition (2019), investigating the potential role tributaries to Bear Creek may have on mercury and methylmercury in the main channel (2020–2021), and evaluating periphyton relationships (2021). These special studies were documented in the *Bear Creek Special Studies Report 2021* (ORNL/SPR-2021/2162).

A 2022 data compilation report for mercury in Bear Creek (ORNL/TM-2023/3069) summarized data from compliance and investigatory studies to begin building a conceptual model to understand the processes affecting mercury transport and transformation in the Bear Creek Watershed and to highlight key knowledge gaps in understanding these processes. This report summarizes historical data from compliance and investigatory studies relevant to mercury contamination and transformation in the Bear Creek Watershed and highlights the relative paucity of water quality data in Bear Creek with respect to mercury concentrations and transformation processes. A recommendation from this study was to install a long-term monitoring station to provide needed data to establish functional linkages among identified components of a conceptual model of mercury transport with measured mercury values from the watershed.

The mercury remediation technology development program at ORNL, which began in 2014, originally focused on understanding and addressing mercury concentrations, flux, and bioaccumulation in EFPC. Research from this program highlighted the importance of periphyton in mercury transformation in stream systems, and in 2021, a task specifically focused on periphyton dynamics and distribution was added to the study plan. Because the aqueous methylmercury concentrations in EFPC are similar to those in nearby Bear Creek despite significantly higher aqueous total mercury concentrations in EFPC, comparing mercury methylation and demethylation processes between the two streams was of particular interest, and Bear Creek became (and remains) a point of focus in the mercury remediation technology development program (ORNL/SPR-2023/3178).

Future work will build on the summary presented in the data compilation report to develop a conceptual model that outlines the key environmental parameters that correlate with methylmercury concentrations

and bioaccumulation in Bear Creek. A conceptual model will help to understand the processes affecting mercury transport and transformation in the Bear Creek Watershed and to highlight key knowledge gaps in understanding these processes. The conceptual model will ultimately provide a strong technical basis for prioritizing new data collection and optimizing potential mitigation actions or best management practices, with the goal of lowering fish tissue mercury concentrations.

#### 1.2.2 Summary of Potential Mercury Source Areas

BCV contains multiple historical waste management and disposal areas that received mercury-contaminated waste streams from Y-12 operations from 1943–1993, in addition to having materials storage areas and construction storage areas (Figure 1.2). The BCV RI and associated decision documents cite mercury as a potential contaminant of concern (COC) in BCV at the following locations:

- Boneyard/Burnyard (BYBY)
- Oil Landfarm, Hazardous Chemical Disposal Area (HCDA)
- S-3 Ponds Site
- Sanitary Landfill 1
- Bear Creek Road Debris Burial
- Creekside Debris Burial

A source control action performed under the BCV Phase I ROD at the remediated BYBY (*Phased Construction Completion Report for the Bear Creek Valley Boneyard/Burnyard Remediation Project at the Y-12 National Security Complex, Oak Ridge, Tennessee* [DOE/OR/01-2077&D2]) focused on excavating mercury-contaminated soil along NT-3. Mercury surface water results in BCV are consistently below Tennessee general AWQC (TDEC 2019); however, until recent years, fish tissue concentrations have remained above or near the EPA-recommended AWQC for mercury (0.3 µg/g in fish). The BCV RI indicated some elevated soil mercury concentrations exist, generally within an order of magnitude of the background criterion (0.34 mg/kg); however, historical mercury and methylmercury data for sediment and soil that may contribute to concentrations in fish are limited. The baseline risk assessment in the BCV RI stated "the sources of mercury and PCBs to the BCV fish are currently unknown."

The Remedial Investigation Report on Bear Creek Valley Operable Unit 2 (Rust Spoil Area, Spoil Area 1, and SY-200 Yard) at the Oak Ridge Y-12 Plant, Oak Ridge, Tennessee (DOE/OR/01-1273&D2; BCV Operable Unit [OU] 2 RI) identified mercury as a COC for human health for the SY-200 Yard, which was a former equipment storage yard used to store nonradioactive contaminated equipment from the 1950s to 1986. Mercury contamination was discovered during construction in 1990, and a soil cover was placed over the site. While other areas (Spoil Area 1 and the Rust Spoil Area) had mercury as a contaminant of potential concern in the BCV OU 2 RI, the baseline risk assessment did not identify mercury as a COC for these areas. The BCV OU 2 RI indicated there were isolated areas of elevated mercury concentrations at the SY-200 Yard; free mercury was observed in some of the borings during the BCV OU 2 RI. The Record of Decision for Bear Creek Operable Unit 2 (Spoil Area 1 and SY-200 Yard) at the Oak Ridge Y-12 Plant, Oak Ridge, Tennessee (DOE/OR/02-1435&D2; BCV OU 2 ROD) identified the SY-200 Yard soil cover are ongoing.

Many of the disposal sites have caps or soil covers that are maintained as described in Table 1.1.

Table 1.1. BCV cap and soil cover areas

Potential mercury source areas	Cap/soil cover details
ВУВУ	BYBY was covered with a 2-ft layer of low-permeability clay and a 6-in. layer of vegetative cover
Oil Landfarm and HCDA	HCDA was covered with a low-permeability RCRA-type cap in 1989; however, this was not a RCRA-regulated facility
	The Oil Landfarm was covered with a multilayer low-permeability cap, consisting of a minimum of 2 ft of compacted clay, a flexible synthetic membrane liner, a geosynthetic drainage net, a geotextile filter fabric, a minimum of 1.5 ft of topsoil, cover vegetation, and cap drains. The final RCRA closure for the Oil Landfarm was certified by TDEC in December 1990
S-3 Ponds Site	The S-3 Ponds Site was covered with a multilayer low-permeability engineered cap, covered with crushed stone, and paved with asphalt for use as a passenger vehicle parking lot. TDEC accepted the final RCRA closure certification for the S-3 Ponds Site on November 15, 1990
Sanitary Landfill 1	The landfill was closed in 1985 by grading to promote drainage, capping with 2 ft of clay and topsoil, and establishing a vegetative cover
Bear Creek Road Debris Burial	The BCV Phase I ROD concluded this area did not contain significant sources of contamination that could pose a risk to health or the environment; therefore, no action was warranted under CERCLA
Creekside Debris Burial	The BCV Phase I ROD concluded this area did not contain significant sources of contamination that could pose a risk to health or the environment; therefore, no action was warranted under CERCLA
SY-200 Yard	A compacted soil cover (3 to 5 ft) was installed over the site beginning in 1986
Spoil Area 1	A 2-ft-minimum vegetative clay soil cover was installed over the site beginning in 1985
Rust Spoil Area	A minimum of 1.5 ft of compacted clay and a 0.5-ft topsoil soil cover were installed over the site beginning in 1983

BCV Phase I ROD = DOE/OR/01-1750&D4. Record of Decision for the Phase I Activities in Bear Creek Valley at the Oak Ridge Y-12 Plant, Oak Ridge, Tennessee, 2000, U.S. Department of Energy, Office of Environmental Management, Oak Ridge, TN.

BCV = Bear Creek Valley

BYBY = Boneyard/Burnyard

CERCLA = Comprehensive Environmental Response, Compensation, and Liability Act of 1980

HCDA = Hazardous Chemical Disposal Area

RCRA = Resource Conservation and Recovery Act of 1976

ROD = Record of Decision

SY = scrapyard

TDEC = Tennessee Department of Environment and Conservation

#### 1.2.3 Summary of Historical Bear Creek Valley Mercury Data

Mercury data for sediment, surface water, and biota in BCV are available in the Oak Ridge Environmental Information System (OREIS). However, prior to this BCV mercury sources RSE, sediment data were limited, and no methylmercury data were available for BCV sediment in OREIS. Twenty-nine data points for total mercury in BCV sediment were available ranging from 1993–2011: 7 locations in Zone 1, 2 locations in Zone 2, and 20 locations in Zone 3. Concentrations ranged from non-detected to 6.9-mg/kg total mercury.

As shown in Table 1.2, under the *Bear Creek Valley Watershed Remedial Action Report Comprehensive Monitoring Plan, Oak Ridge, Tennessee* (DOE/OR/01-2457&D4), surface water and biota sampling have been performed. Results are reported annually in a Remediation Effectiveness Report (RER) or every 5 years in a Five-Year Review Report. Surface water data since 2011 (Figure 1.4) show a steady or declining trend, with mercury below AWQC, with few exceptions. Mercury concentrations are generally higher

upstream and decrease downstream; methylmercury concentrations are more variable upstream to downstream. Additional evaluation of historical mercury and methylmercury surface water data is included in Chapter 4. Fish tissue data are presented in Section 4.2.

Table 1.2. Routine surface water and biota sampling in Bear Creek

Medium	Performance standard	Sampling frequency	Parameter	Monitoring location
	AWQC	Semiannual (Q1 and Q3)	Total mercury and methylmercury	BCK 3.3, BCK 4.55, BCK 9.2, BCK-11.54A, BCK 12.34, NT-3, SS-4, and SS-5
		Semiannual (Q2 and Q4) in year before FYR	Total mercury	BCK 4.55, BCK 9.2, BCK 12.34, NT-3, and NT-8
		Annual in year before FYR	Total mercury	BCK-7.87 and NT-1
Surface water		Annual in year before FYR	Methylmercury	NT-5
	Trend monitoring	Quarterly		NT-1, NT-2, NT-3, SS-4, and SS-5
		Semiannual	Bicarbonate, chloride,	NT-7 and NT-8
		Quarterly in year before FYR	fluoride, and sulfate	NT-5
	Water quality	Semiannual	Total suspended solids	NT-7 and NT-8
		Quarterly in year before FYR	and total dissolved solids	BCK 4.55, BCK-7.87, BCK 9.2, BCK 12.34, NT-1, NT-3, NT-5, and NT-8
Biota	Baseline sampling	Semiannual	Mercury and methylmercury	BCK 3.3, BCK 9.9, and HCK 20.6 (whole-body stoneroller minnows and rock bass fillets); BCK 12.4 (whole-body stoneroller minnows)
	. 0	Annual in year before FYR	Mercury and methylmercury	BCK 9.9 (whole-body caddisflies)

AWQC = ambient water quality criteria BCK = Bear Creek kilometer

FYR = Five-Year Review

HCK = Hinds Creek kilometer NT = North Tributary Q = quarter

SS = surface spring




Figure 1.4. Surface water data for mercury and methylmercury in Bear Creek, NT-3,

and Hinds Creek, 2011–2022.

#### 1.3 REPORT CONTENT

Chapter 2 presents information about implementing the BCV Mercury Sources SAP, including scope of work and changes from the BCV Mercury Sources RSE SAP during field implementation. Chapter 3 summarizes monitoring results for all media collected during the BCV mercury sources RSE. Chapter 4 presents other surface water monitoring data collected by the Water Resources Restoration Program (WRRP) and presents the 2023 biota data collected for the upcoming 2025 RER. The report conclusions and recommendations are presented in Chapter 5. Chapter 6 lists references cited in the report. This report contains seven appendices with supporting information:

- Appendix A. BCV Mercury Sources RSE SAP, including TDEC and EPA comments on the D1 version of the document.
- Appendix B. Photographs of Transect Locations.
- Appendix C. Analytical Data Summary Tables by Media.
- Appendix D. Particle Size Analysis.
- Appendix E. BCV RSE Analyte Correlations.
- Appendix F. WRRP Surface Water Total Mercury and Methylmercury Longitudinal Data Plots for Bear Creek.
- Appendix G. Bear Creek Field Observations.

## 2. IMPLEMENTATION OF THE BEAR CREEK VALLEY MERCURY SOURCES REMEDIAL SITE EVALUATION SAMPLING AND ANALYSIS PLAN

#### 2.1 SCOPE OF WORK

Field implementation, sample collection, laboratory analysis, and data management activities were consistent with the BCV Mercury Sources RSE SAP and were conducted within the framework of plans, procedures, and protocols under the WRRP that help ensure all data collected are managed in a manner consistent with Comprehensive Environmental Response, Compensation, and Liability Act of 1980 requirements. The *Quality Assurance Project Plan for the Water Resources Restoration Program, U.S. Department of Energy, Oak Ridge Reservation, Oak Ridge, Tennessee* (UCOR-4049) identifies and implements quality assurance (QA) requirements for use in sample collection, laboratory analysis, and data management of environmental media monitoring activities. The *Data Management Implementation Plan for the Water Resources Restoration Program, Oak Ridge, Tennessee* (UCOR-4160) serves as the project-level plan for managing all data collected by the WRRP. Together, these plans identify the procedures that are followed in collecting, handling, and maintaining custody of, as well as in verifying, validating, and retaining environmental and laboratory data used by the WRRP to prepare Federal Facility Agreement (FFA) documents. Additional requirements governing fieldwork and sample collection, including QA/quality control samples are specified in the *Quality Assurance Plan for Environmental Characterization and Monitoring, Oak Ridge, Tennessee* (UCOR-4189).

A list of sampling locations along Bear Creek and its tributaries includes transects for the collection of channel sediment, creek bank and floodplain soils, and surface water (Table 2.1). Samples collected per the BCV Mercury Sources RSE SAP and any deviations from the BCV Mercury Sources RSE SAP due to field conditions are shown on Table 2.2. Sample transects are identified by increasing numbered locations (e.g., Bear Creek transect [BCT]1, BCT2) from the most downstream locations northwest of Tennessee Highway 58 to the most upstream location (BCT15) that is located near the western end of the Y-12 facility (Figure 2.1). Location HCTREF is the project reference site on Hinds Creek near Clinton, Tennessee. A conceptual diagram of the transect sampling is included as Figure 2.2.

Table 2.1. List of transect locations in BCV

Sample group	Location
	BCT1 (downstream of BCK 0.6)
LOWBCV	BCT2 (upstream of BCK 0.6; beaver dam previously viewed near this transect was no longer present)
LOWBEV	BCT3 (downstream of BCK 3.3; downstream of beaver dam)
	BCT4 (downstream of BCK 3.3; in beaver pond)
	BCT5 (downstream of BCK 4.55; beaver dam previously viewed near this transect was no longer present)
	BCT6 (downstream of BCK 4.55; beaver dam previously viewed near this transect was no longer present)
BCV ZONE 1	BCT7 (downstream of BCK 7.87 at the confluence of NT-13/Bear Creek; downstream of westernmost beaver dam)
	BCT8 (downstream of BCK 7.87 at the confluence of NT-13/Bear Creek; upstream of westernmost beaver dam)
	BCT9 (downstream of BCK 7.87; upstream of two beaver dams; southeast of Reeves Road/Haul Road)
BCV ZONE 2	BCT10 (downstream of surface water integration point BCK 9.2; upstream of EMDF)

Table 2.1. List of transect locations in BCV (cont.)

Sample group	Location
	BCT11 (upstream of NT-8 and BCK 9.9)
	BCT12A (upstream of NT-3 confluence of NT-3/Bear Creek)
DOW ZONE 2	BCT12B (downstream of BYBY at the confluence of NT-3/Bear Creek)
BCV ZONE 3	BCT13 (upstream of BYBY, EMWMF, and NT-3)
	BCT14 (downstream of SY-200 Yard, Spoil Area 1, S-3 Ponds Site, and BCK 12.34)
	BCT15 (downstream of SY-200 Yard, Spoil Area 1, and S-3 Ponds Site at NT-1)
Hinds Creek	HCTREF (HCK 20.6 reference site)

BCK = Bear Creek kilometer BCT = Bear Creek transect

BCV = Bear Creek Valley

BY BY Boneyard/Burnyard
EMDF = Environmental Management Disposal Facility
EMWMF = Environmental Management Waste Management Facility

HCK = Hinds Creek kilometer

HCTREF = Hinds Creek transect reference site

NT = North Tributary

Table 2.2. Sampling summary by location

T	El l . l	Upper	Lower	CI I		Sequential extraction		
Transect location	Floodplain soil	creek bank soil	creek bank soil	Channel sediment	Surface water	Floodplain soil	Creek bank soil	Sediment
BCT1	X	X	X		X	NS	NS	NS
BCT2	X	X	X		X	NS	NS	NS
BCT3	X	X	X	X	X	NS	NS	NS
BCT4	X	X	X	X	X	NS	NS	NS
BCT5	X	X	X		X	X	X	
BCT6	X	X	X		X	X	X	
BCT7	X	X		X	X	X	X	X
BCT8	X	X		X	X	X	X	X
BCT9	X	X		X	X	X	X	X
BCT10	X	X	X	X	X	NS	NS	NS
BCT11	X	X	X	X	X	X	X	X
BCT12A	X	X	X	X	X	NS	NS	NS
BCT12B	X	X	X	X	X	X	X	X
BCT13	X	X	X	X	X	NS	NS	NS
BCT14	X	X	X	X	X	X	X	X
BCT15	X	X	X	X	X	NS	NS	NS
HCTREF	X	X	X		X	X	X	

<sup>--</sup> Indicates sample unable to be collected either due to lack of fine-grained material or limited creek bank exposure.

BCT = Bear Creek transect

HCTREF = Hinds Creek transect reference site

NS = sequential extraction was not planned for this location; therefore, no sample was collected

x Indicates sample collected.




Figure 2.1. BCV transect locations.




Figure 2.2. BCV RSE transect conceptual diagram.

#### 2.1.1 Channel Sediment Sampling

Channel sediment samples were collected at 12 transect monitoring locations following PROC-ES-2302, *Sediment Sampling*. No channel sediment was sampled at BCT1, BCT2, BCT5, BCT6, or the reference site HCTREF (Hinds Creek kilometer [HCK] 20.6), because no fine-grained sediment was observed at these locations. Sediment was collected to an approximate depth of 0.5 ft and was run through a 1-mm sieve until adequate sample volume was achieved. All channel sediment samples were analyzed for mercury, methylmercury, metals, total organic carbon (TOC), anions, and particle size (Table 2.3). In addition, sequential extraction of mercury was conducted at select transect locations—BCT7, BCT8, BCT9, BCT11, BCT12B, and BCT14. Planned sequential extraction at locations BCT5 and BCT6 and the reference site HCTREF (HCK 20.6) was not possible due to lack of fine-grained sediment at these locations.

Table 2.3. Summary of field and laboratory measurements

Medium	Field measurement	Laboratory measurement
Surface water	Temperature	Dissolved and total mercury
	Dissolved oxygen	Dissolved and total methylmercury
	Turbidity	Metals
	pН	Phosphorous (total)
	Specific conductance (conductivity)	Total organic carbon
	Oxidation-reduction potential	Dissolved organic carbon
		Total dissolved solids
		Total suspended solids
		Anions (chloride, fluoride, nitrate-nitrite as nitrogen, sulfate, and sulfide)
Sediment and	None	Total mercury
soil		Total methylmercury
		Metals
		Total organic carbon
		Particle size analysis
		Anions (chloride, fluoride, nitrate, nitrite, sulfate, and sulfide)
		Sequential extraction of mercury (select locations)

#### 2.1.2 Creek Bank Soil Sampling

Creek bank soil was collected at 16 transect monitoring locations as well as at a single reference site—HCTREF (HCK 20.6)—following PROC-ES-2300, *Soil Sampling*. Except for transects BCT7, BCT8, and BCT9, at which ponding limited the height of the banks, creek bank soil was divided in half into upper and lower sections as follows (Figure 2.2):

- For the upper section of the creek bank soils, samples were collected from the upper half of the exposed bank face on each side of the creek. The upper creek bank samples from both sides were composited into a single sample.
- For the lower section of the creek bank soils, samples were collected from the lower half of the exposed bank face on each side of the creek. The lower creek bank samples from both sides were composited into a single sample.

• At BCT7, BCT8, and BCT9, an aliquot of soil material was collected from each side of the bank and composited into a single sample. These samples were treated as upper bank soil samples for evaluations.

All creek bank soil samples were analyzed for mercury, methylmercury, metals, TOC, anions, and particle size (Table 2.3). In addition, sequential extraction of mercury was conducted at select transect locations—BCT5, BCT6, BCT7, BCT8, BCT9, BCT11, BCT12B, and BCT14—and the reference site HCTREF (HCK 20.6).

#### 2.1.3 Floodplain Soil Sampling

Floodplain soils were collected at each of the 16 transect monitoring locations as well as at a single reference site—HCTREF (HCK 20.6)—following PROC-ES-2300, *Soil Sampling*.

Floodplain soil was collected from the upper 0.5 ft on each side of Bear Creek to generate a single composite sample representing both sides of the floodplain (Figure 2.2). Loose organic material, such as leaves or brush, was removed prior to collection. All floodplain soil samples were analyzed for mercury, methylmercury, metals, TOC, anions, and particle size (Table 2.3). In addition, sequential extraction of mercury was conducted at select transect locations—BCT5, BCT6, BCT7, BCT8, BCT9, BCT11, BCT12B, and BCT14—and the reference site HCTREF (HCK 20.6).

#### 2.1.4 Surface Water Sampling

Surface water samples were collected at each of the 16 transect monitoring locations as well as at a single reference site—HCTREF (HCK 20.6)—following PROC-ES-2203, *Surface Water Sampling – Manual and Automated*.

Because filtered (dissolved) and unfiltered (total) mercury and methylmercury sample volumes were collected for analysis, a peristaltic pump was used to push water through a 0.45-µm filter in addition to the grab method for unfiltered samples. Surface water sampling was conducted before channel sediment was collected to avoid interference between media. All surface water samples were analyzed for total and dissolved mercury and methylmercury, metals, phosphorous, TOC, dissolved organic carbon (DOC), total dissolved solids, total suspended solids, and anions.

#### 2.1.5 Speciation of Mercury via Sequential Extraction

Sequential extractions of soils allow the concentrations of mercury compounds in the soil that belong to different classes (e.g., water soluble, weak acid soluble, organo-complexed, strongly complexed, mineral bound) to be assessed. Understanding the fractionation of mercury into these different classes provides an understanding of the potential solid-phase forms of mercury and, therefore, its propensity to enter the dissolved phase, be transported, and methylated. Eight transect locations (BCT5, BCT6, BCT7, BCT8, BCT9, BCT11, BCT12B, and BCT14), as well as a reference site (HCTREF [HCK 20.6]), were sampled for mercury speciation/sequential extraction analysis. Sufficient mass of solid material from the channel sediment (where available), creek bank soil, and floodplain soil at each selected transect for mercury speciation/sequential extraction was composited into samples from each of the three representative media types (i.e., three composite samples per transect). Sediment was collected to an approximate depth of 0.5 ft and passed through a 1-mm sieve until adequate sample volume was achieved. For creek bank soil, one sample was collected from each side of the bank by removing the upper 0.5 ft of bank soil surface just above the water level and was composited. For floodplain soil, samples were collected from the upper 0.5 ft on each side of the creek to generate a composite sample representing both sides of the floodplain. Planned sequential extraction for channel sediment at locations BCT5 and BCT6 and the reference site HCTREF (HCK 20.6) was not possible due to lack of fine-grained sediment at these locations.

## 2.2 CHANGES FROM THE BEAR CREEK VALLEY MERCURY SOURCES REMEDIAL SITE EVALUATION SAMPLING AND ANALYSIS PLAN

As described in the BCV Mercury Sources RSE SAP, transect locations were adjusted based on field conditions and sampling viability. The following locations were moved during the site reconnaissance prior to field implementation:

- BCT1 and BCT2 were moved further downstream because the beaver dams previously viewed at these locations were no longer in place.
- BCT15 was moved further downstream due to accessibility issues.

During the field sampling, the following samples were not collected (Table 2.2):

- Only particle size analysis samples were collected from channel sediments at BCT1, BCT2, BCT5, BCT6, and the reference site HCTREF (HCK 20.6) due to limited fine-grained sediment observed at these locations. Planned sequential extraction of channel sediment at locations BCT5 and BCT6 and the reference site HCTREF (HCK 20.6) was not performed due to these fine sediment limitations.
- Due to ponding at locations BCT7, BCT8, and BCT9, upper and lower creek bank soil was not divided into separate samples, and a single bank soil sample was collected.

## 3. BEAR CREEK VALLEY MERCURY SOURCES REMEDIAL SITE EVALUATION RESULTS

The following sections summarize results of sampling and analyses conducted for the BCV mercury sources RSE. Figure 2.1 shows the BCV mercury sources RSE transect sampling locations. Table 3.1 lists the sample location names and describes the location habitat characteristic with respect to association with open channel flow condition versus beaver ponds. Appendix B contains photographs of transect locations and substrate at each location.

Table 3.1. Sample location site characteristics

Location identifier	Site characteristic
BCT1	Open channel
BCT2	Open channel
BCT3	Open channel downstream of beaver dam/pond
BCT4	In beaver pond
BCT5	Open channel at washed-out small beaver dam
BCT6	Open channel upstream of former beaver pond
BCT7	Channel immediately downstream of beaver dam complex
BCT8	In large beaver pond downstream of Reeves Road
BCT9	In large beaver pond upstream of Reeves Road
BCT10	Open channel in former beaver pond
BCT11	Open channel upstream of BCK 9.9
BCT12A	Open channel upstream of NT-3
BCT12B	Open channel downstream of NT-3
BCT13	Open channel upstream of closed access road to HCDA (depositional due to culvert inflow restriction)
BCT14	Open channel downstream of BCK 12.34
BCT15	Open channel of NT-1
HCTREF	Open channel of reference site Hinds Creek

BCK = Bear Creek kilometer

BCT = Bear Creek transect

HCDA = Hazardous Chemical Disposal Area

HCTREF = Hinds Creek transect reference site NT = North Tributary

#### 3.1 SURFACE WATER DATA

Field parameters were measured in surface water at each transect location (Table 3.2). Table 3.2 includes the average daily flow rate measured at the Bear Creek integration point (BCK 9.2), which is the most downstream continuous flow monitoring station on Bear Creek. Surface water samples were collected on the dates that sediment and soil samples were collected at each sample transect location (December 2023 through February 2024) and, consequently, temporal variation exists in the dataset. Surface water physicochemical conditions indicate the stream water is well oxygenated, has positive electrochemistry (oxidation-reduction potential [redox]), and is near-neutral in pH. The longitudinal conductivity profile reflects the S-3 Ponds groundwater plume influx at the headwater (near transect BCT15), with generally decreasing conductivity in downstream samples. Because of the wide range of flow conditions during the initial sampling, surface water in Bear Creek was resampled on a single day in April 2024 to measure field parameters and to analyze total and dissolved mercury and methylmercury. Field parameters measured in the April 2024 sample data are included in Table 3.3. The daily average flow rate at BCK 9.2 was 3909 L/min on April 16, 2024.

Table 3.2. Bear Creek mercury sources RSE transects surface water field parameters from the initial sampling event

Location	Date	Conductivity (µmho/cm)	Dissolved oxygen (ppm)	pH (units)	Redox (mV)	Temperature (°C)	Turbidity (NTU)	BCK 9.2 daily average flow (L/min)
BCT15-SW	01/03/24	669	13.36	6.96	182.8	4.2	2	1331
BCT14-SW	12/28/23	534	12.11	7.22	154	9.6	6	4199
BCT13-SW	01/31/24	483	11.01	8.00	149	7.8	6	4796
BCT12A-SW	01/29/24	440	10.16	7.85	140	8.8	7	6892
BCT12B-SW	01/29/24	402	9.79	7.58	108	8.1	9	6892
BCT11-SW	01/31/24	323	10.06	8.12	140	7.8	6	4796
BCT10-SW	01/31/24	332	10.66	7.81	161	9.8	9	4796
BCT9-SW	02/05/24	263	9.9	8.11	139	9.9	5	2726
BCT8-SW	02/05/24	308	11.14	8.01	114	8.9	6	2726
BCT7-SW	02/07/24	338	11.75	8.29	92	5.7	13	2594
BCT6-SW	02/01/24	194	10.54	8.14	118	7.4	6	4419
BCT5-SW	02/01/24	143	9.87	7.78	133	7.4	7	4419
BCT4-SW	01/08/24	230	10.61	8.29	149	7.6	5	1258
BCT3-SW	01/08/24	239	11.76	7.74	179	6.1	4	1258
BCT2-SW	01/04/24	268	11.88	7.73	170.5	6.0	4	1262
BCT1-SW	12/18/23	355.6	10.56	8.22	161	8.2	3	1837
HCTREF-SW	02/02/24	269	10.94	8.07	96	5.9	6	4254

BCK = Bear Creek kilometer

BCT = Bear Creek transect

HCTREF = Hinds Creek transect reference site

NTU = nephelometric turbidity unit

ppm = parts per million redox = oxidation-reduction potential

RSE = remedial site evaluation

SW = surface water

Table 3.3. Bear Creek mercury sources RSE transects surface water field parameters, April 2024

Location	Date	Conductivity (µmho/cm)	Dissolved oxygen (ppm)	pH (units)	Redox (mV)	Temperature (°C)	Turbidity (NTU)
BCT15-SW	04/16/24	556	9.11	7.14	230.6	17.3	9
BCT14-SW	04/16/24	684	8.80	7.46	235.3	16.2	5
BCT13-SW	04/16/24	460	9.23	7.82	206.5	15.9	7
BCT12A-SW	04/16/24	393	9.49	7.96	199.1	15.7	4
BCT12B-SW	04/16/24	349	9.36	7.91	211.1	15.6	7
BCT11-SW	04/16/24	333	9.41	7.96	222.6	15.9	6
BCT10-SW	04/16/24	295	9.59	7.82	219	15.4	5
BCT9-SW	04/16/24	278	7.97	7.56	229.5	16.7	6
BCT8-SW	04/16/24	292	9.83	7.74	15.4	16.8	6
BCT7-SW	04/16/24	283	6.59	7.64	24.1	18.1	9
BCT6-SW	04/16/24	239	8.65	7.69	28.6	16.4	4
BCT5-SW	04/16/24	240	7.60	7.71	19.9	16.4	6
BCT4-SW	04/16/24	232	8.48	7.84	13.2	16.8	6
BCT3-SW	04/16/24	278	8.10	7.66	14.5	16.6	1
BCT2-SW	04/16/24	240	8.18	7.95	-1.7	19.2	5
BCT1-SW	04/16/24	298	10.96	8.02	-7.9	19.0	5
HCTREF-SW	04/17/24	289	9.50	8.10	244.1	18.2	5

BCT = Bear Creek transect

HCTREF = Hinds Creek transect reference site

NTU = nephelometric turbidity unit ppm = parts per million

redox = oxidation-reduction potential

RSE = remedial site evaluation

SW = surface water

Table 3.4 includes analyte results for total and dissolved mercury and methylmercury, nitrate, phosphorus, sulfate, sulfide, TOC, and DOC from the initial surface water sampling event. A comprehensive surface water chemical data summary is included in Appendix C.

Samples collected in this investigation show Bear Creek surface water mercury concentrations were less than the 51-ng/L AWQC level, with the highest measured total mercury concentration of 22 ng/L at BCT13 (dissolved mercury at BCT13 was 3.8 ng/L). Sample transect BCT13 is located upstream of NT-3 that drains the former BYBY, which was the source of significant mercury discharge into Bear Creek prior to its remediation by excavation in the early 2000s. Measured surface water mercury concentrations decrease downstream from BCT13; in the most downstream sample locations (BCT1 and BCT2), total and dissolved mercury are only slightly greater than values at the Hinds Creek reference site.

Surface water TOC and DOC are highest in the Bear Creek headwater samples and decrease in the downstream direction to levels similar to those measured in Hinds Creek.

Nitrate is highest (84.8 mg/L) at the Bear Creek headwater (BCT15) where groundwater from the S-3 Ponds nitrate plume emerges into NT-1. Nitrate concentrations decrease with distance downstream from the headwater source; in the most downstream sample transects (BCT1 and BCT2), the nitrate concentrations are less than those measured in Hinds Creek.

Phosphorus, a nutrient to aquatic and terrestrial plants, was detected in BCT12B at a concentration of 0.037 J mg/L; however, phosphorous was not detected in any other Bear Creek or Hinds Creek surface water samples at the 0.018-mg/L detection limit.

Sulfide, a factor in microbial methylation of mercury, was not detectable in any of the surface water samples at the 0.033-mg/L detection limit. Given the high dissolved oxygen and redox values of approximately 100 mV and greater, the absence of sulfide is expected.

Figure 3.1 is a graph of the mercury and methylmercury (total and dissolved concentrations) in the longitudinal transect from BCT15 downstream to BCT1, with the HCTREF data shown at the extreme right. From the highest measured mercury concentration (22 ng/L) at BCT13, the total mercury concentrations decrease downstream. Dissolved mercury maximum concentrations (4.3 ng/L) occur both upstream and downstream of NT-3 at BCT12A and BCT12B, with decreasing concentrations downstream. Methylmercury exhibits a more complex signature.

Table 3.5 includes the results of the April 2024 surface water sampling event for total and dissolved mercury and methylmercury. Figure 3.2 presents the April 2024 data for total and dissolved mercury and methylmercury graphically. In the April 2024 sampling event, the highest concentrations of dissolved mercury and methylmercury occurred at BCT13 near the entry to the former HCDA and upstream of NT-3. At BCT7, total mercury exhibited a concentration spike that was only weakly reflected in the dissolved mercury data. The total mercury concentration spike is suspected to reflect suspended solids in the sample. In addition to the highest total and dissolved methylmercury concentration at BCT13, elevated concentrations were measured in surface water at BCT9, BCT8, and BCT7, which are associated with the beaver pond complex near the Reeves Road crossing of Bear Creek.

Figure 3.3 shows the ratios of dissolved and total methylmercury to mercury for the April 2024 sampling event. The ratio generally increases in the downstream direction to a high of nearly 18% dissolved methylmercury at BCT2 and a high of approximately 7% total methylmercury at BCT1.

Figure 3.4 shows the nitrate, sulfate, TOC, and DOC surface water concentrations from the initial sampling event at Bear Creek and the Hinds Creek reference site.

Table 3.4. Summary of selected analytes in surface water at BCT locations collected from the initial sampling event

			ВСТ	T15		ВСТ	14		ВСТ	13		BCT	12A		BCT1	2B		ВСТ	<u></u>		ВСТ	10
Chemical name	Units	Filtered	Result <sup>a</sup>		Filtered	Resulta	Lab qual	Filtered	Result <sup>a</sup>	Lab qual	Filtered	Resulta	Lab qual	Filtered	Result <sup>a</sup>	Lab qual	Filtered	Resulta	Lab qual	Filtered	Result <sup>a</sup>	Lab qual
Dissolved organic carbon	mg/L	F	2.62		F	2.7		F	2.04		F	2.36		F	2.212.76		F	1.86		F	1.56	
Total organic carbon	mg/L	UF	2.67		UF	2.62		UF	2.27		UF	2.34		UF	2.59		UF	1.79		UF	1.49	
Mercury	ng/L	UF	1.4		UF	6.9		UF	22		UF	13		UF	13		UF	6.7		UF	4.5	
Mercury	ng/L	F	0.64		F	3.2		F	3.8		F	4.3		F	4.3		F	1.9		F	1.8	
Methylmercury	ng/L	UF	0.066	J	UF	0.17		UF	0.088		UF	0.064	J	UF	0.098		UF	0.063	J	UF	0.051	J
Methylmercury	ng/L	F	0.05	J	F	0.076	J	F	0.074	J	F	0.072	J	F	0.075	J	F	0.043	J	F	0.064	J
Nitrate/Nitrite as nitrogen	mg/L	UF	84.8		UF	19.4		UF	11.3		UF	5.35		UF	3.56		UF	3.48		UF	1.83	
Phosphorous	mg/L	UF	0.018	U	UF	0.037	J	UF	0.018	U	UF	0.018	U									
Sulfate	mg/L	UF	31.5		UF	33.4		UF	26.3		UF	21.7		UF	20.9		UF	20.6		UF	15.9	
Sulfide	mg/L	UF	0.033	U	UF	0.033	U	UF	0.033	U												
	ВСТ9		Г9		BC	Г8		BC	Γ <b>7</b>		BC	Г6		ВСТ	75		BC'	Γ4		ВСТ	Γ3	
Chemical name	Units	Filtered	Result <sup>a</sup>	Lab qual	Filtered	Resulta	Lab qual	Filtered	Result <sup>a</sup>	Lab qual												
		运			臣			臣			运			臣			运			臣		
Dissolved organic carbon	mg/L	F	1.57		F	1.61		F	1.5		F	1.3		F	1.34		F	1.64		F	1.57	
Total organic carbon	mg/L	UF	1.81		UF	1.46		UF	1.4	J	UF	1.26		UF	1.24		UF	1.61		UF	1.64	
Mercury	ng/L	UF	2.3		UF	2.7		UF	4.1	J	UF	2.2		UF	2.5		UF	1.3		UF	1.2	
Mercury	ng/L	F	0.81		F	0.94		F	1.1	J	F	0.96		F	0.8		F	0.65		F	0.52	
Methylmercury	ng/L	UF	0.084		UF	0.076	J	UF	0.14		UF	0.022	UJ	UF	0.022	UJ	UF	0.079	J	UF	0.071	J
Methylmercury	ng/L	F	0.073	J	F	0.09		F	0.09		F	0.031	J	F	0.023	J	F	0.064	J	F	0.048	J
Nitrate/Nitrite as nitrogen	mg/L	UF	2		UF	1.91		UF	1.59		UF	0.735		UF	0.695		UF	1.05		UF	0.994	
Phosphorous	mg/L	UF	0.018	U	UF	0.018	U	UF	0.018	U												
Sulfate	mg/L	UF	18.8		UF	19.3		UF	16.6		UF	9.62		UF	9.6		UF	31.6		UF	30.7	
Sulfide	mg/L	UF	0.033	U	UF	0.033	U	UF	0.033	U												
			BC	Γ2		BC	Γ1		НСТІ	REF												
Chemical name	Units	Filtered	Result <sup>a</sup>	Lab qual	Filtered	Result <sup>a</sup>	Lab qual	Filtered	Result <sup>a</sup>	Lab qual												
Dissolved organic carbon	mg/L	F	1.48		F	1.85		F	1.42		•											
Total organic carbon	mg/L	UF	1.46		UF	1.79		UF	1.15													
Mercury	ng/L	UF	0.99		UF	1.1		UF	0.91													
Mercury	ng/L	F	0.47	J	F	0.57		F	0.39	J												
Methylmercury	ng/L	UF	0.057	J	UF	0.036	J	UF	0.023	J												
Methylmercury	ng/L	F	0.06	J	F	0.044	J	F	0.025	J												
Nitrate/Nitrite as nitrogen	mg/L	UF	0.59		UF	0.555		UF	0.85													
Phosphorous	mg/L	UF	0.018	U	UF	0.018	U	UF	0.018	U												
Sulfate	mg/L mg/L	UF	29.2		UF	30.7		UF	8.04													
Sulfide	mg/L	UF	0.033	U	UF	0.033	U	UF	0.033	U												
Samue	111g/L	i OI	0.055		, 01	0.055	U	i OI	0.055	<u> </u>												

# Table 3.4. Summary of selected analytes in surface water at BCT locations collected from the initial sampling event (cont.)

<sup>a</sup>Result is the maximum of the filtered and unfiltered samples. -- Indicates no qualifier; detection.

-- Indicates no qualifier; detection.

BCT = Bear Creek transect
F = filtered sample
HCTREF = Hinds Creek transect reference site
J = estimated value
Lab = laboratory
qual = qualifier
U = not detected at reported quantitation limit
UF = unfiltered sample
UJ = not detected at estimated value




Table 3.5. Surface water total and dissolved mercury and methylmercury, April 2024 sampling event

				Mercury	(ng/L)			Methylmer	cury (ng/I	٦)		]	Ratios	
Location identifier	Sample	Collection	Т	otal	Dis	solved	1	otal	Dis	solved	Per disso		Methylme	ercury/Mercury
identifier	type	date	Result	Lab qualifier	Result	Lab qualifier	Result	Lab qualifier	Result	Lab qualifier	Total mercury	Methyl- mercury	Total	Dissolved
HCTREF-SW	REG	04/17/24	1		0.33	J	0.042	J	0.053	J	33%	126%	4%	16%
BCT15-SW	REG	04/16/24	5.1		1		0.066	J	0.069	J	20%	105%	1%	7%
BCT14-SW	REG	04/16/24	8.4		1.8		0.085	J	0.058	J	21%	68%	1%	3%
BCT14-SW	FR	04/16/24	8.1		1.9		0.078	J	0.042	J	23%	54%	1%	2%
BCT13-SW	REG	04/16/24	9.8		1.8		0.27	J	0.15	J	18%	56%	3%	8%
BCT12B-SW	REG	04/16/24	7.9		1.8		0.16	J	0.1	J	23%	63%	2%	6%
BCT12B-SW	FR	04/16/24	7.7		1.8		0.16	J	0.12	J	23%	75%	2%	7%
BCT12A-SW	REG	04/16/24	4.2		1.2	UJ	0.12	J	0.022	U	29%	18%	3%	2%
BCT11-SW	REG	04/16/24	4.5		1.4		0.047	J	0.034	J	31%	72%	1%	2%
BCT10-SW	REG	04/16/24	2.6		0.87		0.051	J	0.022	UJ	33%	43%	2%	3%
BCT9-SW	REG	04/16/24	2.1		0.74		0.089	J	0.037	J	35%	42%	4%	5%
BCT8-SW	REG	04/16/24	2.1		0.7		0.13	J	0.085	J	33%	65%	6%	12%
BCT7-SW	REG	04/16/24	4.2		0.87		0.17	J	0.084	J	21%	49%	4%	10%
BCT6-SW	REG	04/16/24	1.6		0.42	J	0.078	J	0.059	J	26%	76%	5%	14%
BCT5-SW	REG	04/16/24	1.5		0.44	J	0.081	J	0.032	J	29%	40%	5%	7%
BCT4-SW	REG	04/16/24	1.9		0.51		0.083	J	0.054	J	27%	65%	4%	11%
BCT3-SW	REG	04/16/24	1.8		0.47	J	0.098	J	0.052	J	26%	53%	5%	11%
BCT2-SW	REG	04/16/24	1.8		0.44	J	0.096	J	0.077	J	24%	80%	5%	18%
BCT1-SW	REG	04/16/24	1.6		0.54		0.11	J	0.083	J	34%	75%	7%	15%
BCV maximum			9.8		1.9		0.27		0.15		35%	105%	7%	18%
BCV average			4.3		1.0		0.11		0.07		27%	61%	3.5%	7.9%
Standard error			0.68		0.13		0.01		0.01		1.2%	4.6%	0.5%	1.1%

0-0

## Table 3.5. Surface water total and dissolved mercury and methylmercury, April 2024 sampling event (cont.)

-- Indicates no qualifier; detection.

BCT = Bear Creek transect

BCV = Bear Creek Valley

FR = field replicate sample HCTREF = Hinds Creek transect reference site


J = estimated value.

Lab = laboratory

REG = regular sample SW = surface water

U = not detected at reported quantitation limit UJ = not detected at estimated value

Figure 3.2. Total and dissolved mercury and methylmercury for the April 2024 surface water sampling event, upstream to downstream.



BCT = Bear Creek transect HCTREF = Hinds Creek transect reference site Hg = mercury MeHg = methylmercury SW = surface water

Figure 3.3. Total and dissolved methylmercury to mercury ratios for the April 2024 surface water sampling event, upstream to downstream.




Figure 3.4. Concentration graphs for selected analytes in Bear Creek (upstream to downstream) and Hinds Creek reference site surface water, original sampling event dates.

### 3.2 FLOODPLAIN SOIL, CREEK BANK SOIL, AND CHANNEL SEDIMENT DATA

#### 3.2.1 Soil and Sediment Particle Size

Soil and sediment samples were collected for particle size analysis at each sample transect location. Lower bank soil samples were not obtained at BCT7, BCT8, and BCT9 due to beaver ponds that limited the height of the banks (Table 2.2). Soils at all locations tend to be silt and clay dominated, while channel sediments are more evenly graded and include medium and fine sand along with silt and generally little clay-size material. Particle size results are included in Table 3.6 and are shown graphically in Figure 3.5. Particle size analysis reports provided by S&ME Inc., are included in Appendix D.

#### 3.2.2 Soil and Sediment Chemical Characteristics

Samples of floodplain soil, creek bank soil, and channel sediment were collected with the following exceptions: lower bank soil was not available at sample locations BCT7, BCT8, and BCT9; and fine-grained channel sediment was not available at BCT1, BCT2, BCT5, BCT6, and HCTREF (Figure 2.1 and Table 2.2).

Table 3.7 includes results for selected analytes measured in the floodplain and creek bank soils and channel sediment at each transect location. Data are arranged in columns by location from upstream to downstream and in rows by floodplain soil, creek bank soil (upper and lower intervals), and channel sediment to allow comparison of the vertical relationships of analyte concentrations at each site. Figures 3.6 through 3.9, respectively, present graphs of the chemical constituents included in Table 3.7 for the floodplain soil, upper creek bank soil, lower creek bank soil, and channel sediment.

Mercury concentrations in the sampled BCV soils and channel sediment are comparatively low. (Note the laboratory reported mercury and methylmercury concentrations in soil and sediment in units of  $\mu g/kg$ , equivalent to parts per billion [ppb]). The maximum measured mercury concentration in floodplain soil was 3500  $\mu g/kg$  at BCT12A, upstream of the NT-3 confluence with Bear Creek. The maximum measured mercury concentration in creek bank soil was 7100  $\mu g/kg$  at BCT13, located near the former HCDA entrance. The maximum measured mercury concentration in channel sediment was 530  $\mu g/kg$  at BCK12A, upstream of the confluence of NT-3 with Bear Creek and downstream of the HCDA.

The highest methylmercury concentrations in the sampled BCV soils and channel sediment are 1.7  $\mu$ g/kg in floodplain soil at BCT12B, downstream of the NT-3 confluence; 0.87  $\mu$ g/kg in creek bank soil at BCT7; and 0.82  $\mu$ g/kg in channel sediment at BCT9 in a beaver pond.

TOC in BCV soil and channel sediment is variable, with generally higher concentrations in the Bear Creek headwaters area than farther downstream. Floodplain soil samples exhibit higher TOC than creek bank soils. Channel sediment TOC concentrations are typically lower than those of the floodplain and creek bank soils.

None of the sediment sample locations downstream of Reeves Road that had sediment suitable for sieving for analysis contained mercury at levels greater than the threshold effect concentration (MacDonald et al. 2000).

Figure 3.10 shows the methylmercury to total mercury ratios for the Bear Creek channel sediment (where samples were available) compared with the methylmercury to total mercury ratios for surface water samples collected on April 16, 2024. The graphs demonstrate generally increasing ratios in the downstream flow direction. Annotations on the graphs show increased methylmercury to total mercury ratios in channel sediment and surface water in and downstream of beaver pond-affected stream reaches.

Table 3.6. Bear Creek and Hinds Creek soil and sediment particle size data

Location	Sample description	Maximum particle size	Percent gravel (4.75 to 75 mm) (%)	Percent coarse sand (2 to 4.75 mm) (%)	Percent medium sand (0.425 to 2 mm) (%)	Percent fine sand (0.075 to 0.425 mm) (%)	Percent silt (0.002 to 0.075 mm) (%)	Percent clay (<0.002 mm) (%)	Percent moisture (%)
			Floodplain s	soil					
BCT1-FP	Dark Brown Clay	#20	0	00	1	10	80.6	8.4	22
BCT2-FP	Light Brown Clay with Sand	#4	0	1	4	20	66.6	8.4	19.9
BCT3-FP	Tan Clay with Sand	3/8 in.	2	2	6	13	65.2	11.8	20.1
BCT4-FP	Brown Sandy Clay	#10	0	00	33	39	49.9	8.1	25.2
BCT5-FP	Brown Silty Sand	3/4 in.	0	1	12	46	30.7	10.3	23.9
BCT6-FP	Brown Silt with Sand	3/4 in.	44	2	3	17	61.2	12.8	21.8
BCT7-FP	Brown Sandy Silt	3/8 in.	0	0	3	37	50	10	36.4
BCT8-FP	Brown Silty Sand with Gravel	3/4 in.	17	77	14	20	34.3	7.7	36.1
BCT9-FP	Brown Clay with Sand	3/8 in.	0	2	5	15	59.9	18.1	28.1
BCT10-FP	Brown Sandy Clay	3/8 in.	11	0	6	33	52.8	7.2	24
BCT11-FP	Brown Clay with Sand	3/8 in.	0	1	1	23	63.1	11.9	22.7
BCT12A-FP	Brown Clay with Sand	3/8 in.	0	1	7	13	65	14	25.3
BCT12B-FP	Brown Sandy Clay	3/4 in.	4	4	14	13	52.5	12.5	23.5
BCT12B-FP	Brown Sandy Clay	3/4 in.	2	2	16	18	50.5	11.5	20
BCT13-FP	Brown Sandy Clay	#4	0	11	10	34	40	15	27.3
BCT14-FP	Brown Clay with Sand	#10	0	00	6	13	65	16	28.6
BCT14-FP	Brown Clay with Sand	#4	0	0	6	15	63.9	15.1	28.7
BCT15-FP	Brown Clay with Sand	3/4 in.	2	2	4	18	60.6	13.4	33.3
HCTREF-FP	Brown Sandy Silt	3/4 in.	6	1	13	24	48.1	7.9	20.9
		Upp	er creek bai	nk soil					
BCT1-BSU	Dark Brown Clay with Sand	#20	0	0	1	23	66.6	9.4	22.4
BCT2-BSU	Brown Clay with Sand	#4	0	00	2	14	70.3	13.7	20
BCT3-BSU	Light Brown Clay with Sand	3/8 in.	66	3	66	3	68.1	13.9	18.2
BCT4-BSU	Brown Sandy Clay	#10	0	0	4	34	53.6	8.4	23.4
BCT5-BSU	Brown Sandy Silt	3/8 in.	2	3	7	36	40.3	11.7	19.6

Table 3.6. Bear Creek and Hinds Creek soil and sediment particle size data (cont.)

Location	Sample description	Maximum particle size	Percent gravel (4.75 to 75 mm) (%)	Percent coarse sand (2 to 4.75 mm) (%)	Percent medium sand (0.425 to 2 mm) (%)	Percent fine sand (0.075 to 0.425 mm) (%)	Percent silt (0.002 to 0.075 mm) (%)	Percent clay (<0.002 mm) (%)	Percent moisture (%)
BCT6-BSU	Brown Sandy Silt	3/4 in.	3	2	6	31	47.1	10.9	20.6
BCT7-BS	Brown Sandy Silt	3/4 in.	11	00	10	32	46.1	10.9	30.2
BCT8-BS	Brown Silty Sand	3/4 in.	13	10	14	25	31.5	6.5	29.7
BCT9-BS	Brown Sandy Silt	3/4 in.	5	4	10	20	48.8	12.2	28.5
BCT10-BSU	Brown Clay with Sand	3/8 in.	1	0	1	21	71.4	5.6	18.8
BCT11-BSU	Brown Clay with Sand	3/8 in.	0	1	2	20	64.9	12.1	20.9
BCT12A-BSU	Brown Clay with Sand	3/4 in.	5	2	3	10	65.5	14.5	22.3
BCT12B-BSU	Brown Clay with Sand	3/8 in.	44	2	4	7	68.9	14.1	20.3
BCT12B-BSU	Brown Sandy Clay	3/4 in.	5	2	5	16	59.2	12.8	21.1
BCT13-BSU	Grayish Brown Sandy Clay	3/8 in.	2	11	66	23	52.8	15.2	27.1
BCT14-BSU	Yellowish Brown Clay	#10	0	0	5	9	62.9	23.1	18.3
BCT14-BSU	Yellowish Brown Clay	#10	0	0	5	9	63.8	22.2	17.9
BCT15-BSU	Brown Clay with Sand	#4	0	11	3	19	64.5	12.5	34.1
HCTREF-BSU	Brown Sandy Silt	3/8 in.	1	2	6	36	45.3	9.7	21.3
		Lov	ver creek ba	nk soil					
BCT1-BSL	Dark Brown Clay with Sand	#4	11	11	6	17	65	10	23.9
BCT2-BSL	Brown Sandy Clay	#4	4	3	11	19	50.5	12.5	22.6
BCT3-BSL	Tan Clay	#10	0	0	2	7	78.6	12.4	21.6
BCT4-BSL	Light Brown Clay with Sand	#4	0	0	5	24	59.9	11.1	27.6
BCT5-BSL	Brown Clay with Sand	#4	0	0	2	24	60.6	13.4	22.7
BCT6-BSL	Brown Silty Sand	3/4 in.	1	1	13	39	33.6	12.4	23.2
BCT10-BSL	Brown Clay with Sand	3/8 in.	1	1	5	20	62.8	10.2	24.8
BCT11-BSL	Brown Clay with Sand	3/8 in.	2	1	4	22	60.7	10.3	23
BCT12B-BSL	Brown Sandy Clay	3/4 in.	1	1	5	24	53.3	15.7	21.2
BCT12B-BSL	Brown Clay	#10	0	0	2	4	78.9	15.1	21.8
BCT12A-BSL	Brown Sandy Clay	3/8 in.	1	1	5	24	52.4	16.6	26.9
BCT13-BSL	Brown Sandy Clay	3/4 in.	3	3	5	24	58.1	6.9	28.3

3-15

Table 3.6. Bear Creek and Hinds Creek soil and sediment particle size data (cont.)

Location	Sample description	Maximum particle size	Percent gravel (4.75 to 75 mm) (%)	Percent coarse sand (2 to 4.75 mm) (%)	Percent medium sand (0.425 to 2 mm) (%)	Percent fine sand (0.075 to 0.425 mm) (%)	Percent silt (0.002 to 0.075 mm) (%)	Percent clay (<0.002 mm) (%)	Percent moisture (%)
BCT14-BSL	Brown Clay	#10	0	0	3	12	59.6	25.4	19.3
BCT14-BSL	Light Brown Clay	#10	0	0	2	10	60	28	19.5
BCT15-BSL	Brown Sandy Clay	#4	0	1	5	25	57.8	11.2	39.3
HCTREF-BSL	Brown Sandy Silt	3/8 in.	1	3	7	35	40.8	13.2	35
		C	hannel sedir	nent					
BCT1-CH	Poorly Graded Gravel with Sand	3/4 in.	61	28	10	00	0	0	
ВСТ2-СН	Brown Silty Sand	#10	00	0	72	11	13.8	3.2	
ВСТ3-СН	Brown Sand with Silt	#10	0	00	57	36	3.3	3.7	27.9
ВСТ4-СН	Brown Silty Sand	#10	0	00	66	19	11.7	3.3	23.1
ВСТ5-СН	Brown Poorly Graded Sand with Silt	#4	0	11	75	15	7.2	1.8	
ВСТ6-СН	Brown Silty Sand	#4	0	0	61	10	26.3	2.7	
ВСТ7-СН	Grayish Brown Silt	#20	0	00	2	12	81.9	4.1	31.5
ВСТ8-СН	Brown Well-graded Sand with Silt	#10	0	0	67	25	4.2	3.8	29.4
ВСТ9-СН	Brown Silty Sand	3/8 in.	0	0	10	21	63.1	5.9	31.6
ВСТ10-СН	Brown Silty Sand	#10	0	00	44	35	18.5	2.5	23.8
BCT11-CH	Brown Well-graded Sand with Silt	3/8 in.	11	11	53	34	66	5	22.9
BCT12B-CH	Brown Poorly Graded Sand with Silt	#10	0	0	44	45	6.3	4.7	29.2
BCT12B-CH	Brown Silty Sand	#10	0	00	41	15	40.2	3.8	30.4
BCT12A-CH	Brown Silty Sand	#4	0	00	40	38	13.5	8.5	31.8
ВСТ13-СН	Brown Well-graded Sand with Silt	3/4 in.	2	00	55	35	2.7	5.3	30.3
BCT14-CH	Yellowish Brown Clay	#10	0	0	77	7	57	29	37
BCT14-CH	Yellowish Brown Clay	#10	0	0	6	4	60.5	29.5	38.2
BCT15-CH	Dark Brown Sandy Clay	#10	0	0	29	19	44.5	7.5	38.6
HCTREF-CH	Brown Well-graded Sand	3/4 in.	2	53	41	1	2.5	0.5	

<sup>--</sup> Indicates no qualifier; detection.

BCT = Bear Creek transect

BSL = lower creek bank soil

BSU = upper creek bank soil

CH = channel sediment

FP = floodplain soil HCTREF = Hinds Creek transect (reference site)




Figure 3.5. Bear Creek floodplain soil, creek bank soil, and channel sediment average particle size distributions.

This page intentionally left blank.

Table 3.7. Summary of selected analyte concentrations in BCTs floodplain soil, upper and lower creek bank soil, and channel sediment samples

Chemical name	Units -	BC	CT15	BC	CT14	ВС	CT13	BC	T12A	BC	Т12В	ВС	CT11	BC	CT10	B	СТ9	B	CT8
Chemical name	Units -	Result	Lab qual	Result	Lab qual	Result	Lab qual	Result	Lab qual	Result	Lab qual	Result	Lab qual						
								Flo	oodplain soil										
Mercury	μg/kg	470		740		3300		3500		3000		810	J	640		380		270	
Methylmercury	μg/kg	0.57		0.48		0.51		0.7		1.7		0.17		0.16	J	0.4		0.51	
Nitrate	mg/kg	15.8		2.82		1.5		2.57		3.78		2.69		1.56		3.99		2.59	
Nitrite	mg/kg	0.445	U	0.448	U	0.446	U	0.429	U	0.422	U	0.421	U	0.423	U	0.441	U	0.488	U
Phosphorous	mg/kg	270		296		244		291		248		223		213		227		250	
Sulfate	mg/kg	16.1		6.64		21.4	J	6		3.65	J	3.85	J	3.08	J	10.6		14.2	
Sulfide	mg/kg	19.2	J	15.2	J	22.2	J	32.6	J	23.4	J	30.4	J	30.7	J	40.9		59.1	
Total organic carbon average	mg/kg	26,600		39,200		25,500		33,600		31,900		15,800		20,200		18,900		24,200	
								Upper	creek bank s	oil									
Mercury	μg/kg	420		59		7100		860		1900		230	J	75		550		180	
Methylmercury	μg/kg	0.17		0.04	J	0.77	J	0.42		0.28		0.13		0.098	J	0.23		0.48	
Nitrate	mg/kg	29.2		0.395	U	1.35		1.33		1.71		2.03		0.403	U	4.4		0.889	J
Nitrite	mg/kg	0.473	U	0.395	U	0.438	U	0.397	U	0.413	U	0.402	U	0.403	U	0.429	U	0.442	U
Phosphorous	mg/kg	255		110		250		202		196		174		114		242		241	
Sulfate	mg/kg	27.8		11.9		25.7	J	3.29	J	3.78	J	4.29	J	2.48	J	9.36		9.28	
Sulfide	mg/kg	39.7		24	J	21.7	J	21.1	J	17.4	J	10.9	U	23.7	J	43.3		36	
Total organic carbon average	mg/kg	26,900		3790		25,000		13,400		10,500		11,200		4810		18,800		22,100	
								Lower	creek bank s	oil									
Mercury	μg/kg	320		41		1600		1600		1300		120	J	150		NA		NA	
Methylmercury	μg/kg	0.25		0.017	UJ	0.71	J	0.26		0.14		0.17		0.056	J	NA		NA	
Nitrate	mg/kg	88.5		23.1		1.7		1.2	J	1.73		2.18		1.28		NA		NA	
Nitrite	mg/kg	0.525	U	0.407	U	0.434	U	0.443	U	0.411	U	0.408	U	0.417	U	NA		NA	
Phosphorous	mg/kg	239		79		215		215		172		174		187		NA		NA	
Sulfate	mg/kg	31.9		74.9		26.8	J	6.02		4.72	J	7.87		6.35	J	NA		NA	
Sulfide	mg/kg	25.9	J	12	J	20	J	32.5	J	21.3	J	28.2	J	23.6	J	NA		NA	
Total organic carbon average	mg/kg	24,200		7950		15,900		16,600		12,700		8920		9000		NA		NA	
								Cha	nnel sedimen	t									
Mercury	μg/kg	430		83		340		530		330		160	J	95	J	440		110	
Methylmercury	μg/kg	0.031	J	0.042	J	0.19		0.4		0.23		0.1		0.27		0.82		0.17	
Nitrate	mg/kg	32.4		14.8		1.33	J	0.975	J	1.01	J	1.43		1.08	J	0.894	J	0.459	U
Nitrite	mg/kg	0.531	U	0.51	U	0.438	U	0.445	U	0.461	U	0.414	U	0.429	U	0.445	U	0.459	U
Phosphorous	mg/kg	245		106		259		185		196		253		375		138		360	
Sulfate	mg/kg	34.3		36.3		18.3		14		13		9.4		33.7		35.5		25	
Sulfide	mg/kg	30.7	J	14	J	17.8	J	12.7	U	12.6	U	11.1	U	12.8	J	36.4		36.7	
Total organic carbon average	mg/kg	21,200		4420		6760		7710		7160		6320		8390		10,800		5080	

Table 3.7. Summary of selected analyte concentrations in BCTs floodplain soil, upper and lower creek bank soil, and channel sediment samples (cont.)

Chemical name	Units	I	BCT7	E	ВСТ6	E	BCT5	E	BCT4	E	ВСТ3	В	SCT2	]	BCT1	HC'	TREF
Chemical name	Onits	Result	Lab qual	Result	Lab qual	Result	Lab qual	Result	Lab qual	Result	Lab qual	Result	Lab qual	Result	Lab qual	Result	Lab qual
							Floc	dplain soi	l								
Mercury	μg/kg	220		62		160		180		86		110		150		40	
Methylmercury	μg/kg	0.6		0.13		0.22		0.37		0.11		0.49		0.28		0.1	
Nitrate	mg/kg	2.79		1.13	J	4.47		2.09		0.938	J	1.76		1.97		0.841	J
Nitrite	mg/kg	0.479	U	0.395	U	0.832	J	0.434	U	0.402	U	0.39	U	0.42	U	0.41	U
Phosphorous	mg/kg	228		158		207		200		162		206		162		220	
Sulfate	mg/kg	10.7		4.42	J	7.66		6.12		6.13		6.99		4.09	J	4.7	J
Sulfide	mg/kg	15.4	J	40		22.9	J	11.6	U	30.3		21.6	J	25.7	J	24.4	J
Total organic carbon average	mg/kg	25,800		8810		25,200		19,400		9760		17,000		14,600		17,400	
							Upper o	reek bank	soil								
Mercury	μg/kg	200		34		200		100		32		120		150		28	
Methylmercury	μg/kg	0.87		0.017	UJ	0.094		0.41		0.025	J	0.24		0.4		0.016	UJ
Nitrate	mg/kg	1.38		1.04	J	0.969	J	1.42		1.09	J	1.43		1.96		1.06	J
Nitrite	mg/kg	0.449	U	0.394	U	0.391	U	0.404	U	0.4	U	0.404	U	0.423	U	0.392	U
Phosphorous	mg/kg	189		145		146		154		158		136		179		113	
Sulfate	mg/kg	11.9		3.46	J	2.8	J	7.66		4.05	J	6.99		3.98	J	9.5	
Sulfide	mg/kg	15	J	26.3	J	22.1	J	17.4	J	18	J	34.9		20.7	J	21.8	J
Total organic carbon average	mg/kg	16,700		5990		10,400		11,800		6640		7350		14,600		6850	
							Lower	reek bank	soil								
Mercury	μg/kg	NA		33		42		66		24	J	41		180		30	
Methylmercury	μg/kg	NA		0.018	UJ	0.018	UJ	0.18		0.033	J	0.062		0.26		0.04	J
Nitrate	mg/kg	NA		0.856	J	0.851	J	1.81		0.41	U	1.09	J	2.11		0.49	U
Nitrite	mg/kg	NA		0.425	U	0.411	U	0.45	U	0.41	U	0.405	U	0.408	U	0.49	U
Phosphorous	mg/kg	NA		133		144		181		144		159		176		184	
Sulfate	mg/kg	NA		10.1		6.28		11.6		16.1		20.8		6.88		8.56	
Sulfide	mg/kg	NA		27.6	J	43.6		31.3	J	23.6	J	11.3	U	15.9	J	31.1	J
Total organic carbon average	mg/kg	NA		5850		7840		10,700		5320		5550		11,200		12,300	
							Chan	nel sedime	nt								
Mercury	μg/kg	86		NA		NA		26	J	29	J	NA		NA		21ª	
Methylmercury	μg/kg	0.25		NA		NA		0.057	J	0.063	J	NA		NA		0.186a	
Nitrate	mg/kg	0.464	U	NA		NA		1.1	J	1.22	J	NA		NA		NA	
Nitrite	mg/kg	0.464	U	NA		NA		0.419	U	0.455	U	NA		NA		NA	
Phosphorous	mg/kg	131		NA		NA		340		239		NA		NA		NA	
Sulfate	mg/kg	20.6		NA		NA		10.4		22.1		NA		NA		NA	
Sulfide	mg/kg	13.1	J	NA		NA		27.6	J	24.4	J	NA		NA		NA	
Total organic carbon average	mg/kg	7250		NA		NA		2500		2440		NA		NA		$11,000^a$	

<sup>&</sup>lt;sup>a</sup>Data provided courtesy of Scott Brooks, Oak Ridge National Laboratory, Environmental Science Division.

Bold indicates channel sediment mercury concentration is greater than sediment threshold effect concentration (180 µg/kg) and is less than probable effect concentration (1060 µg/kg).

<sup>--</sup> Indicates no qualifier; detection.

BCT = Bear Creek transect
J = estimated value
Lab = laboratory
NA = not analyzed
qual = qualifier
U = not detected at reported quantitation limit
UJ = not detected at estimated value






Figure 3.6. BCT (upstream to downstream) and Hinds Creek floodplain soil chemical data.








Figure 3.9. BCT (upstream to downstream) channel sediment chemical data.




Figure 3.10. Bear Creek channel sediment and April 2024 surface water percent methylmercury, shown upstream to downstream.

3.3 BEAR CREEK VALLEY SOIL AND SEDIMENT MERCURY SEQUENTIAL EXTRACTIONS

Mercury sequential extractions were conducted on floodplain soil, creek bank soil, and channel sediment at select Bear Creek transect locations (BCT14, BCT12B, BCT11, BCT9, BCT8, BCT7, BCT6, and BCT5 and the reference site HCTREF) (Figure 2.1) using EPA Method SW846-3200M (modified). Because of very coarse sediment texture, or absence of sediment due to channel scour to bedrock, channel sediment was not suitable for sequential mercury extraction at locations BCT6, BCT5, and HCTREF. Table 3.8 includes the seven mercury extraction fractions and the relative ease of extraction (e.g., Fraction 0, volatile "gaseous" mercury, through Fraction 7, insoluble mercury contained in mineral crystal lattice positions requiring total dissolution of mineral phase for extraction).

Table 3.9 includes the results of the mercury sequential extraction analyses conducted on the Bear Creek transect samples. Measured mercury concentrations in each sample fraction are included along with sample result qualifiers and the percentage of the sum of measured total mercury in the sample. Table 3.9 also includes the average concentration of mercury in each extracted fraction for samples collected at each Bear Creek transect as well as the Hinds Creek reference site. Table 3.9 is organized with data from upstream sample transects to the left and progressing downstream to the right.

Table 3.8. Mercury sequential extraction method fractions and descriptions

Extraction step	Description and presumptive forms
F0 – Volatile Elemental Mercury (gas purging)	Volatile forms
F1 – Water Soluble Mercury (anoxic deionized water)	Water soluble
$F2-pH\ 2$ Soluble Mercury (0.1 M acetic acid + 0.01 M hydrochloric acid)	Low pH soluble mercury forms; human stomach acid equivalent
F3 – 1N Potassium Hydroxide Extractable Mercury	Organically bound mercury
F4 – 12N Nitric Acid Soluble Mercury	Strongly complexed, non-silicate- or non-sulfide-bound mercury
F5 – Aqua Regia Soluble Mercury (3:1 concentrated hydrochloric acid to concentrated nitric acid)	Sulfide-bound mercury
F6 – Mineral-bound Mercury (hydrofluoric, nitric, and hydrochloric acid mixture, high-pressure–high-temperature digestion)	Silicate- or aluminosilicate-bound mercury

F = fraction

M = molarity

N = normality

As summarized, the potassium hydroxide extracted fraction predominates in mercury concentration in most cases for Bear Creek floodplain soil, creek bank soil, and channel sediment.

The mercury sequential extraction results are shown graphically as the percentage extracted in each fraction and the sum of measured mercury (µg/kg or ppb) for floodplain soils (Figure 3.11), creek bank soils (Figure 3.12), and channel sediment (Figure 3.13). In the floodplain soil (Figure 3.11), the dominance of organic-associated mercury extracted in the F3 potassium hydroxide step is obvious in most samples. The Hinds Creek reference site exhibits more mineral-bound (F6) mercury fraction in the floodplain soils than any of the Bear Creek sites. Creek bank soils (Figure 3.12) also exhibit a high proportion of organic-associated mercury, except at the headwater sample location (BCT14) where more strongly bound fraction (F4) mercury dominates. At the Hinds Creek site, a large component of mineral-bound (F6) mercury in creek bank soil was also present, as it was in the floodplain soil. Where channel sediment samples were available (Figure 3.13), organic-associated mercury exhibits an apparent increasing trend in the downstream direction from the headwater (BCT14) to BCT7 located immediately downstream of the Reeves Road crossing of Bear Creek. As noted in Table 3.1, sample locations BCT8 and BCT9 are located within beaver ponds and location BCT7 is located immediately downstream of the BCT8 beaver pond. Channel sediment suitable for the sequential mercury extraction analysis was not available at locations BCT5 and BCT6 or at the Hinds Creek reference site.

Figures 3.14 through 3.22 show data graphs of the sequential mercury extraction results for floodplain soil, creek bank soil, and channel sediment (where available) for each sample location. These graphs portray the vertical distribution of the mercury forms at each sample location.

Table 3.9. Results of mercury sequential extraction analyses from floodplain soil, creek bank soil, and channel sediment at eight BCT locations and at the Hinds Creek reference site

			BCT14			BCT12B			BCT11			ВСТ9			ВСТ8	
Chemical name	Units	Result	Percent of sum	Lab qual												
					Floo	dplain soil										
Mercury (F0) – Volatile Elemental Mercury	μg/kg	0.18	0%	UJ	0.18	0%	UJ	0.17	0%	UJ	0.56	0%	J	0.22	0%	UJ
Mercury (F1) – Water Soluble Mercury	μg/kg	20	6%	J	54	3%	J	25	3%		13	6%		3.9	3%	J
Mercury (F2) – pH 2 Soluble Mercury	μg/kg	9.7	3%	J	40	3%	J	17	2%		3.3	2%	J	1.9	2%	J
Mercury (F3) – 1N Potassium Hydroxide Extractable Mercury	μg/kg	270	79%	J	850	53%	J	550	73%		160	73%		99	76%	
Mercury (F4) – 12N Nitric Acid Soluble Mercury	μg/kg	37	11%	J	560	35%	J	110	15%		31	14%		18	14%	J
Mercury (F5) – Aqua Regia Soluble Mercury	μg/kg	4.7	1%	J	53	3%	J	45	6%		11	5%		3.6	3%	J
Mercury (F6) – Mineral-bound Mercury	μg/kg	1.7	1%	J	2.6	0%	J	0.73	0%	J	1	1%	J	0.4	0%	J
Mercury (FS) <sup>a</sup> – Total Mercury by Summation	μg/kg	340	101%	J	1600	98%	J	750	100%		220	100%	J	130	98%	J
					Ba	ınk soil										
Mercury (F0) – Volatile Elemental Mercury	μg/kg	0.17	0%	UJ	0.17	0%	UJ	0.18	0%	J	0.18	0%	UJ	0.2	0%	UJ
Mercury (F1) – Water Soluble Mercury	μg/kg	1.1	0%	UJ	26	3%	J	7.1	4%		8.8	4%		2.8	3%	J
Mercury (F2) – pH 2 Soluble Mercury	μg/kg	1.1	0%	UJ	20	2%	J	3.5	2%	J	4.9	2%		2.1	3%	J
Mercury (F3) – 1N Potassium Hydroxide Extractable Mercury	μg/kg	2.2	0%	UJ	390	46%	J	120	75%		200	80%		65	77%	
Mercury (F4) – 12N Nitric Acid Soluble Mercury	μg/kg	17	81%	J	150	18%	J	15	9%	J	25	10%		12	14%	J
Mercury (F5) – Aqua Regia Soluble Mercury	μg/kg	3.1	15%	J	250	30%	J	9.3	6%		6.8	3%		3	4%	J
Mercury (F6) – Mineral-bound Mercury	μg/kg	1.3	6%	J	6.8	1%	J	0.93	1%	J	0.82	0%	J	0.36	0%	UJ
Mercury (FS) <sup>a</sup> – Total Mercury by Summation	μg/kg	21	102%	J	840	100%	J	160	98%	J	250	99%	J	85	100%	J
					Chann	el sediment										
Mercury (F0) – Volatile Elemental Mercury	μg/kg	7.4	14%	J	0.19	0%	J	0.52	1%	J	0.18	0%	UJ	0.2	0%	UJ
Mercury (F1) – Water Soluble Mercury	μg/kg	1.3	0%	UJ	6.5	3%		5.3	6%		8.6	4%		3.4	4%	J
Mercury (F2) – pH 2 Soluble Mercury	μg/kg	1.3	0%	UJ	5.6	3%	J	4.3	4%	J	4.8	2%		1.3	2%	J
Mercury (F3) – 1N Potassium Hydroxide Extractable Mercury	μg/kg	2.7	0%	UJ	99	45%		37	38%		170	71%		70	80%	
Mercury (F4) – 12N Nitric Acid Soluble Mercury	μg/kg	32	58%	J	20	9%		8.3	9%	J	43	18%		9.6	11%	J
Mercury (F5) – Aqua Regia Soluble Mercury	μg/kg	23	42%	J	85	39%		40	41%		11	5%		2.7	3%	J
Mercury (F6) – Mineral-bound Mercury	μg/kg	0.46	1%	J	0.91	0%	J	1.9	2%	J	1.2	1%	J	0.71	1%	J
Mercury (FS) <sup>a</sup> – Total Mercury by Summation	μg/kg	55	101%	J	220	99%	J	97	100%	J	240	99%	J	88	100%	J

Table 3.9. Results of mercury sequential extraction analyses from floodplain soil, creek bank soil, and channel sediment at eight BCT locations and at the Hinds Creek reference site (cont.)

			BCT7			BCT6			BCT5		Bear Cre	ek average		HCTREF	
Chemical name	Units	Result	Percent of sum	Lab qual	Result	Percent of sum	Lab qual	Result	Percent of sum	Lab qual	Average result	Average percent	Result	Percent of sum	Lab qual
					Floodpla	in soil									
Mercury (F0) – Volatile Elemental Mercury	μg/kg	0.2	0%	UJ	1.4	5%	J	0.17	0%	UJ	0.25	1%	0.22	1%	J
Mercury (F1) – Water Soluble Mercury	μg/kg	3.4	3%	J	2.5	8%	J	3.2	3%	J	15.6	4%	1.1	0%	UJ
Mercury (F2) – pH 2 Soluble Mercury	μg/kg	1.6	1%	J	1.2	0%	UJ	2	2%	J	9.4	2%	1.1	0%	UJ
Mercury (F3) – 1N Potassium Hydroxide Extractable Mercury	μg/kg	110	85%		26	84%		83	84%		269	76%	18	62%	
Mercury (F4) – 12N Nitric Acid Soluble Mercury	μg/kg	15	12%	J	5.9	0%	UJ	6.2	6%	J	97	13%	5.5	0%	UJ
Mercury (F5) – Aqua Regia Soluble Mercury	μg/kg	2.9	2%	J	2.5	8%	J	3.5	4%	J	16	4%	1.2	4%	J
Mercury (F6) – Mineral-bound Mercury	μg/kg	0.47	0%	J	0.42	1%	J	0.76	1%	J	1.0	1%	10	35%	J
Mercury (FS) <sup>a</sup> – Total Mercury by Summation	μg/kg	130	103%	J	31	102%	J	99	100%	J		101%	29	101%	J
					Bank	soil									
Mercury (F0) – Volatile Elemental Mercury	μg/kg	0.19	0%	UJ	0.18	0%	UJ	0.93	1%	J	0.1	0%	0.37	1%	J
Mercury (F1) – Water Soluble Mercury	μg/kg	3.3	3%	J	1.5	6%	J	4.4	5%	J	6.7	4%	2.8	9%	J
Mercury (F2) – pH 2 Soluble Mercury	μg/kg	1.3	1%	J	1.2	0%	UJ	1.6	2%	J	4.2	2%	1.1	0%	UJ
Mercury (F3) – 1N Potassium Hydroxide Extractable Mercury	μg/kg	86	78%		19	76%		68	79%		118.5	64%	13	39%	
Mercury (F4) – 12N Nitric Acid Soluble Mercury	μg/kg	14	13%	J	5.9	0%	UJ	6.8	8%	J	30	19%	5.3	0%	UJ
Mercury (F5) – Aqua Regia Soluble Mercury	μg/kg	2.2	2%	J	4.5	18%		4.4	5%		35.4	10%	1.2	4%	J
Mercury (F6) – Mineral-bound Mercury	$\mu g/kg$	0.34	0%	UJ	0.48	2%	J	0.72	1%	J	1.4	1%	16	49%	J
Mercury (FS) <sup>a</sup> – Total Mercury by Summation	$\mu g/kg$	110	97%	J	25	102%	J	86	100%	J		100%	33	100%	J
					Channel s	ediment									
Mercury (F0) – Volatile Elemental Mercury	<u>μ</u> g/kg	0.21	0%	UJ	NA			NA			1.4	2%	NA		
Mercury (F1) – Water Soluble Mercury	μg/kg	1.6	3%	J	NA			NA			4.25	3%	NA		
Mercury (F2) – pH 2 Soluble Mercury	μg/kg	1.6	3%	J	NA			NA			2.9	2%	NA		
Mercury (F3) – 1N Potassium Hydroxide Extractable Mercury	$\mu g/kg$	45	88%		NA			NA			70.2	54%	NA		
Mercury (F4) – 12N Nitric Acid Soluble Mercury	$\mu g/kg$	6.6	0%	UJ	NA			NA			18.8	17%	NA		
Mercury (F5) – Aqua Regia Soluble Mercury	$\mu g/kg$	2.4	5%	J	NA			NA			27.4	22%	NA		
Mercury (F6) – Mineral-bound Mercury	μg/kg	0.38	0%	UJ	NA			NA			0.86	1%	NA		
Mercury (FS) <sup>a</sup> – Total Mercury by Summation	μg/kg	51	99%	J	NA			NA				100%	NA		

"Mercury (FS) sum does not include Mercury (F0) aliquot because volatile mercury is measured on a separate sample aliquot. Laboratory reporting at a two-significant-figure level accounts for summed percentages greater than or less than 100%.

<sup>--</sup> Indicates no qualifier; detection or not applicable.

BCT = Bear Creek transect

F = fraction
HCTREF = Hinds Creek transect reference site

J = estimated value


Lab = laboratory

N = normality

NA = not analyzed


qual = qualifier

UJ = not detected at estimated value




BCT = Bear Creek transect HCTREF = Hinds Creek transect reference site Hg = mercury J = estimated value ppb = parts per billion UJ = not detected at estimated value

Figure 3.11. Results of Bear Creek floodplain soil sequential mercury extraction.



BCT = Bear Creek transect HCTREF = Hinds Creek transect reference site Hg = mercury J = estimated value ppb = parts per billion UJ = not detected at estimated value

Figure 3.12. Results of Bear Creek bank soil sequential mercury extraction.



BCT = Bear Creek transect Hg = mercury J = estimated value ppb = parts per billion UJ = not detected at estimated value

Figure 3.13. Results of Bear Creek channel sediment sequential mercury extraction.




Figure 3.14. Sequential mercury extraction results at location BCT5.




Figure 3.15. Sequential mercury extraction results at location BCT6.




Figure 3.16. Sequential mercury extraction results at location BCT7.




Figure 3.17. Sequential mercury extraction results at location BCT8.




Figure 3.18. Sequential mercury extraction results at location BCT9.






Figure 3.19. Sequential mercury extraction results at location BCT11.



\_\_\_\_

Figure 3.20. Sequential mercury extraction results at location BCT12B.



See Table 3.8 for definition of mercury extraction fractions

Hg = mercury J = estimated value ppb = parts per billion UJ = not detected at estimated value

Figure 3.21. Sequential mercury extraction results at location BCT14.




Figure 3.22. Sequential mercury extraction results at the Hinds Creek reference site (HCTREF).

### 3.3.1 Analyte Correlation Evaluations

Results of analyses of Bear Creek floodplain and bank soil samples (combined and separately) and Bear Creek stream sediment samples were evaluated for analyte pairwise and multivariate correlation. Correlation results are included in Appendix E. The pairwise correlation tables contain the correlation coefficient for each analyte pair and use a color scheme to reflect the degree of significance of each correlation pair. Combinations that result in correlations that are significant at the 0.05 significance level are highlighted. The pairwise correlations showed many strong and significant correlations among metals, including mercury.

# 3.4 BEAR CREEK VALLEY MERCURY COMPARISON TO LOWER EAST FORK POPLAR CREEK

From 1953–1983, use of mercury for lithium isotope separation processes at Y-12 created mercury-contaminated waste that was disposed in BCV waste disposal areas as well as large releases of mercury to the EFPC. Section 1.2.2 summarized information regarding mercury-contaminated waste disposals in BCV. The UEFPC Watershed and the BCV Watershed are separated by a watershed divide located on the eastern side of BCV (Figure 1.2).

Mercury concentrations in floodplain and creek bank soil and channel sediment in Bear Creek are much lower than those of the Lower East Fork Poplar Creek (LEFPC). The purposes of this comparison are to contrast the conditions between the two watersheds and to highlight some differences thought to contribute to apparent higher bioaccumulation rates for mercury in fish in Bear Creek. Data are not amenable to statistical comparisons due to discrepancies in data availability and sparse sample results from several studies that limit statistical test feasibility.

Sequential mercury extraction data for Bear Creek show a different partitioning of mercury than those reported for LEFPC (Crowther et al., 2021). Table 3.10 includes the average percentages of extraction fractions F3, F4, and F5 for channel sediment in both watersheds. Channel sediment in Bear Creek contains a much greater percentage (54.44%) of mercury associated with the organic component extraction step (F3) than is reported for LEFPC (4.63%). The fraction of mercury removed in extraction step F3 has been correlated with mercury methylation potential in aquatic sediments (ORNL/TM-2016/578). By extension of this concept to the conditions extant in Bear Creek, and the strong association of mercury with the F3 extraction step, its organic carbon association (Table 3.9 and Figures 3.11 through 3.13) may be a key factor in the observed Bear Creek mercury bioaccumulation.

Table 3.10. Comparison of channel sediment sequential mercury extraction steps F3, F4, and F5 averages in Bear Creek and LEFPC

Extraction step	Bear Creek	LEFPC <sup>a</sup>
F3 – 1N Potassium Hydroxide Extractable Mercury	54.44%	4.63%
F4 – 12N Nitric Acid Soluble Mercury	19.60%	18.20%
F5 – Aqua Regia Soluble Mercury (3:1 concentrated hydrochloric acid to concentrated nitric acid)	22.34%	76.39%

<sup>&</sup>quot;Averages calculated from combined data in Tables S6.3, S7.1, and S7.2 for three sampling locations in LEFPC reported in Crowther et al., 2021, "Supporting information for use of sequential extraction and mercury stable isotope analysis to assess remobilization of sediment-bound legacy mercury," *Environ. Sci.: Processes Impacts*, 23:756-775.

F = fraction

LEFPC = Lower East Fork Poplar Creek


N = normality

In addition to readily binding mercury, organic matter can serve as an electron donor to mercury-methylating bacteria. Bravo et al. (2017) found that, in boreal lakes, phytoplankton-derived organic compounds (relatively low molecular weight) enhance the mercury-methylation rates; whereas, methylation rates in lakes dominated with terrigenous organic compounds (relatively high molecular weights) had lower methylation rates but contained higher methylmercury concentrations.

# 4. WATER RESOURCES RESTORATION PROGRAM BEAR CREEK MONITORING RESULTS

#### 4.1 SURFACE WATER DATA

The WRRP began regular semiannual sampling of total mercury and methylmercury in surface water in 2011. These surface water data are collected approximately in sync with the Biological Monitoring and Abatement Program biota sampling in Bear Creek. Six sample locations are located in Zone 3 where continuous flow is measured, one is located in Zone 3 where flow is not measured (surface spring SS-4), and two are located farther downstream where flow is not measured coincident with sampling (BCK 4.6 and BCK 3.3) (Figure 4.1). While this dataset is not inclusive of all surface water mercury and methylmercury values for the time period, it is the Environmental Management Program dataset that provides same-day, snapshot data from stations arrayed along the length of Bear Creek, inclusive of key tributary and spring inputs to the Bear Creek mainstem. Appendix F provides the WRRP surface water total mercury and methylmercury longitudinal data plots for Bear Creek.




AWQC = ambient water quality criteria BCK = Bear Creek kilometer Hg = mercury HgT = total mercury Max = maximum NT = North Tributary SS = surface spring StDev = standard deviation

Figure 4.1. Summary of WRRP surface water total mercury data, 2011–2023.

Figure 4.1 summarizes total mercury data, with average, standard deviation of the average, and maximum concentrations and the 51-ng/L surface water AWQC as a reference. As shown in this dataset, the measured maximum concentration has exceeded AWQC only at BCK 12.34 at the Bear Creek headwater, at NT-3, and in spring SS-4, all three of which are in the upper half of Zone 3. The maximum detected mercury result at NT-3 (August 2020) contained 17.1 mg/L of suspended solids, which raises the possibility that

particle-associated mercury is a possible mode of mercury transport in NT-3. Figure 4.2 summarizes methylmercury results. Methylmercury non-detected results are included at the reported value.





BCK = Bear Creek kilometer J = estimated value Max = maximum MeHg = methylmercury NT = North Tributary SS = surface spring StDev = standard deviation UJ = not detected at estimated value

Figure 4.2. Summary of WRRP surface water methylmercury data, 2011–2023.

Figures 4.3 through 4.5 show the flow rate versus total mercury and methylmercury concentrations for BCK 12.34, NT-3 (BC-NT3), BCK 11.54A, NT-8 (BC-NT8), SS-5, and BCK 9.2. In most of the graphs, there is little apparent correlation between daily average flow rate and concentration. However, higher flow rates cause increased sediment transport, and with the strong particle retention of mercury, its mass transport can be greatly increased during high flow events.


Mercury and methylmercury flux rates (mg/day) were calculated using the daily average flow rate from the continuous monitoring records and the reported mercury concentration for each site. Figure 4.6 shows the calculated total mercury flux rate for each monitoring location on each sampled date. The daily average flow rate at BCK 9.2 is shown as an indicator of watershed flow conditions. For the BCK 9.2 flux rate data, labels indicate the total mercury concentration and the calculated daily flux rate. Figure 4.6 shows the responsiveness of the flux rate for all locations to the total flow state as indicated at BCK 9.2.





 $BCK = Bear\ Creek\ kilometer \qquad HgT = total\ mercury \qquad J = estimated\ value \qquad MeHgT = total\ methylmercury \\ JN = matrix\ spike\ outside\ limits;\ estimated\ value \qquad N = matrix\ spike\ outside\ limits \qquad NT = North\ Tributary$ 

Figure 4.3. Flow rate versus total mercury and methylmercury concentrations at BCK 12.34 and Bear Creek NT-3.






Figure 4.4. Flow rate versus total mercury and methylmercury concentrations at BCK 11.54A and Bear Creek NT-8.

JN = matrix spike outside limits; estimated value NU = matrix spike outside limits; not detected at reported quantitation limit NT = North Tributary




Figure 4.5. Flow rate versus total mercury and methylmercury concentrations at spring SS-5 and BCK 9.2.




Figure 4.6. Daily total mercury flux rate for surface water flow measurement stations.

The proportions of calculated daily mercury flux are shown in Figure 4.7. Monitoring location contributions are stacked, with the most upstream location (BCK 12.34) at the bottom and subsequent downstream monitoring locations stacked upward in sequence. Although the NT-3 stream was a historic major source of mercury to Bear Creek prior to BYBY remediation, the post-remediation mercury flux contribution to Bear Creek is minimal. Also confirmed is the negligible contribution from NT-8. The mercury flux measured at BCK 12.34 originates from contaminant discharges associated with the S-3 Ponds groundwater plume.

The flux increase observed between the small NT-3 contribution and the BCK 11.54A flume location is attributed largely to the general increase in flow, because the average and maximum mercury concentration decreases at BCK 11.54A as compared to the upstream locations (Figure 4.1). Spring SS-5 contributes a significant flux of mercury from the Maynardville Limestone karst flow system that is a subsurface channel for contaminant migration originating from multiple upstream sources. At BCK 9.2, the Zone 3 BCV Phase I ROD surface water integration point, the increased flow and decreased average mercury concentration relative to upstream locations accounts for approximately 50% of the Zone 3 surface water total mercury flux.

Figure 4.8 shows the estimated daily flux rate of total methylmercury at the monitored locations. Similar to the total mercury flux rate, the methylmercury flux rate is dominated by water flow variability. Figure 4.9 shows the stacked flux estimates for total methylmercury. Laboratory data qualifiers are shown on Figure 4.9 as a note of caution in interpreting the figure, because many of the laboratory results are estimated or non-detected results.

The WRRP snapshot total mercury data for Bear Creek and tributary water sources indicate concentrations are generally less than the 51-ng/L AWQC level, with occasional higher concentrations in the headwater reach. Sampling conducted during the BCV mercury sources RSE demonstrated that, at baseflow conditions, an average of 27% (+/- 5%) of mercury in surface water was dissolved, with the remaining 73% being associated with filterable solids. Surface water concentrations higher than 51 ng/L are associated with presence of elevated solids in samples.

The distribution of total mercury flux in Zone 3 Bear Creek monitoring data does not point to a specific strong source of mercury entering Bear Creek. The increasing flux in the downstream flow direction is dominated by the increasing flow volume as concentration data tend to gradually decrease. This characteristic suggests mercury entering Bear Creek downstream of the known source associated with the S-3 Ponds area is derived from dispersed mercury that may be associated with secondary contamination of creek bank and floodplain soil.

#### 4.2 BIOLOGICAL DATA

Aquatic biological monitoring of stream sites in the BCV Watershed is used to evaluate stream ecological conditions over time, providing an important measure of the effectiveness of both past and potential future remedial and abatement actions in the watershed. Long-term trends of monitored sunfish show that, despite significant fluctuation, mean mercury concentrations in fish (Figure 4.10) have been generally declining in the last 10 years and are at, below, or just above the fish tissue criterion (0.3 µg/g). Virtually all (approximately 100%) of the mercury in fish fillets in Bear Creek is methylmercury.

Mercury concentrations in rock bass (*Ambloplites rupestris*) collected in Bear Creek remained low in 2023, with concentrations at BCK 3.3 and BCK 9.9 approaching and dropping below the EPA-recommended fish-based AWQC of 0.3 μg/g in October 2023 (Figure 4.10). The fish-based AWQC of 0.3 μg/g is derived for human health risks and is not a BCV Phase I ROD-specified goal; it is, thus, used for comparison purposes only. However, concentrations remain slightly elevated with respect to fish collected from the reference site (HCK 20.6; Figure 4.10). Decreases in fish tissue mercury concentrations have coincided with decreases in aqueous methylmercury concentrations in Bear Creek (ORNL/SPR-2021/2162). The decrease in aqueous methylmercury concentrations and availability of larger fish could be attributed to the significant changes in habitat due to fluctuations in beaver activity over the past few years.





Figure 4.7. Bear Creek Zone 3 surface water total mercury flux proportions measured at various monitoring stations.

BCK 9.2 Daily MeHg Flux (mg/d)






Figure 4.9. Bear Creek Zone 3 estimated total methylmercury flux proportions measured at various monitoring stations.




Figure 4.10. Average concentrations of mercury in Bear Creek fish.

The habitat through the middle-lower stretches (BCK 9.9 to BCK 4.5) of the stream had historically been poor for rock bass such that, in the early 2010s, this species could not be found and redbreast sunfish (*Lepomis auritus*) were collected as a surrogate species. Starting in 2015, as beaver-impounded sections of the middle-lower stretches of the stream created deeper pools, rock bass were present in larger numbers and in larger sizes for bioaccumulation collection. The lack of large beaver dams in FYs 2021–2023 coincided with the smaller numbers and size of rock bass available for collection. In addition, overharvesting is a concern in smaller streams like Bear Creek. Projects that require continual monitoring of larger fish or specific investigations that require increased harvesting can lead to the temporary extirpation of larger size classes of fish from sections of stream. In 2023, the sample sizes and species of fish used for bioaccumulation monitoring were significantly lower than in previous years, as fish were not available for collection despite multiple collection events. Populations of targeted fish species may require additional time to recover. Under equivalent exposure conditions in the stream, larger fish are expected to have higher tissue mercury concentrations. Future monitoring will determine whether fish mercury concentrations remain low as fish populations recover and whether the overall decreasing trend observed throughout the stream continues.

Figures 4.11 and 4.12 compare total mercury and methylmercury concentrations at BCK 9.9 and BCK 3.3, respectively, relative to fish tissue concentrations. Surface water concentrations for mercury and methylmercury are consistently well below the AWQC of 51 ng/L.

Fish tissue concentrations for rock bass at BCK 3.3 have decreased since the maximum in the early 2010s, and redbreast sunfish concentrations at BCK 9.9 have declined over the same period. Rock bass concentrations have fluctuated at BCK 9.9 over the same period but have declined sharply since the maximum in 2017–2018.




Figure 4.11. Mercury and methylmercury concentrations in surface water and total mercury fish tissue concentrations at BCK 9.9.




Figure 4.12. Mercury and methylmercury concentrations in surface water and total mercury fish tissue concentrations at BCK 3.3.

### 5. CONCLUSIONS AND RECOMMENDATIONS

This report presents results of the BCV mercury sources RSE conducted in FY 2024. The objective of the BCV mercury sources RSE is to evaluate potential sources of mercury and methylmercury within the BCV Watershed, as included in the mercury-management approach for Bear Creek in the EMDF ROD, to determine if active remediation is warranted. DOE collected and evaluated the BCV mercury sources RSE data, including channel sediment, creek bank and floodplain soil, and surface water, at transects along Bear Creek. All sampling was conducted as planned between December 2023 and April 2024, per the BCV Mercury Sources RSE SAP, with the exception of a few changes due to field conditions and stream morphology.

#### 5.1 CONCLUSIONS

Transect data showed total mercury in surface water was highest at BCT13 (9.8 ng/L) upstream of NT-3, with decreasing concentrations downstream. Dissolved mercury was highest (1.2 ng/L to 1.9 ng/L) in the stream reach that includes BCT14, BCT13, BCT12A, BCT12B, and BCT11, with decreasing concentrations downstream. Total and dissolved mercury exhibited very little change in concentration in Bear Creek downstream of BCT7 (total mercury 1.5 ng/L to 1.9 ng/L and dissolved mercury 0.42 ng/L to 0.54 ng/L). Dissolved mercury comprised an average of 27% of the total mercury concentration, with the remaining 73% of the mercury being associated with filterable solids. All Bear Creek surface water samples collected in this investigation had mercury concentrations less than the 51-ng/L AWQC level.

Methylmercury in surface water exhibited concentration behavior similar to the pattern for total mercury, with the highest concentration measured at BCT13 (0.27 J ng/L). A significant secondary peak of methylmercury was measured in surface water at sample locations BCT9, BCT8, and BCT7, all of which are associated with beaver ponds. Data collected in June and August 2017 and in August 2018 (ORNL/SPR-2018/902) suggest the beaver dams located along Bear Creek are promoting conditions favorable for methylmercury generation. The ratio of methylmercury to total mercury increased in the downstream portion of Bear Creek. The ratio of dissolved methylmercury to total mercury was higher than for total methylmercury to total mercury. The maximum ratio was measured at BCT2, where the dissolved methylmercury was nearly 18% of the concentration of total mercury.

Floodplain and creek bank soil mercury and methylmercury concentrations were highest downstream from BCK 12.34 and upstream of NT-3 (BCT12B, BCT12A, and BCT13), and decrease farther downstream. Locally, near beaver ponds at the Reeves Road crossing (BCT7, BCT8, and BCT9), there is increased methylmercury in soil (especially in upper bank soils). TOC in floodplain soils is highest at BCT14 (39,200 mg/kg), with decreasing concentrations downstream. TOC concentrations furthest downstream at BCT1, BCT2, and BCT3 (9760 mg/kg to 17,000 mg/kg) were similar to those at the Hinds Creek reference site (17,400 mg/kg). Channel sediment exhibited the highest mercury concentration at BCT12A (530  $\mu$ g/kg) upstream of the NT-3 confluence, with a second peak at BCT9 (440  $\mu$ g/kg) in a beaver pond upstream of the Reeves Road crossing. Channel sediment TOC was highest at location BCT15 (21,200 mg/kg), located on NT-1. Due to stream channel morphology and scour, suitable fine-grained channel sediment was not available to sample at the Hinds Creek reference site and at Bear Creek transects BCT1, BCT2, BCT5, and BCT6.

Results of mercury sequential extraction analyses in Bear Creek floodplain soil samples revealed the majority of mercury was associated with extraction of organic-associated (F3) mercury. Most of the creek bank soil mercury sequential extraction analyses showed the largest fraction of mercury to be extracted with the organic (F3) extraction step; however, the majority of mercury in the creek bank soil at BCT14 was found in the F4 extraction step, thus suggesting mercury sulfide. Mercury sequential extraction analyses of channel sediment similarly showed a strong presence of organic-associated (F3) mercury, with the majority of mercury at BCT14 found in the F4 (presumed sulfide) step, although a significant fraction at BCT14 was extracted in the F5 step that indicates very strongly bound mercury. The F5 fraction was also

significant in channel sediment as far downstream as BCT11, which is downstream of the NT-7 confluence with Bear Creek near BCK 9.9.

The WRRP total mercury and methylmercury flux evaluation at gauged locations in the eastern half of BCV (Zone 3) suggests the dominant effect at play under varying flow conditions is transport of particle-associated mercury that increases with increasing flow rates.

Despite low aqueous mercury concentrations in Bear Creek, the fish fillet concentrations are relatively high, likely due to high methylation efficiency and efficient trophic transfer (ORNL/TM-2024/3735). Fish fillet concentrations were approaching and dropping below the EPA-recommended fish-based AWQC of 0.3 µg/g in fillets in October 2023, but as previously mentioned, mercury concentrations in fish in this creek have fluctuated significantly in recent years with habitat changes. Heavy sampling in this creek has led to a decline in fish numbers and sizes in 2023; the fish collected in 2023 were smaller than average, which could affect mercury concentrations. If populations recover and/or habitat changes in the creek to facilitate mercury methylation, concentrations in fish may increase above AWQC. Future monitoring will determine whether fish mercury concentrations remain low as fish populations recover and whether the overall decreasing trend observed throughout the stream continues.

Overall, mercury contamination of surface water, floodplain and creek bank soil, and channel sediment is highest in the upper reaches of Bear Creek, with decreasing concentrations downstream. The increasing mercury flux in the downstream flow direction is dominated by the increasing flow volume, as concentration data tend to gradually decrease. This characteristic suggests mercury entering Bear Creek downstream of the known source associated with the S-3 Ponds area is derived from dispersed mercury that may be associated with secondary contamination of creek bank and floodplain soil. Although mercury is detected at low levels in sampled media, concentrations are much lower than other ORR mercury-contaminated sites and fish concentrations continue to decrease. This BCV mercury sources RSE did not identify a source of mercury in the media collected in and along Bear Creek that indicates active remediation or an RI of mercury sources is needed. However, the presence of mercury in fish tissue indicates additional studies to understand methylation in Bear Creek are necessary to further refine the CSM.

#### 5.2 **RECOMMENDATIONS**

While the historical water quality and discharge measurement dataset is extensive in Bear Creek, there are far fewer coordinated water-quality-plus-discharge-measurement campaigns with a specific emphasis on total versus dissolved mercury and methylmercury. Although the baseflow analysis of total versus dissolved mercury and methylmercury conducted in this investigation showed approximately 73% of the mercury and 39% of the methylmercury were particle-associated, there are no event-based (i.e., precipitation-driven flows), coordinated, discharge and water quality sampling campaigns. To assess total and dissolved fluxes of mercury and other solutes in addition to those available at established stations in Zone 3, the following additional coordinated flow measurement and sampling locations are needed:

- A BCK 1 monitoring station equipped with a pressure transducer and multiparameter sonde (temperature, pH, specific conductance, dissolved oxygen, and nitrate) was installed in FY 2024. These continuous monitoring instruments may be supplemented with manual or programmed auto-sampling for mercury and methylmercury or other constituents. These measurements can provide flux data at one location.
- Additional gaging stations would be required to fully assess loading at additional locations in Bear Creek. Dedicated, coupled, concentration-discharge measurements under both baseflow conditions and over several flood hydrographs would be ideal to better understand mass loading. Several flow-control

structures at road crossings exist along Bear Creek downstream of existing location BCK 9.2. Evaluating potential instrumentation of selected existing structures may identify opportunities for relatively cost-effective flow measurement and sampling stations for additional flux-measurement monitoring.

In addition to obtaining better discharge measurements in Bear Creek, understanding the biological factors controlling mercury methylation and trophic transfer in this creek is important. In the BCV Watershed, where aqueous mercury concentrations are low but fish tissue concentrations are relatively elevated, the path to achieving and maintaining fish tissue guidelines must include more than a simple assessment of aqueous total mercury concentrations. Mercury, especially in the methylmercury form, biomagnifies through the food web, leading to elevated concentrations in fish. The drivers for mercury methylation in the creek, as well as for mercury bioaccumulation, need to be understood. Because periphyton are likely the substrate that enables mercury methylation and are also the base of the food web, serving to concentrate methylmercury concentrations in the dietary pathways leading to fish, future monitoring should focus on periphyton dynamics.

Although periphyton are understood to play an important role in mercury methylation in stream systems, how the periphyton community changes spatially and temporally in Bear Creek and how these changes relate to methylmercury concentrations in the stream are yet to be understood. Recommendations to evaluate periphyton in Bear Creek in support of the CSM under the mercury remediation technology development program follow:

- Based on guidance provided by Bravo et al. (2017), evaluating organic material characteristics that may affect mercury methylation and biouptake is required to understand the mixes of organic compound origins and molecular weights and to inform attempts to discern potential aquatic habitat characteristics that influence mercury methylation, accumulation in media, and bioaccumulation. This information is critical to any best management watershed-scale practices aimed at reducing methylmercury concentrations in fish.
- To gain a better understanding of mercury methylation for decision-making, data from the aforementioned continuous monitoring instruments should also be supplemented with manual sampling data for measures of periphyton abundance and community structure.
- To understand spatial and temporal periphyton dynamics over time, deploying nutrient (e.g., the nitrate sensor at the BCK 1 monitoring station), light, and turbidity sensors to augment the discharge and water quality measurements collected at additional future monitoring station installations is required. The continuous measurement of periphyton- and mercury-relevant water quality parameters may also be enriched by installing relatively low-cost field cameras capable of estimating aquatic vegetation distribution and abundance in proximity to the monitoring station on a daily basis. Nutrients and light are both important drivers of periphyton distribution and abundance, along with temperature, depth, velocity, and substrate type. Integrated telemetry and web-based data portals to allow these contemporaneous measurements, continuous monitoring stations, field cameras, and stream conditions to be viewed in near real-time would also improve data accessibility and usability and add a new capability to the toolset used to assess the BCV Watershed.

Additional BCV investigations under the mercury remediation technology development program will be documented in a report that is updated annually. The initial document will be issued as a secondary DOE document with an associated FFA Appendix E milestone that presents the results of historical studies, the current CSM for mercury methylation in BCV, and the data gaps that will be addressed in future studies to support further development of the CSM. An annual addendum to that report (also with an FFA Appendix E milestone [i.e., the ORNL Technical Memorandum]) will be issued and will summarize completed activities

and additional investigation recommendations based on the results of those activities. The report will be appended to the annual EMDF Phased Construction Completion Report. An annual roundtable meeting will be held with the FFA parties (DOE, EPA, and TDEC) to provide the status of the investigations and to discuss recommendations for additional studies. While no remedial action is required at this time, the scientific investigations proposed may indicate an opportunity to reduce mercury in Bear Creek in the future. DOE will determine if Bear Creek is in compliance with applicable water quality standards prior to EMDF operations. If Bear Creek is determined to not be in compliance, then DOE commits to a schedule of actions as approved by the FFA parties (DOE, EPA, and TDEC).

### 6. REFERENCES

- Bravo et al. 2017. "Molecular composition of organic matter controls methylmercury formation in boreal lakes," *Nature Communications*, DOI: 10.1038/ncomms14255.
- Crowther et al. 2021. "Use of sequential extraction and mercury stable isotope analysis to assess remobilization of sediment-bound legacy mercury," *Environ. Sci.: Processes Impacts*, 23:756–775.
- DOE/OR/01-1273/D2. Remedial Investigation Report on Bear Creek Valley Operable Unit 2 (Rust Spoil Area, Spoil Area 1, and SY-200 Yard) at the Oak Ridge Y-12 Plant, Oak Ridge, Tennessee, 1995, U.S. Department of Energy, Office of Environmental Restoration and Waste Management, Oak Ridge, TN.
- DOE/OR/01-1455&D2. Report on the Remedial Investigation for Bear Creek Valley at the Oak Ridge Y-12 Plant, Oak Ridge, Tennessee, 1997, U.S. Department of Energy, Office of Environmental Management, Oak Ridge, TN.
- DOE/OR/01-1750&D4. Record of Decision for the Phase I Activities in Bear Creek Valley at the Oak Ridge Y-12 Plant, Oak Ridge, Tennessee, 2000, U.S. Department of Energy, Office of Environmental Management, Oak Ridge, TN.
- DOE/OR/01-2077&D2. Phased Construction Completion Report for the Bear Creek Valley Boneyard/Burnyard Remediation Project at the Y-12 National Security Complex, Oak Ridge, Tennessee, 2004, U.S. Department of Energy, Office of Environmental Management, Oak Ridge, TN.
- DOE/OR/01-2457&D4. Bear Creek Valley Watershed Remedial Action Report Comprehensive Monitoring Plan, Oak Ridge, Tennessee, 2019, U.S. Department of Energy, Office of Environmental Management, Oak Ridge, TN.
- DOE/OR/01-2794&D2/R2. Record of Decision for Comprehensive Environmental Response, Compensation, and Liability Act Oak Ridge Reservation Waste Disposal at the Environmental Management Disposal Facility, Oak Ridge, Tennessee, 2022, U.S. Department of Energy, Office of Environmental Management, Oak Ridge, TN.
- DOE/OR/01-2958&D2. Bear Creek Valley Mercury Sources Remedial Site Evaluation Sampling and Analysis Plan, Oak Ridge, Tennessee, 2024, U.S. Department of Energy, Office of Environmental Management, Oak Ridge, TN.
- DOE/OR/02-1435&D2. Record of Decision for Bear Creek Operable Unit 2 (Spoil Area 1 and SY-200 Yard) at the Oak Ridge Y-12 Plant, Oak Ridge, Tennessee, 1997, U.S. Department of Energy, Environmental Restoration and Waste Management Division, Oak Ridge, TN.
- MacDonald et al. 2000. "Development and Evaluation of Consensus-Based Sediment Quality Guidelines for Freshwater Ecosystems," *Archives of Environmental Contamination and Toxicology*, 39:20–31.
- ORNL/SPR-2018/902. *Bear Creek Special Studies Report 2018*, 2018, Oak Ridge National Laboratory, Oak Ridge, TN.

- ORNL/SPR-2021/2162. Bear Creek Special Studies Report 2021, 2021, Oak Ridge National Laboratory, Oak Ridge, TN.
- ORNL/SPR-2023/3178. Mercury Remediation Technology Development for Lower East Fork Poplar Creek-FY 2023 Update, 2023, Oak Ridge National Laboratory, Oak Ridge, TN.
- ORNL/TM-2016/578. Mercury Content of Sediments in East Fork Poplar Creek: Current Assessment and Past Trends, 2017, Oak Ridge National Laboratory, Oak Ridge, TN.
- ORNL/TM-2023/3069. *Knowledge Summary and Research Needs for Mercury Dynamics in Bear Creek, Oak Ridge, Tennessee*, 2022, Oak Ridge National Laboratory, Oak Ridge, TN.
- ORNL/TM-2024/3735. *Mercury trophic transfer in Bear Creek*, 2024, Oak Ridge National Laboratory, Oak Ridge, TN.
- PROC-ES-2203. Surface Water Sampling Manual and Automated, latest revision, United Cleanup Oak Ridge LLC, Oak Ridge, TN.
- PROC-ES-2300. Soil Sampling, latest revision, United Cleanup Oak Ridge LLC, Oak Ridge, TN.
- PROC-ES-2302. Sediment Sampling, latest revision, United Cleanup Oak Ridge LLC, Oak Ridge, TN.
- TDEC 2019. State of Tennessee Water Quality Standards, Chapter 0400-40-03, "General Water Quality Criteria," Tennessee Department of Environment and Conservation, Division of Water Pollution Control.
- UCOR-4049. Quality Assurance Project Plan for the Water Resources Restoration Program, U.S. Department of Energy, Oak Ridge Reservation, Oak Ridge, Tennessee, latest revision, United Cleanup Oak Ridge LLC, Oak Ridge, TN.
- UCOR-4160. Data Management Implementation Plan for the Water Resources Restoration Program, Oak Ridge, Tennessee, latest revision, United Cleanup Oak Ridge LLC, Oak Ridge, TN.
- UCOR-4189. Quality Assurance Plan for Environmental Characterization and Monitoring, Oak Ridge, Tennessee, latest revision, United Cleanup Oak Ridge LLC, Oak Ridge, TN.
- Xiaotian et al. 2002. "Modeling the high-mercury wet deposition in the southeastern US with WRF-GC-Hg v1.0," *Geosci. Model Dev.*, 15, 3845–3859.

## APPENDIX A.

D2 VERSION OF THE BEAR CREEK VALLEY MERCURY SOURCES REMEDIAL SITE EVALUATION SAMPLING AND ANALYSIS PLAN, OAK RIDGE, TENNESSEE (DOE/OR/01-2958&D2) THAT INCLUDES TENNESSEE DEPARTMENT OF ENVIRONMENT AND CONSERVATION AND U.S. ENVIRONMENTAL PROTECTION AGENCY COMMENTS ON THE D1 VERSION OF THE DOCUMENT

This page intentionally left blank.


# Bear Creek Valley Mercury Sources Remedial Site Evaluation Sampling and Analysis Plan Oak Ridge, Tennessee



DOCUMENT IS UNCLASSIFIED, NON-SENSITIVE APPROVED FOR PUBLIC RELEASE

Name/Organization: Dave Lannom

Date: 07/29/2024 eIRO#: 6713




# Bear Creek Valley Mercury Sources Remedial Site Evaluation Sampling and Analysis Plan Oak Ridge, Tennessee

Date Issued—August 2024

Prepared by the Water Resources Restoration Program United Cleanup Oak Ridge LLC

Prepared for the U.S. Department of Energy Office of Environmental Management

United Cleanup Oak Ridge LLC under contract 89303322DEM000067



# **CONTENTS**

TA	BLES	v			
AC	RONYMS	Vii			
1.	SAMPLING OBJECTIVES				
	1.1 OBJECTIVE	1			
2.	PROJECT INFORMATION				
	2.1 INTRODUCTION	3			
	2.2 BACKGROUND				
	2.2.1 Site Description				
	2.2.2 Summary of Potential Mercury Source Areas				
	2.3 SUMMARY OF EXISTING BEAR CREEK VALLEY MERCURY DATA				
3.	DATA QUALITY OBJECTIVES SUMMARY	11			
	3.1 DATA QUALITY OBJECTIVE STEP 1: STATE THE PROBLEM	11			
	3.2 DATA QUALITY OBJECTIVE STEP 2: IDENTIFY GOALS OF THE STUDY				
	3.3 DATA QUALITY OBJECTIVE STEP 3: IDENTIFY INFORMATION INPUTS	11			
	3.4 DATA QUALITY OBJECTIVE STEP 4: DEFINE THE STUDY AREA	10			
	BOUNDARY	12			
	3.5 DATA QUALITY OBJECTIVE STEP 5: DEVELOP THE ANALYTICAL APPROACH	12			
	3.6 DATA QUALITY OBJECTIVE STEP 6: SPECIFY THE PERFORMANCE OR	12			
	ACCEPTANCE CRITERIA	12			
	3.7 DATA QUALITY OBJECTIVE STEP 7: DEVELOP THE PLAN FOR OBTAINING				
	DATA	13			
4.	SAMPLING AND ANALYSIS PLAN – SAMPLE LOCATION SELECTION	15			
	4.1 SEDIMENT				
	4.2 SOIL				
	4.3 SURFACE WATER				
	4.4 SEQUENTIAL EXTRACTION OF MERCURY COMPOUNDS				
	4.5 FINAL SELECTION OF SAMPLING LOCATIONS	20			
5.	SAMPLING AND ANALYSIS PLAN – ANALYTICAL PROGRAM	21			
	5.1 SUITE OF ANALYTES AND METHODS				
	5.2 LABORATORY-DEFINED VALUES AND REQUESTED REPORTING LIMITS	21			
	5.2.1 Laboratory-Defined Values	21			
	5.2.2 Requested Reporting Limits				
	5.3 ANALYTICAL METHODS	22			
	5.4 FIELD ANALYTICAL SAMPLING AND LABORATORY QUALITY	22			
	ASSURANCE/QUALITY CONTROL	23			
6.	SAMPLING AND ANALYSIS PLAN – DATA MANAGEMENT				
	6.1 DATA VERIFICATION AND VALIDATION				
	6.2 DATA STORAGE	25			
7.	PROJECT ORGANIZATION, SCHEDULE, AND REPORTING	2.7			

	7.1	ORGANIZATION	27
	7.2	SCHEDULE	27
	7.3	REPORTING	27
8.	REF	ERENCES	29
APF	END	IX A. DATA QUALITY OBJECTIVE PRESENTATION AND MEETING MINUTES FO	R
		THE BEAR CREEK VALLEY MERCURY SOURCES REMEDIAL SITE	
		EVALUATION	A-1
APF	END	IX B. BEAR CREEK VALLEY MERCURY SOURCES REMEDIAL SITE EVALUATIO	N
		SAMPLING AND ANALYSIS PLAN TABLES	B-1

# **FIGURES**

Figure 2.1. BCV Watershed zones and potential source areas.		
Figure 2.2. Surface water data for mercury and methylmercury in Bear Creek, NT-3, and Hinds		
Creek, 2011–2022		
Figure 2.3. Average concentrations of mercury in Bear Creek fish	9	
Figure 4.1. BCV transect sampling locations.	17	
Figure 4.2. BCV RSE transect diagram.	19	
TABLES		
Table 2.1. Current surface water and biota sampling in Bear Creek	7	
Table 4.1. List of transect locations in BCV		
Table 5.1. Summary of field and analytical parameters	21	

This page intentionally left blank.

### **ACRONYMS**

AWQC ambient water quality criteria

BCK Bear Creek kilometer
BCT Bear Creek transect
BCV Bear Creek Valley
BRA baseline risk assessment
BYBY Boneyard/Burnyard

CERCLA Comprehensive Environmental Response, Compensation, and Liability Act of 1980

COC contaminant of concern CSM conceptual site model

DMIP Data Management Implementation Plan

DOE U.S. Department of Energy DQO data quality objective

EMDF Environmental Management Disposal Facility
EPA U. S. Environmental Protection Agency

FFA Federal Facility Agreement
HCK Hinds Creek kilometer
MDL method detection limit
NT North Tributary

OREIS Oak Ridge Environmental Information System

ORR Oak Ridge Reservation

OU operable unit

PQL practical quantitation limit

QA quality assurance

QAPP Quality Assurance Project Plan

RI remedial investigation
ROD Record of Decision
RRL requested reporting limit
RSE remedial site evaluation
SAP Sampling and Analysis Plan
SMO Sample Management Office

TDEC Tennessee Department of Environment and Conservation

UEFPC Upper East Fork Poplar Creek

WRRP Water Resources Restoration Program Y-12 Y-12 National Security Complex

This page intentionally left blank.

### 1. SAMPLING OBJECTIVES

#### 1.1 OBJECTIVE

The objective of the Bear Creek Valley (BCV) mercury sources remedial site evaluation (RSE) is to evaluate potential sources of mercury and methylmercury within the BCV Watershed, as outlined in the mercury-management approach for Bear Creek in the *Record of Decision for Comprehensive Environmental Response, Compensation, and Liability Act Oak Ridge Reservation Waste Disposal at the Environmental Management Disposal Facility, Oak Ridge, Tennessee* (DOE/OR/01-2794&D2/R2; Environmental Management Disposal Facility [EMDF] Record of Decision [ROD]). The BCV Watershed is located in the north-central portion of the Oak Ridge Reservation (ORR) west of the Y-12 National Security Complex (Y-12). Y-12 began operations in the 1940s as part of the Manhattan Project for the purpose of enriching uranium for the first atomic bombs. Since that time, the Y-12 missions have changed, and in the 1950s, new processes for separating lithium used large amounts of mercury. Although process functions were performed adjacent to BCV in the Y-12 Main Plant Area, waste from operations at Y-12 were disposed in pits, trenches, and burial grounds in the 2800-acre BCV Watershed.

Prior remedial investigations (RIs) (e.g., Report on the Remedial Investigation of Bear Creek Valley at the Oak Ridge Y-12 Plant, Oak Ridge, Tennessee [DOE/OR/01-1455&D2; BCV RI]) and decision documents in BCV cite mercury as a potential contaminant of concern (COC). A source control action performed under the Record of Decision for the Phase I Activities in Bear Creek Valley at the Oak Ridge Y-12 Plant, Oak Ridge, Tennessee (DOE/OR/01-1750&D4; BCV Phase I ROD) at the remediated Boneyard/Burnyard (BYBY) (Phased Construction Completion Report for the Bear Creek Valley Boneyard/Burnyard Remediation Project at the Y 12 National Security Complex, Oak Ridge, Tennessee [DOE/OR/01-2077&D2]) focused on excavating mercury-contaminated soil along North Tributary (NT)-3. Mercury surface water results in BCV are consistently below Tennessee general ambient water quality criteria (AWQC) (TDEC 2019); however, fish tissue concentrations remain above or near the U.S. Environmental Protection Agency (EPA)-recommended AWQC for mercury (0.3 μg/g in fish). Mercury and methylmercury data for sediment and soil that may contribute to concentrations in fish are limited.

This BCV Mercury Sources RSE Sampling and Analysis Plan (SAP) identifies the locations, media, and sampling methodology that will support the RSE objectives. Impacts of source areas and hydrology on mercury concentrations will be assessed in associated channel sediment, creek bank and floodplain soils, and surface water at multiple sampling transects throughout the length of the stream. Results of this evaluation will be combined with biota data to evaluate potential mercury source areas in BCV to support the RSE and any recommendations made therein.

This page intentionally left blank.

#### 2. PROJECT INFORMATION

#### 2.1 INTRODUCTION

This section contains information about BCV and potential sources of mercury in BCV and summarizes existing mercury data in BCV. Information in this section serves to provide a context for the sampling discussed in later sections of this BCV Mercury Sources RSE SAP.

#### 2.2 BACKGROUND

#### 2.2.1 Site Description

The BCV Watershed is located at the western end of Y-12 in the north-central portion of the ORR west of the Upper East Fork Poplar Creek (UEFPC) Watershed (Figure 2.1). BCV contains closed and active waste disposal facilities. The boundary between the BCV and UEFPC Watersheds is defined by a surface water divide between eastward-flowing East Fork Poplar Creek and westward-flowing Bear Creek. The integration point for Bear Creek is at Bear Creek kilometer (BCK) 9.2 where more than 99% of the available water from the eastern portion of BCV passes through this location either as surface water or groundwater. As illustrated in Figure 2.1, the BCV Watershed is subdivided into three zones based on end use. The subareas of BCV to be investigated under this BCV Mercury Sources RSE SAP represent geographic areas located at or downstream from potential U.S. Department of Energy (DOE) on-site source areas. Based on the EMDF ROD, the end use for Zone 1 and Zone 2 will be revised to restricted recreational and controlled industrial, respectively, which will be codified in an upcoming modification to the BCV Phase I ROD.

#### 2.2.2 Summary of Potential Mercury Source Areas

BCV contains multiple historical waste management and disposal areas that received mercury-contaminated waste streams from Y-12 operations from 1943–1993, in addition to having materials storage areas and construction storage areas (Figure 2.1). There are two RODs for BCV that identify mercury as a COC—the BCV Phase I ROD; and the *Record of Decision for Bear Creek Operable Unit 2 (Spoil Area 1 and SY-200 Yard) at the Oak Ridge Y-12 Plant, Oak Ridge, Tennessee* (DOE/OR/02-1435&D2; BCV Operable Unit [OU]2 ROD). The BCV Phase I ROD cited mercury as a COC posing environmental hazards due to migration from BYBY. BYBY is a former mercury source area that was remediated using hydraulic controls and excavation of visible waste material; Bear Creek tributary NT-3 runs through BYBY. The BCV RI identified mercury as a COC for human health for the following: BYBY, Oil Landfarm, Hazardous Chemical Disposal Area, S-3 Ponds Site, Sanitary Landfill 1, Bear Creek Road Debris Burial, and Creekside Debris Burial. The BCV RI indicated some elevated soil mercury concentrations exist, generally within an order of magnitude of the background criterion (0.34 mg/kg). The baseline risk assessment (BRA) in the BCV RI stated "the sources of mercury and PCBs to the BCV fish are currently unknown."

The Remedial Investigation Report on Bear Creek Valley Operable Unit 2 (Rust Spoil Area, Spoil Area 1, and SY-200 Yard) at the Oak Ridge Y-12 Plant, Oak Ridge, Tennessee (DOE/OR/01-1273&D2; BCV OU2 RI) identified mercury as a COC for human health for the SY-200 Yard, which was a former equipment storage yard used to store nonradioactive contaminated equipment from the 1950s to 1986. Mercury contamination was discovered during construction in 1990, and a soil cover was placed over the site. While other areas (Spoil Area 1 and the Rust Spoil Area) had mercury as a contaminant of potential concern in the BCV OU2 RI, the BRA did not identify mercury as a COC for these areas. The BCV OU2 RI indicated mercury concentrations were elevated at the SY-200 Yard but were generally within an order of magnitude of background; however, free mercury was seen in some of the borings at the SY-200 Yard during the BCV OU2 RI. The BCV OU2 ROD identified the SY-200 Yard as the area with mercury.




Figure 2.1. BCV Watershed zones and potential source areas.

#### 2.3 SUMMARY OF EXISTING BEAR CREEK VALLEY MERCURY DATA

Mercury data for sediment, surface water, and biota in BCV are available in the Oak Ridge Environmental Information System (OREIS). However, sediment data are limited, and no methylmercury data are available for BCV sediment in OREIS. Twenty-nine data points for total mercury in BCV sediment are available ranging from 1993–2011: 7 locations in Zone 1, 2 locations in Zone 2, and 20 locations in Zone 3. Concentrations range from non-detect to 6.9 mg/kg total mercury.

As shown in Table 2.1, under the *Bear Creek Valley Watershed Remedial Action Report Comprehensive Monitoring Plan, Oak Ridge, Tennessee* (DOE/OR/01-2457&D4), surface water and biota sampling have been performed. Results are reported annually in a Remediation Effectiveness Report or every 5 years in a Five-Year Review. Surface water data since 2011 (Figure 2.2) show a steady or declining trend, with mercury below AWQC, with very few exceptions. Mercury concentrations are generally higher upstream and decrease downstream; methylmercury concentrations are more variable upstream to downstream. Long-term trends of monitored sunfish show that, despite significant fluctuation, mean mercury concentrations in fish (Figure 2.3) have been generally declining in the last 10 years and are below or just above the fish tissue criterion (0.3 µg/g) as of 2022.

Table 2.1. Current surface water and biota sampling in Bear Creek

Medium	Performance standard	Sampling frequency	Parameter	Monitoring location
Surface water	AWQC	Semiannual (Q1 and Q3)	Total mercury and methylmercury	BCK 3.3, BCK 4.55, BCK 9.2, BCK-11.54A, BCK 12.34, NT-3, SS-4, and SS-5
		Semiannual (Q2 and Q4) in year before FYR	Total mercury	BCK 4.55, BCK 9.2, BCK 12.34, NT-3, and NT-8
		Annual in year before FYR	Total mercury	BCK-7.87 and NT-1
		Annual in year before FYR	Methylmercury	NT-5
	Trend monitoring	Quarterly Semiannual Quarterly in year before FYR	Bicarbonate, carbonate, chloride, fluoride, and sulfate	NT-1, NT-2, NT-3, SS-4, and SS-5 NT-7 and NT-8 NT-5
	Water quality	Semiannual Quarterly in year before FYR	Total suspended solids and total dissolved solids	NT-7 and NT-8 BCK 4.55, BCK-7.87, BCK 9.2, BCK 12.34, NT-1, NT-3, NT-5, and NT-8
Biota	Baseline sampling	Semiannual	Mercury and methylmercury	BCK 3.3, BCK 9.9, and HCK 20.6 (whole-body stoneroller minnows and rock bass fillets); BCK 12.4 (whole-body stoneroller minnows)
		Annual in year before FYR	Mercury and methylmercury	BCK 9.9 (whole-body caddisflies)

AWQC = ambient water quality criteria BCK = Bear Creek kilometer

FYR = Five-Year Review

HCK = Hinds Creek kilometer

NT = North Tributary Q = quarter

SS = surface spring




Figure 2.2. Surface water data for mercury and methylmercury in Bear Creek, NT-3, and Hinds Creek, 2011–2022.




Figure 2.3. Average concentrations of mercury in Bear Creek fish.

Recently, the Oak Ridge National Laboratory conducted special studies of Bear Creek to better understand the biotic and abiotic factors contributing to mercury concentrations in fish in Bear Creek. These field studies focused on gaining an understanding of the processes controlling mercury methylation and bioaccumulation, with beaver dams and periphyton being key areas of interest. Studies included understanding the role of beaver dams in contributing to mercury dynamics in Bear Creek (2017–2018), evaluating the effects of fine-grained sediment deposition (2019), investigating the potential role tributaries to Bear Creek may have on mercury and methylmercury in the main channel (2020–2021), and evaluating periphyton relationships (2021). These special studies were documented in the *Bear Creek Special Studies Report 2021* (ORNL/SPR-2021/2162).

#### 3. DATA QUALITY OBJECTIVES SUMMARY

The Data Quality Objective (DQO) Process provides systematic planning for decision-making and is an important tool for defining the type, quality, and quantity of data needed to make defensible decisions. EPA developed the *Guidance on Systematic Planning Using the Data Quality Objectives Process* (EPA/240/B-06/001) for implementing the DQO Process as part of its Quality System, an Agency-wide program for environmental data. The DQO Process is a seven-step iterative planning approach used to prepare plans for environmental data-collection activities. It provides a systematic approach for defining criteria that a data-collection design should satisfy, including identifying when, where, and how to collect samples or measurements; determining tolerable decision error rates; and specifying the number of samples or measurements that should be collected. DQOs define the purpose of the data-collection effort, clarify what the data should represent to satisfy this purpose, and specify the performance requirements for the quality of information to be obtained from the data. These outputs, which are developed in the first six steps, are then used in the seventh and final step of the DQO Process to develop a data-collection design that meets all performance criteria and other design requirements and constraints.

DOE, EPA, and Tennessee Department of Environment and Conservation (TDEC) representatives attended a DQO meeting held on June 29, 2023. Appendix A provides the meeting minutes and a copy of the presentation. The BCV Mercury Sources RSE project DQOs are summarized below.

#### 3.1 DATA QUALITY OBJECTIVE STEP 1: STATE THE PROBLEM

The first step in the DQO Process is to concisely describe the problem to be studied. Review of prior studies and existing information is necessary to gain a sufficient understanding to define the problem (i.e., conceptual site model [CSM]). The problem statement identified during the DQO meeting is:

There are insufficient data along Bear Creek and its tributaries to determine if there are potential sources
of mercury and methylmercury in channel sediment and creek bank and floodplain soils that may be
contributing to exceedances of fish tissue criterion in prior years.

#### 3.2 DATA QUALITY OBJECTIVE STEP 2: IDENTIFY GOALS OF THE STUDY

Step 2 of the DQO Process is to identify how data will be used to meet the objectives and what questions the study will attempt to resolve. The goals of the BCV Mercury Sources RSE project are to:

- Determine if there are areas (channel sediment and creek bank and floodplain soils) along Bear Creek and its tributaries that are potential sources of mercury and methylmercury that may affect fish.
- Obtain data from various hydrologic settings (pools, beaver ponds, etc.) that may contribute to mercury
  methylation and its bioaccumulation in the environment of Bear Creek and a reference location (e.g.,
  Hinds Creek).

#### 3.3 DATA QUALITY OBJECTIVE STEP 3: IDENTIFY INFORMATION INPUTS

This step is to identify the information that needs to be obtained and the measurements that need to be taken to achieve the goals of the study. This information is necessary so that proper data may be collected to resolve the problem statement. The information inputs for the BCV Mercury Sources RSE project are to:

- Review potential sources of mercury and methylmercury in Bear Creek and its tributaries.
- Review existing historical biota, surface water, sediment, and special studies data in Bear Creek, its tributaries, and a reference site (Hinds Creek kilometer [HCK] 20.6).

- Collect additional surface water, channel sediment, creek bank soil, and floodplain soil data along Bear Creek and its tributaries from selected transects.
- Collect additional surface water, channel sediment, creek bank soil, and floodplain soil data from the reference site (HCK 20.6).

#### 3.4 DATA QUALITY OBJECTIVE STEP 4: DEFINE THE STUDY AREA BOUNDARY

The purpose of this step is to clarify the site characteristics that the environmental measurements are intended to represent. In this step, time periods and spatial area to which decisions will apply (i.e., determine when and where data will be collected) are specified. Practical constraints that could interfere with sampling also are identified in this step. The BCV Mercury Sources RSE study area boundaries follow:

- Spatial Bear Creek, its tributaries, and a reference location and limited surrounding creek bank soil and floodplain soil.
- Vertical shallow channel sediment, creek bank soil, and floodplain soil (0 to 0.5 ft).
- Temporal samples anticipated to be collected in December 2023 to meet the RSE milestone of September 2024.

#### 3.5 DATA QUALITY OBJECTIVE STEP 5: DEVELOP THE ANALYTICAL APPROACH

This step is to develop an analytic approach that will guide how the study results are analyzed and conclusions are drawn from the data. The key steps for the analytical approach are to:

- Prepare an initial CSM to include all available information on potential mercury sources and historic sediment and surface water monitoring data from Bear Creek.
- Field-locate transects in potential source areas and pool areas (e.g., upstream of beaver dams) in Bear Creek and the mouths of tributaries (e.g., NT-3) based on the reconnaissance survey. Field-locate a reference site.
- Collect surface water, channel sediment, creek bank soil, and floodplain soil samples at transects to determine mercury and methylmercury concentrations along Bear Creek, its tributaries, and the reference site (HCK 20.6).
- Assess and document physical stream conditions (e.g., channel morphology, substrate) at each transect.
- Analyze samples for mercury, methylmercury, nutrients (e.g., sulfate, total phosphorus, nitrate-nitrite as nitrogen, organic carbon), particle size analysis, and mercury speciation/sequential extraction at select locations.
- Screen mercury surface water data against applicable TDEC AWQC.
- Compare concentrations in channel sediment and creek bank and floodplain soil in and around Bear Creek and its tributaries to the reference site (HCK 20.6) concentrations.

### 3.6 DATA QUALITY OBJECTIVE STEP 6: SPECIFY THE PERFORMANCE OR ACCEPTANCE CRITERIA

The purpose of this step is to derive the performance or acceptance criteria that the collected data will need to achieve to minimize the possibility of either making erroneous conclusions or failing to keep uncertainty in estimates to within acceptable levels. Sampling uncertainty and associated decision errors are managed by increasing the number of field samples, which is more effective than controlling measurement uncertainty by repeated laboratory analyses. By designing the data-collection process appropriately, the

level of uncertainty in the data can be controlled to achieve acceptable results. Thus, errors in decisions based on environmental data may be managed effectively by complying with the requirements of the *Quality Assurance Project Plan for the Water Resources Restoration Program*, U.S. Department of Energy, Oak Ridge Reservation, Oak Ridge, Tennessee (UCOR-4049; Water Resources Restoration Program [WRRP] Quality Assurance Project Plan [QAPP]). New data will be obtained under approved WRRP procedures and quality programs and will be archived in OREIS.

#### 3.7 DATA QUALITY OBJECTIVE STEP 7: DEVELOP THE PLAN FOR OBTAINING DATA

The purpose of this step is to identify a field investigation sampling design that meets performance criteria, as specified in the preceding steps of the DQO Process. The output of this step is development of this BCV Mercury Sources RSE SAP. The sampling and analysis approach (Chapters 4 and 5) presents the plan for generating data for the BCV mercury sources RSE that satisfies the DQO and is sufficient to make decisions that achieve RSE requirements.

## 4. SAMPLING AND ANALYSIS PLAN – SAMPLE LOCATION SELECTION

The work contained within this BCV Mercury Sources RSE SAP is consistent with a framework of plans, procedures, and protocols under the WRRP that help ensure all data collected are managed in a manner consistent with Comprehensive Environmental Response, Compensation, and Liability Act of 1980 (CERCLA) requirements. In accordance with this overall objective, the WRRP has developed the WRRP QAPP to identify and implement quality assurance (QA) requirements for use in sample collection, laboratory analysis, and data management of environmental media monitoring activities. The *Data Management Implementation Plan for the Water Resources Restoration Program, Oak Ridge, Tennessee* (UCOR-4160; WRRP Data Management Implementation Plan [DMIP]) serves as the project-level plan for managing all data collected by the WRRP. Together, these plans identify the procedures that are followed in collecting, maintaining custody of, and handling samples, as well as in verifying, validating, and retaining environmental and laboratory data used by the WRRP in preparation of Federal Facility Agreement (FFA) documents.

Sample collection, laboratory analysis, and data management activities performed under this BCV Mercury Sources RSE SAP will follow the requirements of approved, relevant WRRP procedures, as detailed in the WRRP QAPP and WRRP DMIP. Additional requirements governing fieldwork and sample collection, specified in the *Quality Assurance Plan for Environmental Characterization and Monitoring, Oak Ridge, Tennessee* (UCOR-4189), will also apply as appropriate. Per EPA's Uniform Federal Policy, a SAP/QAPP checklist will be submitted under separate cover for EPA approval. The approved checklist will be retained in Appendix F of the WRRP QAPP.

A list of sampling locations along Bear Creek and its tributaries includes transects meeting the requirements of DQO Process Steps 4 and 5 (define the study area boundary and develop the analytical approach) for the collection of channel sediment, creek bank and floodplain soils, and surface water, which was identified in the DQO meeting (Table 4.1 and Figure 4.1). A conceptual diagram of the transect sampling plan is included as Figure 4.2 and is described in Sections 4.1 through 4.4.

#### 4.1 SEDIMENT

Channel sediment samples will be collected at each of the 16 transect monitoring locations as well as at a single reference site (HCK 20.6) following PROC-ES-2302, *Sediment Sampling*. Sediment will be collected to an approximate depth of 0.5 ft and run through a 1-mm sieve until adequate sample volume is achieved.

#### **4.2 SOIL**

Both creek bank and floodplain soils will be collected at each of the 16 transect monitoring locations as well as at a single reference site (HCK 20.6) following PROC-ES-2300, *Soil Sampling*. However, collection of these two soil types will vary as follows:

- Creek bank soils will be divided in half into upper and lower sections based on in-field conditions as follows (Figure 4.2):
  - For the upper section of the creek bank soils, samples will be collected by removing the surface soil on the upper half of each side of the bank. The upper creek bank samples on each side will be composited into a single sample.
  - For the lower section of the creek bank soils, samples will be collected by removing the surface soil on the lower half (above the creek level) of each side of the bank. The lower creek bank samples on each side will be composited into a single sample.
- Floodplain soil will be collected from the upper 0.5 ft on either side of Bear Creek to generate a composite sample representing both sides of the floodplain (Figure 4.2). Loose organic material, such as leaves or brush, will be removed prior to collection.

Table 4.1. List of transect locations in BCV

Sample group	Location		
LOWBCV	BCT1 (upstream of BCK 0.6; downstream of beaver dam)		
	BCT2 (upstream of BCK 0.6; upstream of beaver dam)		
	BCT3 (downstream of BCK 3.3; downstream of beaver dam)		
	BCT4 (downstream of BCK 3.3; upstream of beaver dam)		
BCV ZONE 1	BCT5 (downstream of BCK 4.55; downstream of beaver dam)		
	BCT6 (downstream of BCK 4.55; upstream of beaver dam)		
	BCT7 (downstream of BCK 7.87 at the confluence of NT-13/Bear Creek; downstream of westernmost beaver dam)		
	BCT8 (downstream of BCK 7.87 at the confluence of NT-13/Bear Creek; upstream of westernmost beaver dam)		
	BCT9 (downstream of BCK 7.87; upstream of two beaver dams; southeast of Reeves Road/Haul Road)		
BCV ZONE 2	BCT10 (downstream of surface water integration point BCK 9.2; upstream of EMDF)		
	BCT11 (upstream of NT-8 at BCK 9.9)		
	BCT12A (upstream of NT-3 at the confluence with Bear Creek)		
DOW ZONE 2	BCT12B (downstream of BYBY at the confluence of NT-3/Bear Creek)		
BCV ZONE 3	BCT13 (upstream of BYBY, EMWMF, and NT-3)		
	BCT14 (downstream of SY-200 Yard, Spoil Area 1, and S-3 Ponds Site)		
	BCT15 (downstream of S-3 Ponds Site)		
Hinds Creek	HCTREF (HCK 20.6 reference site)		

BCK = Bear Creek kilometer

BCK = Bear Creek kilometer
BCT = Bear Creek transect
BCV = Bear Creek Valley
BYBY = Boneyard/Burnyard
EMDF = Environmental Management Disposal Facility
EMWMF = Environmental Management Waste Management Facility

HCK = Hinds Creek kilometer

HCTREF = Hinds Creek transect reference site

NT = North Tributary




Figure 4.1. BCV transect sampling locations.




Figure 4.2. BCV RSE transect diagram.

#### 4.3 SURFACE WATER

Surface water samples will be collected at each of the 16 transect monitoring locations as well as at a single reference site (HCK 20.6) following PROC-ES-2203, *Surface Water Sampling – Manual and Automated*.

Because filtered and unfiltered mercury and methylmercury sample volumes are to be collected for analysis, a peristaltic pump will be required for filtration in addition to the grab method. Surface water sampling should be conducted before channel sediment is collected to avoid interference between media.

#### 4.4 SEQUENTIAL EXTRACTION OF MERCURY COMPOUNDS

Eight transect locations, as well as a reference location (HCK 20.6), will be sampled for mercury speciation/sequential extraction analysis. Sufficient mass of solid material from the channel sediment, creek bank soil, and floodplain soil at each selected transect for mercury speciation/sequential extraction will be composited into samples from each of the three representative media types (i.e., three composite samples per transect). Sediment will be collected to an approximate depth of 0.5 ft and passed through a 1-mm sieve until adequate sample volume is achieved. For creek bank soil, one sample will be collected from each side of the bank by removing the upper 0.5 ft of bank soil surface just above the water level and composited. For floodplain soil, samples will be collected from the upper 0.5 ft on each side of the creek to generate a composite sample representing both sides of the floodplain.

Locations for this analysis are shown on Figure 4.1 and include one at the reference site (HCK 20.6); one downstream of the SY-200 Yard, Spoil Area 1, and S-3 Ponds Site (Bear Creek transect [BCT]14); one downstream of BYBY at the confluence of Bear Creek and NT-3 (BCT12B); one upstream of NT-8 at BCK 9.9 (BCT11); three downstream of BCK 7.87 at the confluence of Bear Creek and NT-13 in proximity to two prominent beaver dams (BCT7, BCT8, and BCT9); and two downstream of BCK 4.55 in proximity to a beaver dam (BCT5 and BCT6).

#### 4.5 FINAL SELECTION OF SAMPLING LOCATIONS

Table 4.1 presents the transect sample locations for the BCV mercury sources RSE. These locations along Bear Creek were selected based on their location downgradient of and/or in the vicinity of potential source areas, where biota and surface water sampling have historically occurred, and in the vicinity of ponds. Physical stream conditions (e.g., channel morphology, substrate) at each transect were assessed in the selection process.

Actual field locations may be adjusted based on field conditions and sampling viability. Deviations from this BCV Mercury Sources RSE SAP will be documented in the field logbook and in the BCV mercury sources RSE.

#### 5. SAMPLING AND ANALYSIS PLAN – ANALYTICAL PROGRAM

#### 5.1 SUITE OF ANALYTES AND METHODS

The planned suite of analytes and methods of analysis for all samples to be collected during the BCV mercury sources RSE are listed in Appendix B (Tables B.1 through B.3) and summarized in Table 5.1. The suite is based on discussions and input received during the DQO Process; consideration of primary COCs mercury and methylmercury from potential mercury source areas within the BCV Watershed; and the standard suite of analytes and analytical methods used for sediment, soil, and surface water by the WRRP. As such, results for the analyte suite will be consistent with and comparable to the water quality database for the ORR that is maintained in OREIS. As shown in Appendix B, each of the parameter groups for identified analytes corresponds with a table in the latest version of the WRRP QAPP, which has been revised to add the BCVRSE, S-BCVRSE, and HGSEQ parameter groups for this project.

Table 5.1. Summary of field and analytical parameters

Medium	Field parameter	Analytical parameter
Surface water	Temperature	Dissolved and total mercury
	Dissolved oxygen	Dissolved and total methylmercury
	Turbidity	Metals
	pН	Phosphorous (total)
	Specific conductance (conductivity)	Total organic carbon
	Oxidation-reduction potential	Dissolved organic carbon
		Total dissolved solids
		Total suspended solids
		Anions (chloride, fluoride, nitrate-nitrite as nitrogen, sulfate, and sulfide)
Sediment and soil	None	Total mercury
		Total methylmercury
		Metals
		Total organic carbon
		Particle size analysis
		Anions (chloride, fluoride, nitrate, nitrite, sulfate, and sulfide)
		Sequential extraction of mercury compounds

#### 5.2 LABORATORY-DEFINED VALUES AND REQUESTED REPORTING LIMITS

To develop the analytical program, different values were considered for each analyte.

#### 5.2.1 Laboratory-Defined Values

Laboratory-defined values for the BCV mercury sources RSE analytes and analytical methods are listed in Appendix B (Tables B.2 and B.3) and discussed below.

#### 5.2.1.1 MDLs

Method detection limits (MDLs) apply to non-radionuclide analytes and are defined as the minimum concentration of an analyte that can be measured and reported with a 99% confidence that the analyte is present in the sample with a concentration greater than zero. Analyte concentrations at the MDL have a 50% chance of being reported as a non-detect or a false negative, and analyte concentrations near the MDL cannot be quantified with statistical rigor. Values above the MDL but below the practical quantitation limit (PQL) indicate the analyte is likely present in the sample, although at concentrations below those that can be quantified with statistical significance (DOD/DOE 2013).

#### 5.2.1.2 PQLs

PQLs apply to non-radionuclide analytes and are defined as the lowest concentration of an analyte that produces a quantitative result within specified limits of precision and bias. The PQL is typically greater than the MDL. PQLs are dependent on the acceptance limits for precision and bias selected for the requirements of the program. For many projects, the PQL is required to be at or above the lowest concentration of the laboratory standards used in method calibration for an analyte. Measurements falling between the MDL and PQL assure the presence of an analyte with confidence, but their numeric values are estimates and not quantified numbers (DOD/DOE 2013).

#### **5.2.2** Requested Reporting Limits

Requested reporting limits (RRLs), referred to as reporting limits in the *Consolidated Quality Systems Manual (QSM) for Environmental Laboratories* (DOD/DOE 2013), are concentration levels for specific constituents within a sample that are specified by the project. The RRLs are defined so that obtained sediment, soil, and surface water data meet all project requirements for reporting quantitative data with known precision and bias for a specific analyte in a specific matrix. For the BCVRSE, S-BCVRSE, and HGSEQ parameter groups, the laboratory is being requested to report detections with respect to the MDLs, which are generally lower than the RRLs. The RRLs, if met, ensure project data can be successfully screened against appropriate criteria and standards. For most WRRP projects, laboratories are requested to report detections of chemical analytes with respect to the MDL.

#### 5.3 ANALYTICAL METHODS

Analytical methods for sediment, soil, and surface water analyses are summarized in Appendix B, Table B.2 (water) and Table B.3 (soil and sediment) and correspond to methods listed by parameter group in the latest version of the WRRP QAPP for each analyte. All analytical methods are EPA standard procedures routinely employed by Oak Ridge Sample Management Office (SMO) contract laboratories.

Discussions during the DQO Process resulted in development of the BCVRSE, S-BCVRSE, and HGSEQ parameter groups (WRRP QAPP) which are unique to the BCV mercury sources RSE. These parameter groups were developed to specify analytes (e.g., mercury, methylmercury, nutrients, particle size analysis, and total organic carbon) and methods for the sequential extraction of mercury in sediment and soil for the BCV mercury sources RSE. For surface water samples collected, AWQC may be used for comparison purposes only, but are not a required screening level.

## 5.4 FIELD ANALYTICAL SAMPLING AND LABORATORY QUALITY ASSURANCE/QUALITY CONTROL

All relevant QA and quality control procedures and requirements specified in the WRRP QAPP (field collection) and for the SMO (laboratory analyses) are incorporated by reference for compliance. No changes in WRRP and SMO procedures incorporated under the WRRP QAPP are anticipated for the BCV mercury sources RSE.

Appendix B provides the planning tables that will be used for the BCV mercury sources RSE, including locations, sampling methods, frequencies, analyses, and reporting levels. Final selection of locations will be decided as described in Section 4.5 of this BCV Mercury Sources RSE SAP.

#### 5.5 ANALYTICAL MEDIA CONSIDERATIONS

Surface water (Appendix B, Table B.2) will be analyzed for dissolved and total mercury, dissolved and total methylmercury, total phosphorous, dissolved organic carbon, total dissolved solids, total suspended solids, total organic carbon, anions (e.g., chloride, fluoride, nitrate-nitrite as nitrogen, sulfate, and sulfide), and metals. Both filtered and unfiltered surface water samples will be collected for mercury and methylmercury as part of this suite. Field parameters collected at the time of sampling are temperature, dissolved oxygen, turbidity, pH, specific conductivity, and oxidation-reduction potential.

Sediment and soil (Appendix B, Table B.3) will be analyzed for total mercury and methylmercury, particle size analysis, total organic carbon, anions (e.g., chloride, fluoride, nitrate, nitrite, sulfate, and sulfide), and metals. Sediment and soil will also have additional analysis performed for mercury speciation. This analysis will provide data for volatile elemental mercury, water soluble mercury, pH2 soluble mercury, 1N potassium hydroxide extractable mercury, 12N nitric acid soluble mercury, aqua regia soluble mercury residue, and mineral-bound mercury.

#### 6. SAMPLING AND ANALYSIS PLAN – DATA MANAGEMENT

#### 6.1 DATA VERIFICATION AND VALIDATION

All data will be verified following WRRP QAPP and WRRP DMIP requirements. All mercury and methylmercury data will be validated following the WRRP QAPP and WRRP DMIP. Level 4 data packages will be required for all analyses completed under the BCV Mercury Sources RSE project. Verification and validation will be conducted by United Cleanup Oak Ridge LLC/RSI Entech staff and/or their validation subcontractor.

#### **6.2 DATA STORAGE**

All data will be stored in the Project Environmental Measurements System following required procedures and WRRP QAPP requirements and will be archived in OREIS.

#### 7. PROJECT ORGANIZATION, SCHEDULE, AND REPORTING

#### 7.1 ORGANIZATION

The EMDF ROD outlined the mercury-management approach for Bear Creek that included an RSE. The DOE Environmental Management Program is the responsible organization for implementing the RSE under the CERCLA process, with coordination and approval by EPA and TDEC in accordance with the FFA. The Project Team is comprised of representatives from DOE, EPA, and TDEC. The DOE Environmental Management Program will use the WRRP, a contractor-implemented organization, for support in executing BCV Mercury Sources RSE project monitoring. The WRRP has comprehensive procedures for sampling and provides data for use in making watershed-management decisions related to remedial action effectiveness and contaminant trends on the ORR. WRRP support will include QA, sampling and analysis, and data management resources. Additional details about WRRP organizations, roles, and responsibilities are provided in the WRRP QAPP.

#### 7.2 SCHEDULE

Fieldwork described in this BCV Mercury Sources RSE SAP is anticipated to be conducted in December 2023. Data evaluation and preparation of the BCV Mercury Sources RSE Report will occur between January and September 2024 (FFA Appendix E milestone: September 30, 2024).

#### 7.3 REPORTING

Sampling activities, sampling results, and data evaluation will be summarized in the BCV Mercury Sources RSE Report that has an FFA Appendix E milestone date of September 30, 2024.

#### 8. REFERENCES

- DOD/DOE 2013. Consolidated Quality Systems Manual (QSM) for Environmental Laboratories, version 5.0, U.S. Department of Defense/U.S. Department of Energy, Washington, D.C.
- DOE/OR/01-1273&D2. Remedial Investigation Report on Bear Creek Valley Operable Unit 2 (Rust Spoil Area, Spoil Area I, and SY-200 Yard) at the Oak Ridge Y-12 Plant, Oak Ridge, Tennessee, 1995, U.S. Department of Energy, Office of Environmental Restoration and Waste Management, Oak Ridge, TN.
- DOE/OR/01-1455&D2. Report on the Remedial Investigation of Bear Creek Valley at the Oak Ridge Y-12 Plant, Oak Ridge, Tennessee, 1997, U.S. Department of Energy, Office of Environmental Management, Oak Ridge, TN.
- DOE/OR/01-1750&D4. Record of Decision for the Phase I Activities in Bear Creek Valley at the Oak Ridge Y-12 Plant, Oak Ridge, Tennessee, 2000, U.S. Department of Energy, Office of Environmental Management, Oak Ridge, TN.
- DOE/OR/01-2077&D2. Phased Construction Completion Report for the Bear Creek Valley Boneyard/Burnyard Remediation Project at the Y-12 National Security Complex, Oak Ridge, Tennessee, 2004, U.S. Department of Energy, Office of Environmental Management, Oak Ridge, TN.
- DOE/OR/01-2457&D4. Bear Creek Valley Watershed Remedial Action Report Comprehensive Monitoring Plan, Oak Ridge, Tennessee, 2019, U.S. Department of Energy, Office of Environmental Management, Oak Ridge, TN.
- DOE/OR/01-2794&D2/R2. Record of Decision for Comprehensive Environmental Response, Compensation, and Liability Act Oak Ridge Reservation Waste Disposal at the Environmental Management Disposal Facility, Oak Ridge, Tennessee, 2022, U.S. Department of Energy, Office of Environmental Management, Oak Ridge, TN.
- DOE/OR/02-1435&D2. Record of Decision for Bear Creek Operable Unit 2 (Spoil Area 1 and SY-200 Yard) at the Oak Ridge Y-12 Plant, Oak Ridge, Tennessee, 1997, U.S. Department of Energy, Environmental Restoration and Waste Management Division, Oak Ridge, TN.
- EPA/240/B-06/001. Guidance on Systematic Planning Using the Data Quality Objectives Process, 2006, U.S. Environmental Protection Agency, Washington, D.C.
- ORNL/SPR-2021/2162. Bear Creek Special Studies Report 2021, 2021, Oak Ridge National Laboratory, Oak Ridge, TN.
- PROC-ES-2203. Surface Water Sampling Manual and Automated, latest revision, United Cleanup Oak Ridge LLC, Oak Ridge, TN.
- PROC-ES-2300. Soil Sampling, latest revision, United Cleanup Oak Ridge LLC, Oak Ridge, TN.
- PROC-ES-2302. Sediment Sampling, latest revision, United Cleanup Oak Ridge LLC, Oak Ridge, TN.

- TDEC 2019. State of Tennessee Water Quality Standards, Chapter 0400-40-03, "General Water Quality Criteria," Tennessee Department of Environment and Conservation, Division of Water Pollution Control.
- UCOR-4049. Quality Assurance Project Plan for the Water Resources Restoration Program, U.S. Department of Energy, Oak Ridge Reservation, Oak Ridge, Tennessee, latest revision, United Cleanup Oak Ridge LLC, Oak Ridge, TN.
- UCOR-4160. Data Management Implementation Plan for the Water Resources Restoration Program, Oak Ridge, Tennessee, latest revision, United Cleanup Oak Ridge LLC, Oak Ridge, TN.
- UCOR-4189. Quality Assurance Plan for Environmental Characterization and Monitoring, Oak Ridge, Tennessee, latest revision, United Cleanup Oak Ridge LLC, Oak Ridge, TN.

# APPENDIX A. DATA QUALITY OBJECTIVE PRESENTATION AND MEETING MINUTES FOR THE BEAR CREEK VALLEY MERCURY SOURCES REMEDIAL SITE EVALUATION

## BEAR CREEK VALLEY (BCV) MERCURY SOURCES REMEDIAL SITE EVALUATION (RSE) DATA QUALITY OBJECTIVES (DQO) MEETING MINUTES BCV

**DATE:** June 29, 2023; 2 p.m.

#### **ATTENDEES:**

Sam Scheffler – DOE

Roger Petrie – DOE

Dana Casey – TDEC

Diana Turner – UCOR

Bob Bock – UCOR (ph)

Lynn Sims – UCOR

Lynn Sims – UCOR

Cody Juneau – TDEC

Courtney Thomason – TDEC

Brad Stephenson – TDEC

Randy Young– TDEC

Jana Dawson – EPA (ph)

Eddie Arnold – UCOR, presenter

Annette Primrose – UCOR (ph)

Scott Brooks – ORNL-ESD

Natalie Landry – ORNL-ESD

Chris DeRolph – ORNL-ESD (ph)

Terry Mathews – ORNL-ESD

Sally Absher – Leidos (ph)

Sid Garland – UCOR

**PURPOSE:** The objective of the meeting is to review the history and sources of mercury in Bear Creek and to present DQOs and proposed sampling for the BCV Mercury Sources RSE.

#### **AGENDA (Slide 2):**

- Introduction, Safety Topic
- BCV Mercury Sources RSE Milestone
- Site Background
- Previous Investigations
- DQO steps
- Proposed transects and analytes
- RSE schedule

**INTRODUCTION, SAFETY TOPIC (Slide 3):** Eddie Arnold introduced the participants in the conference room and online and presented a brief safety topic about fireworks in anticipation of the upcoming 4<sup>th</sup> of July holiday.

#### **BCV Mercury Sources RSE Milestone (Slide 4)**

The BCV Mercury Sources RSE Milestone is 9/30/2024 as part of the Federal Facility Agreement (FFA) Appendix E. The RSE is being conducted per an agreement as part of the Environmental Management Disposal Facility (EMDF) Record of Decision (ROD).

**NOTE:** The remainder of these minutes only includes notes for slides in which there was additional discussion. For slides on which there were no additional questions, comments, or discussions, only the slide title is presented. The final DQO Presentation is attached to these minutes.

#### **Site Background - BCV History (Slide 5)**

#### Oak Ridge Reservation [figure] (Slide 6)

#### **BCV Site Location [figure] (Slide 7)**

- Randy Young prompted DOE to explain why the RSE/milestone was in place.
- Roger Petrie explained that Bear Creek is currently listed as impaired and under the antidegradation rule, no new discharges of mercury from EMDF are allowed; this is not possible, so to

construct the EMDF Treatment Facility, DOE agreed to follow a sequence of events in the Comprehensive Environmental Response, Compensation, and Liability Act (CERCLA) to address mercury offsets to get the EMDF ROD signed. The first CERCLA step is the RSE, which DOE has committed to perform. The purpose of the DQO is to present the activities required to complete the RSE.

- Randy Young also clarified for EPA that there was a short discussion while waiting for others to join the call between Randy Young and Roger Petrie about polychlorinated biphenyls (PCBs). DOE stated that PCBs will not be an issue for this facility because DOE expects the PCB inventory to be less than that for EMWMF and for the wastewater treatment to remove all PCBs before discharge to Bear Creek. Therefore, they are not addressed in this approach. Roger Petrie also stated that DOE will develop an offset approach for PCBs if needed. Randy Young asked about potential risks to the cleanup program if landfill operations are impacted by PCB discharge and advised DOE to not take unnecessary risks. Roger Petrie maintained that DOE does not believe it will be an issue and is willing to take the risk in this approach of developing offsets later if necessary. NOTE: During comment resolution on the meeting minutes DOE confirmed that EPA Method 1668 (congener method) will be utilized for comparison of surface water PCB results to the AWQC as applicable.
- Roger Petrie clarified that because the EMDF facility is new, the anti-degradation rule applies, but other existing facilities' discharges are grandfathered in and the anti-degradation rule doesn't apply.

#### BCV Decision Documents – BCV Mercury Sources [BCV Phase 1 RI and ROD] (Slide 8)

- Randy Young had a question about the second bullet on the slide: has the BCV Phase I RI been looked at enough to know how much of a problem mercury is at Sanitary Landfill 1?
- UCOR clarified that mercury was not a contaminant of concern (COC) for the landfill in the BCV Phase I ROD.

#### BCV Decision Documents – BCV Mercury Sources [BCV OU2 RI and ROD] (Slide 9)

Potential BCV Mercury Sources [figure] (Slide 10)

#### **BCV Mercury Sources – SY-200 Yard (Slide 11)**

- Randy Young asked for clarification on the timing/approval of the cover/cap at the SY-200 Yard
- UCOR and DOE responded that it was part of the BCV OU-2 ROD which was pre-FFA

#### **BCV Mercury Sources – Spoil Area 1 (Slide 12)**

**BCV Mercury Sources – NT-3 (Slide 13)** 

#### **BCV Mercury Sources – Others (Slide 14)**

- Dana Casey asked what defines minor level of mercury.
- Eddie Arnold responded that it is likely over background, but that they were generally very minor exceedances
- Brad Stephenson asked for further clarification in the case of 58-83 mg/kg, if that was also considered minor
- Eddie Arnold clarified that at the time it was considered minor, as the wording is from the BCV Phase I RI

#### Previous Investigations – Current Surface Water and Biota Sampling in Bear Creek [fig.] (Slide 15)

Previous Investigations—Current Surface Water and Biota Sampling in Bear Creek [table] (Slide 16)

#### Previous Investigations – Summary of Historical Surface Water Data (Slide 17)

- Jana Dawson asked for clarification about mercury vs methylmercury, generation vs release.
- Eddie Arnold clarified that the evaluation is from the perspective of the effect on biota.

#### **Previous Investigations – Summary of Historical Biota Data (Slide 18)**

- Eddie Arnold invited input from ORNL.
- Terry Mathews mentioned that there had been recent habitat changes that affected mercury methylation but that mercury in fish has been trending downward in the last few years. Additionally, mercury (in fish) at the reference site (where there is no DOE input) has increased; Bear Creek is now around background.
- Randy Young asked if there were any other things that might be addressed regarding habitat in a remedy for mercury or methylmercury.
- Terry Mathews clarified that best management practices and beaver management are being followed beavers are not necessarily bad for habitat but exacerbate mercury methylation, so beaver management is one of the best things to do. Terry added that mercury methylation is discussed later in the presentation and that the focus is on watershed-scale practices.

#### Previous Investigations – Summary of Historical Sediment Data [figure] (Slide 19)

#### Previous Investigations – Summary of Historical Sediment Data (Slide 20)

- Courtney Thomason asked about details regarding historical sediment samples if they were grab samples, what was the depth, etc.
- Eddie Arnold responded that the few samples were mostly surface grab samples (under 6 inches), with a few deeper samples. Concentrations were low and no methylmercury data was collected except during the ORNL special studies.

#### **Previous Investigations – Summary of Special Studies Data (Slide 21)**

#### **Previous Investigations – Summary of Special Studies Data (Slide 22)**

- Scott Brooks presented special studies data (slides 22-26) and oriented viewers to figures.
- Although concentrations are elevated at NT-3 and at the borrow area near Highway 95 due to beaver impoundment, mass loading is thought to be low due to the small amount of discharge.

#### **Previous Investigations – Summary of Special Studies Data (Slide 23)**

#### **Previous Investigations – Summary of Special Studies Data (Slide 24)**

#### **Previous Investigations – Summary of Special Studies Data (Slide 25)**

- There are two properties of methylmercury that are important in this case: 1) that it is bioaccumulative 2) that periphyton is known to be a source of methylmercury generation; both factors lead to high levels of methylmercury in periphyton.
- Cody Juneau asked if methylmercury at NT-3 (tributaries) is much higher.
- Scott Brooks clarified that the tributaries are represented by the orange triangles and confirmed that methylmercury is very high at NT-3 relative to other samples, but also cautioned that it is only a single sample and difficult to separate from pond muck. More data are needed to draw conclusions.

#### **Previous Investigations – Summary of Special Studies Data (Slide 26)**

- Terry Mathews followed up on Randy Young's previous questions about other actions that may be performed watershed-wide. Studies have indicated that periphyton is a potential contributor of methylmercury to the creek. Chris DeRolph has been using drones to look spatially at periphyton communities and habitat throughout the creek.
- Courtney Thomason asked if microbes have been evaluated separately from periphyton.
- Scott Brooks responded that they have in East Fork but not in BCV.
- Brad Stephenson asked how often surface water is sampled.
- Scott Brooks responded that NT-3 was sampled once or twice; Eddie Arnold responded that WRRP does regular quarterly sampling there.

#### **DQO Steps (Slide 27)**

• Eddie Arnold resumed presenting.

#### **DQO Step 1. State the Problem (Slide 28)**

#### DQO Step 2. Identify the Goals of the Study (Slide 29)

- TDEC and Roger Petrie discussed that the goal of the study was to find mercury sources to offset, if possible, not to do an RI, but that data show finding mercury sources to offset may be difficult.
- Discussion continued that mercury concentrations in fish need to be below 0.3 ppt because of the anti-degradation rule. So in addition to the RSE, fish need to be monitored to see if they remain below 0.3 ppt. It's not strictly about mass of mercury produced by EMDF as that will be very small.

#### **DQO Step 3. Identify Information Inputs (Slide 30)**

#### DQO Step 4. Define the Study Area Boundary (Slide 31)

#### **DQO Step 5. Develop the Analytical Approach (Slide 32)**

- Cody Juneau asked that mercury speciation be quickly explained.
- Terry Mathews responded that it is sequential extraction an iterative process with increasingly harsh digestion which results in a percentage of mercury coming off at each step; this determines how tightly bound the mercury is.

#### DQO Step 6. Specify the Performance or Acceptance Criteria (Slide 33)

#### **DQO** Step 7. Develop the Plan for Obtaining Data (Slide 34)

#### **Proposed Transect Locations (Slide 35)**

• Eddie Arnold clarified that there is a general idea of locations but transects will be field-located based on access.

#### **Proposed Transect Locations [figure] (Slide 36)**

#### Beaver Dams near BCK 7.0 [figure] (Slide 37)

- Courtney Thomason asked about sediment deposition not associated with beavers.
- Eddie Arnold responded that overbank depositional areas are limited and the upper portion of Bear Creek is often dry.
- Courtney Thomason asked if there will be any effort to locate any non-beaver depositional areas in the lower portion of Bear Creek.
- Eddie Arnold responded that there will be an effort but added that the substrate doesn't lend itself to fine-grained sediment deposition in non-beaver areas.

#### **Transect Sampling (Slide 38)**

- Dana Casey asked if there is reason to think that there would be mercury deeper than 6 inches that could connect to the surface water.
- Eddie Arnold agreed that it is a possibility but the groundwater data do not indicate that.
- TDEC, UCOR, DOE, and ORNL participated in a discussion of shallow groundwater and concluded it is out of the scope for the RSE but would be an interesting topic for a separate investigation. NOTE: During comment resolution on the meeting minutes TDEC clarified the recommendation that shallow groundwater sampling should be included in the project scope.
- There was discussion to clarify the goal of this RSE and whether that was to find mercury sources, methylmercury sources, or sources of methylation, and DOE reiterated that this is a source investigation for mercury and methlymercury.

#### **Transect Sampling Diagram (Slide 39)**

#### **Analytical Suite (Slide 40)**

#### RSE Schedule (Slide 41)

• Eddie Arnold reviewed the schedule and TDEC asked about what happens afterward/schedule going forward.

- Roger Petrie and TDEC discussed hypothetical future actions, and UCOR mentioned that those discussions are better left until after the results of the RSE.
- Roger Petrie reiterated that this is a very tight schedule but that it can be met. Other valid questions that arose during today's discussion may eventually be addressed but cannot be added to this RSE due to schedule.
- Randy Young agreed but anticipated that TDEC will have several comments. Brad Stephenson also mentioned that TDEC wants EMDF to succeed and this RSE is a big part of getting there and of signing the ROD.

#### Wrap Up

- There were no additional comments or questions on the BCV RSE DQO presentation.
- TDEC said that they had some data they would share.
- Eddie Arnold mentioned that minutes might be delayed due to the upcoming July 4 holiday.

There were no further questions or comments. The meeting was adjourned at 3:33 pm.

Respectfully submitted



Bear Creek Valley (BCV)
Mercury Sources
Remedial Site
Evaluation (RSE) Data
Quality Objectives
(DQOs)







June 29, 2023




### **Agenda**

- 1. Introductions, Safety Topic
- 2. BCV Mercury Sources RSE Milestone
- 3. Site Background
  - BCV History
  - BCV Decision Documents
  - Potential BCV Mercury Sources
- 4. Previous Investigations
  - Current surface water and biota sampling in Bear Creek
  - Summary of historical data
  - Summary of special studies data
- 5. DQO steps
- 6. Proposed transects and analytes
- 7. RSE Schedule



## **Safety Topic – Fireworks!!!**







#### **BCV** Mercury Sources RSE Milestone

## FY 2023 – 2025 Federal Facility Agreement Milestones Appendix E

Project/Subproject	FY 2023 Milestone	FY 2024 Milestone	FY 2025 Milestone
BCV Mercury Sources		RSE 9/30/24	

As part of the Environmental Management Disposal Facility Record of Decision (EMDF ROD; DOE/OR/01-2794&D2/R2) it was agreed to conduct a RSE (40 CFR 300.420)




## **Site Background - BCV History**

- BCV contains multiple historical waste management and disposal areas that received mercury contaminated waste streams from Y-12 operations from 1943 to 1993 in addition to having materials storage areas and construction storage areas.
- East of the BCV Watershed is the Upper East Fork Poplar Creek (UEFPC) Watershed including the operational portion of the Y-12 plant. The boundary between the two watersheds is defined by a surface water divide that is between the eastward-flowing EFPC and westward-flowing Bear Creek.
- The Integration Point (IP) for Bear Creek is at BCK 9.2 where more than 99% of the available water from the eastern portion of BCV passes through this location either as surface water or groundwater.
- BCV has two RODs that identify mercury as a constituent of concern (COC): BCV OU2 ROD and BCV Phase I ROD.




## Oak Ridge Reservation



#### **BCV Site Location**







#### **BCV Decision Documents – BCV Mercury Sources**

BCV Phase I ROD (DOE/OR/01-1750&D4) and BCV Phase I Remedial Investigation (RI) (DOE/OR/01-1455&D2)

- The BCV Phase I ROD cited mercury as a COC posing environmental hazards due to migration from the Boneyard/Burnyard (BYBY). No other mention of mercury in the BCV Phase I ROD.
- The BCV Phase I RI identified mercury as a COC (human health) for BYBY, Oil Landfarm (OLF), Hazardous Chemical Disposal Area (HCDA), S-3 site, Sanitary Landfill 1, Bear Creek Road Debris Burial, and Creekside Debris Burial.
- The BCV Phase I RI indicated there were some elevated soil mercury concentrations, generally within an order of magnitude of background criterion (0.34 mg/kg).
- The Baseline Risk Assessment (BRA) in the BCV Phase I RI stated: the sources of mercury and PCBs to the BCV fish are currently unknown.




#### **BCV Decision Documents – BCV Mercury Sources**

BCV OU2 ROD (DOE/OR/02-1435&D2) and BCV OU2 RI (DOE/OR/01-1273&D2)

- The BCV OU2 ROD identified two areas with mercury, SY-200 and Spoil Area 1. No other mention of mercury in the BCV OU2 ROD.
- The BCV OU2 RI identified mercury as a COC (human health) for SY-200. While Spoil Area 1 and the Rust Spoil Area had mercury as a contaminant of potential concern (COPC), the BRA did not identify mercury as a COC for these areas.
- The BCV OU2 RI indicated that mercury concentrations were elevated at SY-200 but were generally within an order of magnitude of background; however, free mercury was seen in some of the borings during the BCV OU2 RI.




## **Potential BCV Mercury Sources**





### **BCV Mercury Sources – SY-200 Yard**





- SY-200 Yard was a former equipment storage yard used to store nonradioactive contaminated equipment, mercury flasks, etc. from the 1950s to 1986
- In 1990, construction of the Environmental Support Facility began at the site. During construction, mercury was detected at high levels (up to 816 mg/kg) in excavated soils and visible mercury was noted
- Construction was paused and a 3 to 5 ft soil cover was placed across the site
- Bear Creek shown in blue
- Red area is high mercury area at SY-200 Yard from human health risk assessment
- Soil borings in the red area had visible mercury

(DOE/OR/01-1273&D2)



## **BCV Mercury Sources- Spoil Area 1**



- Spoil Area 1 was used for the disposal of what was characterized as uncontaminated construction debris from Y-12, but soil and groundwater studies confirmed the presence of heavy metals and radionuclides
- Mercury exceeded its MCL in a groundwater sample collected from a small intermittent seep near the base of the landfill; no constituents exceeded risk-based levels in surface water samples collected at the site
- Mercury was elevated in soil relative to background; however, mercury was not identified as a COC in soil in the BCV OU2 RI for Spoil Area 1

(DOE/OR/01-1273&D2, DOE/OR/01-2895&D2)



### **BCV Mercury Sources – NT-3**

NT-3 at the Boneyard/Burnyard (BYBY) is a remediated former strong mercury source

- BYBY was a visual cleanup
- Surface water sample in August 2020 had a mercury concentration (147 ng/L) above the AWQC
- It was concluded that it was a statistical outlier based on the available data and attributed to mercury adsorbed to suspended sediment (TDS 17.1 mg)

BEGINNING OF OIL LANDEARY

- · Bear Creek shown in dark blue
- NT-3 tributary in light blue
- BYBY outlined in black
- NT-03 sampling point shown as red star

(DOE/OR/01-2895&D2/V1)



## **BCV Mercury Sources - Others**

#### S-3 Pond Pathway 3 to NT-1

 Minor mercury contributions from mercury-contaminated fill materials (58-83 mg/kg in soil)


## Other sites (BCBG, OLF, HCDA, Sanitary Landfill 1, Bear Creek Road Debris Burial, Creekside Debris Burial, and Rust Spoil Area)

Minor mercury contributions

(DOE/OR/01-1455/V1&D2)

## UCOR

## **Previous Investigations - Current Surface Water** and **Biota Sampling in Bear Creek**





## **Previous Investigations - Current Surface Water and Biota Sampling in Bear Creek**


Media	Performance standard	Sampling frequency	Parameters	Monitoring Locations
Surface water	AWQC	Semi-annual (Q1 and Q3)	Total Hg and MeHg	BCK 3.3, BCK 4.55, BCK 9.2, BCK-11.54A, BCK 12.34, NT-03, NT-08, SS-4, and SS-5
		Semi-annual (Q2 and Q4) (year before FYR)	Total Hg	BCK 4.55, BCK 9.2, BCK 12.34, NT-03, and NT-08
		Annual (year before FYR)	Total Hg	BCK-07.87 and NT-01
		Annual (year before FYR)	МеНд	NT-05
	Trend monitoring	Quarterly	Bicarbonate, carbonate, chloride, fluoride, and sulfate	NT-01, NT-02, NT-03, SS-4, and SS-5
		Semi-annual	Bicarbonate, carbonate, chloride, fluoride, and sulfate	NT-07 and NT-08
		Quarterly (year before FYR)	Bicarbonate, carbonate, chloride, fluoride, and sulfate	NT-05
Water q	Water quality	Semi-annual	Total suspended solids and total dissolved solids.	NT-07 and NT-08
		Quarterly (year before FYR)	Total suspended solids and total dissolved solids.	BCK 9.2, BCK-07.87, BCK 4.55, NT-03, BCK-12.34, NT- 01, NT-05, and NT-08
Biota	Baseline sampling	Semi-annual	Hg and MeHg	BCK 3.3, BCK 9.9, and HCK 20.6 (whole-body stoneroller minnows and rock bass fillets) BCK 12.4 (whole-body stoneroller minnows)
		Annual (year before FYR)	Hg and MeHg	BCK-9.9 (whole body caddisflies)

#### A-2

## **Previous Investigations - Summary of Historical Surface Water Data**




Surface Water Hg & MeHg in Bear Creek, NT-3, and Hinds Creek (2011-2022)



## **Previous Investigations - Summary of Historical Biota Data**




- FY 2022 total mercury concentrations were well below the mercury AWQC (51 ng/L) in surface water
  - BCK 3.3 (3.59 ng/L), BCK 4.6 (2.34 ng/L), BCK 9.2 (4.78 ng/L),
     BCK 11.54A (9.76 ng/L), BCK 12.34 (10 ng/L), and NT-03 (19.7 ng/L)
  - Fish tissue concentrations in Bear Creek remain near the fish tissue criterion (0.3 μg/g).



## **Previous Investigations - Summary of Historical Sediment Data**





## **Previous Investigations - Summary of Historical Sediment Data**



Limited historical sediment data for Bear Creek are available in OREIS (primarily sampled 2011 and earlier)

- Zone 1: 7 locations, 6 in May 1995 and 1 in June 2005
  - Concentrations ranged from 0.14U 0.97 mg/kg total mercury
- Zone 2: 2 locations sampled in May 1995
  - Concentrations were ND and 0.16 mg/kg total mercury
- Zone 3: 20 locations sampled December 1993 April 2011
  - Concentrations ranged from 0.0189J 6.9 mg/kg total mercury

No methylmercury data are available for sediment with the exception of limited special studies data discussed later in this presentation.



Special studies of Bear Creek were conducted between 2017 and 2021 to better understand the biotic and abiotic factors contributing to mercury concentrations in fish in Bear Creek. The focus of these field studies was to gain an understanding of the processes controlling mercury methylation and bioaccumulation with beaver dams and periphyton being key areas of interest. Studies included:


- Understanding the role of beaver dams in contributing to mercury dynamics in Bear Creek (2017-2018)
- Evaluating the effects of fine-grained sediment deposition (2019)
- Investigation of the potential role that tributaries to Bear Creek may have on mercury and methylmercury in the main channel (2020-2021)
- Periphyton relationships (2021)


PAGE 22


## **Previous Investigations - Summary of Special Studies Data**



#### **Bear Creek Surface Water mercury**








- Total mercury and dissolved mercury concentrations decrease downstream in Bear Creek
- Higher concentrations in NT-03 and beaver pond but mass loading likely small


PAGE 23


## **Previous Investigations - Summary of Special Studies Data**



#### **Bear Creek Surface Water methylmercury**








- Total methylmercury, dissolved methylmercury, particulate methylmercury concentrations increase downstream during this study
- Higher concentrations in NT-03 and beaver pond but mass loading likely small
- Effect of beaver dam at BCK 7 evident



#### Total mercury in sediments (A) and periphyton (B) along Bear Creek




Flow is from left to right on each panel. The vertical dashed line marks the confluence of Bear Creek with EFPC.

- Sediment mercury concentration decreases downstream
- Effect of the former beaver dam at BCK 7 is evident
- Tributary sediments are comparable to Bear Creek
- Total mercury concentrations in periphyton in lower section of Bear Creek is higher than in the upper section
- Total mercury in periphyton is, on average, 1.4 times greater than in co-located sediment



#### Methylmercury in sediments (A) and periphyton (B) along Bear Creek




Flow is from left to right on each panel. The vertical dashed line marks the confluence of Bear Creek with EFPC.

- Sediment methylmercury concentration is variable with no strong spatial trend
- Effect of the former beaver dam at BCK 7 is evident
- Methylmercury in periphyton is higher in lower Bear Creek compared to upper
- Periphyton methylmercury in NT-3 and outlet of beaver pond is substantially greater than other locations
- Total methylmercury in periphyton is, on average, 5.6 times greater than in co-located sediment



#### Sulfate concentrations along Bear Creek



- Sulfate concentrations were consistent within the sampled reach but elevated relative to NT-09, NT-10, and NT-11
- The higher sulfate concentrations in lower Bear Creek coincide with relatively higher periphyton methylmercury concentrations in those sample locations




### **DQO Steps**

United States Environmental Protection Agency Office of Environmental Information Washington, DC 20460 EPA/240/B-06/001 February 2006



# Guidance on Systematic Planning Using the Data Quality Objectives Process

EPA QA/G-4



that meets the performance criteria



### **DQO Step 1. State the Problem**

There are insufficient data along Bear Creek and its tributaries to determine if there are potential sources of mercury and methylmercury in sediment and floodplain soils that may be contributing to exceedances of fish tissue criterion in prior years



### **DQO Step 2. Identify the Goals of the Study**

- Determine if there are areas (channel sediment, creek bank, and floodplain soils) along Bear Creek and its tributaries that are potential sources of mercury and methylmercury that may affect fish
- Obtain data from various hydrologic settings
   (i.e., pools, beaver ponds, etc.) that may contribute to
   mercury methylation and its bioaccumulation in the
   environment of Bear Creek and a reference location
   (e.g., Hinds Creek)



### **DQO Step 3. Identify Information Inputs**

- Review potential sources of mercury and methylmercury in Bear Creek and its tributaries
- Review existing historical biota, surface water, sediment, and special studies data in Bear Creek, its tributaries, and reference site
- Collect additional surface water, channel sediment, creek bank, and floodplain soils data along Bear
   Creek and its tributaries from selected transects
- Collect additional surface water, channel sediment, creek bank, and floodplain soils data from the reference site



### **DQO Step 4. Define the Study Area Boundary**

- Spatial
  - The study boundary is Bear Creek, its tributaries, and a reference location and limited surrounding creek bank sediment and floodplain soil
- Temporal
  - Collect samples in Fall 2023 to meet the RSE milestone of September 2024



#### **DQO Step 5. Develop the Analytical Approach**

- Collect surface water, channel sediment, creek bank, and floodplain soils to determine mercury and methylmercury concentrations along Bear Creek, its tributaries, and the reference site
- Analytical parameters will include mercury, methylmercury, nutrients (e.g., sulfate, phosphate, nitrogen, organic carbon, etc.), particle size analysis (PSA), and mercury speciation at select locations
- Transects will be field-located in potential source areas and pool areas (e.g., upstream of beaver dams) in Bear Creek and the mouths of tributaries (e.g., NT-3) based on a reconnaissance survey



# **DQO Step 6. Specify the Performance or Acceptance Criteria**

 New data will be obtained under UCOR/RSI approved procedures and quality programs and will be archived in OREIS.





#### **DQO Step 7. Develop the Plan for Obtaining Data**

- Compile all available information on potential mercury and methylmercury sources, existing surface water, sediment, and biota data (BMAP)
- Evaluate existing data
- Conduct additional characterization fieldwork:
  - Identify locations to conduct surface water, channel sediment, creek bank, and floodplain soil transect sampling.
  - Identify reference site location for surface water, channel sediment, creek bank, and floodplain soil sampling.




## **Proposed Transect Locations**

- Proposed transects along Bear Creek are based on the following\*:
  - Locations downgradient and in the vicinity of potential source areas
  - Locations where sampling for biota and surface water have historically occurred
  - Locations in the vicinity of beaver ponds

<sup>\*</sup>Exact transect locations are subject to change based on access and other field factors.




### **Proposed Transect Locations**





### **Beaver Dams near BCK 7.0**





## **Transect Sampling**

### Channel sediment

 Collect 1-2 samples of channel sediment (0 – 0.5 ft) at each transect (number of samples at each transect will be based on width of Bear Creek at each location)

### Creek bank sediment

 Collect 2 samples of bank sediment (0 – 0.5 ft) at each transect (one on each bank)

### Floodplain soil


 Collect 2 samples of floodplain soil (0 – 0.5 ft) in the vicinity of Bear Creek (one on each side of Bear Creek)

### Surface water

Collect 1 surface water sample at each transect



### **Transect Sampling Diagram**





## **Analytical Suite**

- Analyze channel sediment, creek bank, and floodplain soils for mercury, methylmercury, nutrients, PSA, and organic carbon
- Analyze surface water for mercury, methylmercury, nutrients, and organic carbon
- Additional mercury speciation partitioning in select sampling transects based on hydrologic setting



### **RSE Schedule**

Activity	Date(s)
Historical Data Review and DQO Preparation	May/June 2023
DQO Meeting	June 2023
Prepare and Review RSE Sampling and Analysis Plan (SAP; FFA secondary document)	July/August 2023
Submit RSE SAP	August 2023
Perform RSE Sediment and Surface Water Sampling	September - November 2023
Data Evaluation (SED, SW, and 2023 Fish Tissue)	January - March 2024
Prepare and Review RSE D1	April - September 2024
Submit BCV Mercury Sources RSE D1	FFA App E: September 30, 2024

This page intentionally left blank.

# APPENDIX B. BEAR CREEK VALLEY MERCURY SOURCES REMEDIAL SITE EVALUATION SAMPLING AND ANALYSIS PLAN TABLES

This page intentionally left blank.

Table B.1. Sample groups for the BCV mercury sources RSE

Sample group <sup>a</sup>	Location <sup>b</sup>	Sampling point <sup>b</sup>	Monitoring frequency <sup>c</sup>	Matrix <sup>d</sup>	Sample type <sup>e</sup>	<b>Dup</b> <sup>f</sup>	Analyte/parameter group <sup>g</sup>
		BCT1-BSL	<u> </u>	SO	С		S-BCVRSE
	DOTAL CONTRACT OF STREET	BCT1-BSU	1 ! !	SO	С	} ! !	S-BCVRSE
	BCT1 (upstream of BCK 0.6; downstream of beaver dam)	BCT1-CH	Q1	SE	С		S-BCVRSE
	beaver dam)	BCT1-FP		SO	С		S-BCVRSE
		BCT1-SW	1 ! !	WS	G		BCVRSE(+F)
		BCT2-BSL		SO	С	]   	S-BCVRSE
	DCT2 ( , , , , , , , , , , , , , , , , , ,	BCT2-BSU		SO	С		S-BCVRSE
	BCT2 (upstream of BCK 0.6; upstream of beaver dam)	ВСТ2-СН	Q1	SE	С		S-BCVRSE
	beaver dam)	BCT2-FP	]   	SO	С		S-BCVRSE
LOWBCV		BCT2-SW	] 	WS	G		BCVRSE(+F)
LOWBEV		BCT3-BSL	] 	SO	С		S-BCVRSE
	DCT2 (1	BCT3-BSU		SO	С		S-BCVRSE
	BCT3 (downstream of BCK 3.3; downstream of beaver dam)	ВСТ3-СН	Q1	SE	C		S-BCVRSE
	downstream of beaver damy	BCT3-FP		SO	С		S-BCVRSE
		BCT3-SW	] !	WS	G		BCVRSE(+F)
		BCT4-BSL		SO	C		S-BCVRSE
	DCT4 (1transf-DCV-2-2	BCT4-BSU	]   	SO	C		S-BCVRSE
	BCT4 (downstream of BCK 3.3; upstream of beaver dam)	ВСТ4-СН	Q1	SE	С		S-BCVRSE
	beaver dam,	BCT4-FP	] }	SO	C		S-BCVRSE
	İ	BCT4-SW	i 	WS	G		BCVRSE(+F)
		BCT5-BSL	 	SO	C		S-BCVRSE
		BCT5-BSU		SO	C		S-BCVRSE
	BCT5 (downstream of BCK 4.55;	ВСТ5-СН	Q1	SE	C		S-BCVRSE, HGSEQ
	downstream of beaver dam)	BCT5-FP	Q1	SO	C		S-BCVRSE, HGSEQ
		BCT5-SW	<u> </u>	WS	G	<u> </u> 	BCVRSE(+F)
BCV ZONE 1		BCT5-BS	<u> </u>	SO	C	   	HGSEQ
DC V ZOIVE I		BCT6-BSL	]   	SO	C		S-BCVRSE
		BCT6-BSU		SO	C		S-BCVRSE
	BCT6 (downstream of BCK 4.55; upstream	ВСТ6-СН	Q1	SE	С	   	S-BCVRSE, HGSEQ
	of beaver dam)	BCT6-FP	Ųι	SO	С	     	S-BCVRSE, HGSEQ
		BCT6-SW		WS	G	, i i !	BCVRSE(+F)
		BCT6-BS	! !	SO	С	! !	HGSEQ

Table B.1. Sample groups for the BCV mercury sources RSE (cont.)

Sample group <sup>a</sup>	Location <sup>b</sup>	Sampling point <sup>b</sup>	Monitoring frequency <sup>c</sup>	Matrix <sup>d</sup>	Sample type <sup>e</sup>	Dup <sup>f</sup>	Analyte/parameter group <sup>g</sup>
		BCT7-BSL	I I I	SO	С	! ! !	S-BCVRSE
		BCT7-BSU	] !	SO	С	] !	S-BCVRSE
	BCT7 (downstream of BCK 7.87 at the confluence of NT-13/Bear Creek; downstream of westernmost beaver dam)	ВСТ7-СН	Q1	SE	С		S-BCVRSE, HGSEQ
		BCT7-FP	Q1	SO	С	 	S-BCVRSE, HGSEQ
	de militare de mesterimiest seu en dumi,	BCT7-SW	] <u>!</u> <u>!</u>	WS	G	: :	BCVRSE(+F)
		BCT7-BS	] !	SO	С	] 	HGSEQ
		BCT8-BSL	 	SO	С	ļ	S-BCVRSE
		BCT8-BSU		SO	С	i i	S-BCVRSE
RCV ZONE 1	BCT8 (downstream of BCK 7.87 at the confluence of NT-13/Bear Creek; upstream	ВСТ8-СН	Q1	SE	С		S-BCVRSE, HGSEQ
BC V ZONE I	of westernmost beaver damn)	BCT8-FP	Q1	SO	C	! ! !	S-BCVRSE, HGSEQ
į	or westerminost seaver dammy	BCT8-SW	] !	WS	G	} !	BCVRSE(+F)
		BCT8-BS		SO	С	<u> </u>	HGSEQ
		BCT9-BSL	] 	SO	С		S-BCVRSE
	BCT9 (downstream of BCK 7.87; upstream of two beaver dams; southeast of Reeves	BCT9-BSU		SO	С	   	S-BCVRSE
		ВСТ9-СН	Q1	SE	С		S-BCVRSE, HGSEQ
	Road/Haul Road)	BCT9-FP	Q1	SO	С		S-BCVRSE, HGSEQ
		BCT9-SW	]   	WS	G		BCVRSE(+F)
		BCT9-BS		SO	С		HGSEQ
		BCT10-BSL	]	SO	С		S-BCVRSE
	BCT10 (downstream of surface water	BCT10-BSU		SO	С		S-BCVRSE
BCV ZONE 2	integration point BCK 9.2; upstream of	BCT10-CH	Q1	SE	С		S-BCVRSE
	EMDF)	BCT10-FP		SO	С	]	S-BCVRSE
		BCT10-SW	] 	WS	G		BCVRSE(+F)
		BCT11-BSL	] 	SO	C	 	S-BCVRSE
		BCT11-BSU	¦ 	SO	С	<b></b>   	S-BCVRSE
BCV ZONE 3	BCT11 (upstream of NT-8 at BCK 9.9)	ВСТ11-СН	Q1	SE	С	<b>-</b>     	S-BCVRSE, HGSEQ
BC V ZONE 3	De 111 (upsucam of 1v1-6 at DCK 9.9)	BCT11-FP	ζ,	SO	С	, <b></b> i i	S-BCVRSE, HGSEQ
		BCT11-SW	i ¦	WS	G	i i !	BCVRSE(+F)
		BCT11-BS	! ! !	SO	С	   	HGSEQ

Table B.1. Sample groups for the BCV mercury sources RSE (cont.)

Sample group <sup>a</sup>	Location <sup>b</sup>	Sampling point <sup>b</sup>	Monitoring frequency <sup>c</sup>	Matrix <sup>d</sup>	Sample type <sup>e</sup>	<b>Dup</b> <sup>f</sup>	Analyte/parameter group <sup>g</sup>
		BCT12A-BSL		SO	С	<u> </u>	S-BCVRSE
	DOTION ( CNT 2 C 34	BCT12A-BSU		SO	С	] !	S-BCVRSE
	BCT12A (upstream of NT-3 confluence with Bear Creek)	BCT12A-CH	Q1	SE	С		S-BCVRSE
	Bear Creek)	BCT12A-FP		SO	С		S-BCVRSE
		BCT12A-SW		WS	G	 	BCVRSE(+F)
		BCT12B-BSL		SO	C	ļ	S-BCVRSE
		BCT12B-BSU		SO	С	j	S-BCVRSE
	BCT12B (downstream of BYBY at the	BCT12B-CH	Q1	SE	С	X	S-BCVRSE, HGSEQ
	confluence of NT-3/Bear Creek)	BCT12B-FP	Q1	SO	C	A	S-BCVRSE, HGSEQ
		BCT12B-SW		WS	G	}	BCVRSE(+F)
		BCT12B-BS		SO	C	<u> </u>	HGSEQ
		BCT13-BSL		SO	C		S-BCVRSE
	BCT13 (upstream of BYBY, EMWMF, and	BCT13-BSU	]   	SO	C		S-BCVRSE
BCV ZONE 3	NT-3)	ВСТ13-СН	Q1	SE	C	! !	S-BCVRSE
		BCT13-FP		SO	C		S-BCVRSE
		BCT13-SW	i 	WS	G		BCVRSE(+F)
		BCT14-BSL		SO	C	<u>[</u>	S-BCVRSE
		BCT14-BSU		SO	С	į	S-BCVRSE
	BCT14 (downstream of SY-200 Yard, Spoil	BCT14-CH	Q1	SE	C	X	S-BCVRSE, HGSEQ
	Area 1, and S-3 Ponds Site)	BCT14-FP	Q1	SO	C	Λ	S-BCVRSE, HGSEQ
		BCT14-SW	i i	WS	G	]	BCVRSE(+F)
		BCT14-BS		SO	С	! ! !	HGSEQ
		BCT15-BSL		SO	C	<u> </u>	S-BCVRSE
		BCT15-BSU		SO	C	i ! J	S-BCVRSE
	BCT15 (downstream of S-3 Ponds Site)	BCT15-CH	Q1	SE	С	! ! {	S-BCVRSE
		BCT15-FP		SO	C	! ! !	S-BCVRSE
		BCT15-SW		WS	G	<u> </u>	BCVRSE(+F)
		HCTREF-BSL	 ! !	SO	С	ļ	S-BCVRSE
		HCTREF-BSU	! ! !	SO	C	   	S-BCVRSE
Hinds Creek	HCTREF	HCTREF-CH	Q1	SE	С	! !	S-BCVRSE, HGSEQ
Timas Creek	(HCK 20.6 reference site)	HCTREF-FP	ν,	SO	С	i 	S-BCVRSE, HGSEQ
		HCTREF-SW	! !	WS	G	<u> </u>	BCVRSE(+F)
		HCTREF-BS		SO	C	<u> </u>	HGSEQ

#### Table B.1. Sample groups for the BCV mercury sources RSE (cont.)

Sample group BCV = Bear Creek Valley Watershed sample group number LOWBCV = Lower Bear Creek Valley Samples in each group will be collected during as short a time period as possible, following the schedule provided Location and sampling point BCK = Bear Creek kilometer EMWMF = Environmental Management Waste Management Facility FP = floodplain soil BCT = Bear Creek transect BS = creek bank soil HCK = Hinds Creek BSL = creek bank soil (lower) HCTREF = Hinds Creek transect reference site BSU = creek bank soil (upper) NT = northern tributaryBYBY = Boneyard/Burnyard SW = surface water CH = channel sediment SY = scrap yardEMDF = Environmental Management Disposal Facility Monitoring frequency Q = quarter of the fiscal year (e.g., Q1, Q2, Q3, Q4) Matrix SE = sedimentWS = surface waterSO = soilSample type G = grab sampleC = composite sample

<sup>j</sup> Duplicate

X = field duplicate sample will be collected for all analyte/parameter groups except HGSEQ

Field duplicates samples will be collected concurrently with the investigative samples and sent to the laboratory responsible for analyses of the investigative sample. Field duplicates will be collected at a frequency of 10% of the samples collected (i.e., 1 to 10 total samples collected equal 1 field duplicate; 1 to 20 total samples collected equal 2 field duplicates) or as specified in the task-specific work control document (e.g., Sampling and Analysis Plans [SAPs]), in accordance with the *Quality Assurance Project Plan for the Water Resources Restoration Program, U.S. Department of Energy, Oak Ridge Reservation, Oak Ridge, Tennessee* (UCOR-4049; Water Resources Restoration Program [WRRP] Quality Assurance Project Plan [QAPP]). Deviations from this Remedial Site Evaluation (RSE) SAP will be documented in the field logbook and in the BCV Mercury Sources RSE. Changes will be documented, as appropriate in the field, as well as in the Project Environmental Measurements System

g Analyte/parameter group

See Tables D.56 through D.58 in the WRRP QAPP for a list of parameter groups and analytes

BCVRSE(+F) = Both a filtered and unfiltered sample are obtained by sampling personnel for the designated metals analysis to be performed by the laboratory. Otherwise, only an unfiltered sample is obtained and analyzed for metals

WRRP									Sc	reening lev	vels <sup>d</sup>		
$QAPP^a$	Analyte	CAS	Analytical method <sup>b</sup>	Units	·	Laboratory	$RRL^c$	Surface water					
parameter group	111111,11	number	ramay cour moulou		MDL	PQL	1442	DWS	CCC	CMC	W&O	ООС	
BCVRSE				Field paran	neters				1				
	Water temperature	NA	NA	°C	i	i	i i		i	i			
	Dissolved oxygen	7782-44-7	NA	mg/L	L 	   			L				
	Turbidity	NA	NA	NTU									
	рН	NA	NA	рН	∱   	<del> </del>   	 		; 	}   	<del> </del>		
	Specific conductance (conductivity)	NA	NA	μmhos/cm					 				
	Oxidation-reduction potential	NA	NA	mV	 	 	 		 	 			
				Metals	5	•				•			
	Mercury	7439-97-6	EPA-1631	ng/L	0.08	0.5	0.5	2000	770	1400	50	51	
	Methylmercury	22967-92-6	EPA-1630	ng/L	0.026	0.08	0.02						
	Aluminum	7429-90-5	SW846-6010 or SW846-6020	mg/L	0.0193	0.05	0.05						
	Antimony	7440-36-0	SW846-6010 or SW846-6020	mg/L	0.001	0.003	0.003	0.006		<u> </u>	0.0056	0.64	
	Arsenic	7440-38-2	SW846-6010 or SW846-6020	mg/L	0.005	0.03	0.005	0.01	0.15	0.34	0.01	0.01	
	Barium	7440-39-3	SW846-6010 or SW846-6020	mg/L	0.001	0.005	0.005	2					
	Beryllium	7440-41-7	SW846-6010 or SW846-6020	mg/L	0.0002	0.0005	0.001	0.004					
	Boron	7440-42-8	SW846-6010 or SW846-6020	mg/L	0.0052	0.015	0.01		ļ				
	Cadmium	7440-43-9	SW846-6010 or SW846-6020	mg/L	0.0003	0.001	0.00013	0.005	0.00072	0.0018			
	Calcium	7440-70-2	SW846-6010 or SW846-6020	mg/L	0.05	0.2	0.01						
	Chromium	7440-47-3	SW846-6010 or SW846-6020	mg/L	0.003	0.01	0.005	0.1	0.074	0.57			
	Cobalt	7440-48-4	SW846-6010 or SW846-6020	mg/L	0.0003	0.001	0.005			<u> </u>			
	Copper	7440-50-8	SW846-6010 or SW846-6020	mg/L	0.0003	0.002	0.005		0.009	0.013			
	Iron	7439-89-6	SW846-6010 or SW846-6020	mg/L	0.033	0.1	0.01						
	Lead	7439-92-1	SW846-6010 or SW846-6020	mg/L	0.0005	0.002	0.002	0.005	0.0025	0.065			
	Lithium	7439-93-2	SW846-6010 or SW846-6020	mg/L	0.003	0.01	0.01						
	Magnesium	7439-95-4	SW846-6010 or SW846-6020	mg/L	0.11	0.3	0.05						
	Manganese	7439-96-5	SW846-6010 or SW846-6020	mg/L	0.001	0.005	0.005						
	Nickel	7439-98-7	SW846-6010 or SW846-6020	mg/L	0.0006	0.002	0.01	0.1	0.052	0.47	0.61	4.6	
	Potassium	7440-02-0	SW846-6010 or SW846-6020	mg/L	0.05	0.15	0.025						
	Selenium	7440-09-7	SW846-6010 or SW846-6020	mg/L	0.006	0.03	0.0025	0.05	0.0031	0.02	0.17	4.2	
	Silicon	7782-49-2	SW846-6010 or SW846-6020	mg/L	0.025	0.1	0.01						
	Silver	7440-22-4	SW846-6010 or SW846-6020	mg/L	0.0003	0.001	0.0015			0.0032			
	Sodium	7440-23-5	SW846-6010 or SW846-6020	mg/L	0.1	0.3	0.01			<u></u>			
	Strontium	7440-24-6	SW846-6010 or SW846-6020	mg/L	0.002	0.01	0.005						
	Thallium	7440-28-0	SW846-6010 or SW846-6020	mg/L	0.0006	0.002	0.001	0.002			0.00024	0.00047	
	Uranium	7440-61-1	SW846-6010 or SW846-6020	mg/L	0.000067	0.0002	0.004			<u> </u>			
	Vanadium	7440-62-2	SW846-6010 or SW846-6020	mg/L	0.001	0.005	0.01			<u> </u>			
	Zinc	7440-66-6	SW846-6010 or SW846-6020	mg/L	0.0033	0.02	0.01		0.12	0.12	7.4	26	

Table B.2. Analytes, RRLs, and screening levels for water for the BCV mercury sources RSE (cont.)

WRRP								Screening levels <sup>d</sup>						
QAPP <sup>a</sup>	Analyte	CAS number	Analytical method <sup>b</sup>	Units	Laboratory MDL	Laboratory	$\mathbf{RRL}^c$		Surface water					
parameter group		number			MIDL	PQL		DWS	CCC	CMC	W&O	ООС		
BCVRSE			Mi	scellaneous p	arameters									
(cont)	Phosphorus (total)	7723-14-0	SW846-6020	mg/L	0.018	0.05	0.05		ļ					
	Total organic carbon	E701250	SW846-9060	mg/L	0.33	1	1			<u> </u>				
	Dissolved organic carbon	E701250	SW846-9060	mg/L	0.33	1	1							
	Total dissolved solids	E1642222	SM-2540 C	mg/L	3.4	5	10	500	 					
	Total suspended solids	E1642818	SM-2540 D	mg/L	1.14	5	5			<u> </u>				
				Anion	S									
	Chloride	16887-00-6	EPA-300.0	mg/L	0.067	0.2	0.01		<u> </u>					
	Fluoride	16984-48-8	EPA-300.0	mg/L	0.033	0.1	0.05			I				
	Sulfate	14808-79-8	EPA-300.0	mg/L	0.133	0.4	0.1							
	Sulfide	18496-25-8	SM-4500-S2 D	mg/L	0.033	0.1	0.1			Ī				
	Nitrate-nitrite as nitrogen	E701177	EPA-353.2	mg/L	0.017	0.05	0.1	10		<del></del>				

<sup>&</sup>lt;sup>a</sup>UCOR-4049. Ouality Assurance Project Plan for the Water Resources Restoration Program, U.S. Department of Energy, Oak Ridge Reservation, Oak Ridge, Tennessee, latest revision, United Cleanup Oak Ridge, TN.

-- = not available or not applicable

BCV = Bear Creek Valley

CAS = Chemical Abstracts Service

CCC = Tennessee Department of Environment and Conservation fish and aquatic life criterion continuous concentration criteria, Chapter 1200-4-3-.03(3)

CMC = Tennessee Department of Environment and Conservation fish and aquatic life criterion maximum concentration criteria, Chapter 1200-4-3-.03(3)

DWS = Tennessee Department of Environment and Conservation domestic water supply criteria, Chapter 1200-4-3-.03(1).

EPA = U.S. Environmental Protection Agency

MDL = method detection limit

NA = not applicable

NTU = nephelometric turbidity unit

OOC = Tennessee Department of Environment and Conservation recreation organisms only criteria, Chapter 1200-4-3-.03(4)

PQL = practical quantitation limit

QAPP = Quality Assurance Project Plan

RRL = requested reporting limit

RSE = remedial site evaluation

SW846 = EPA test methods for evaluating solid waste, physical/chemical methods

W&O = Tennessee Department of Environment and Conservation recreation water and organisms criteria, Chapter 1200-4-3-.03(4)

WRRP = Water Resources Restoration Program

<sup>&</sup>lt;sup>b</sup> An alternative method or alternate technique may be used to achieve the RRLs.

<sup>&</sup>lt;sup>c</sup>RRLs are defined so that the data obtained meet program/project requirements for reporting quantitative data. For this parameter group, the laboratory is being requested to report detections with respect to the MDLs, which are generally lower than the RRLs.

<sup>&</sup>lt;sup>d</sup> Screening levels listed here are for potential comparison purposes only and are not required performance goals.

Table B.3. Analytes, RRLs, and screening levels for sediment and soil for the BCV mercury sources RSE

WRRP  QAPP <sup>a</sup> parameter  group	Analyte	CAS number	Analytical method <sup>b</sup>	Units	Laboratory MDL	Laboratory PQL	$RRL^c$	Screening level
S-BCVRSE				Metals	l.	l		
	Mercury	7439-97-6	SW846-7471	mg/kg	0.009	0.03	0.1	Compare to reference site
	Methylmercury	22967-92-6	EPA-1630 (modified)	ng/g	0.017	0.058	0.017	Compare to reference site
	Aluminum	7429-90-5	SW846-6010 or SW846-6020	mg/kg	6.8	20	1	Compare to reference site
	Antimony	7440-36-0	SW846-6010 or SW846-6020	mg/kg	0.33	2	0.5	Compare to reference site
	Arsenic	7440-38-2	SW846-6010 or SW846-6020	mg/kg	0.338	1	0.5	Compare to reference site
	Barium	7440-39-3	SW846-6010 or SW846-6020	mg/kg	0.1	0.5	0.5	Compare to reference site
	Beryllium	7440-41-7	SW846-6010 or SW846-6020	mg/kg	0.1	0.5	0.1	Compare to reference site
	Boron	7440-42-8	SW846-6010 or SW846-6020	mg/kg	1	5	1	Compare to reference site
	Cadmium	7440-43-9	SW846-6010 or SW846-6020	mg/kg	0.02	0.2	0.1	Compare to reference site
	Calcium	7440-70-2	SW846-6010 or SW846-6020	mg/kg	8	25	5	Compare to reference site
	Chromium	7440-47-3	SW846-6010 or SW846-6020	mg/kg	0.15	1	0.5	Compare to reference site
	Cobalt	7440-48-4	SW846-6010 or SW846-6020	mg/kg	0.15	0.5	0.5	Compare to reference site
	Copper	7440-50-8	SW846-6010 or SW846-6020	mg/kg	0.3	2	0.5	Compare to reference site
	Iron	7439-89-6	SW846-6010 or SW846-6020	mg/kg	8	25	1	Compare to reference site
	Lead	7439-92-1	SW846-6010 or SW846-6020	mg/kg	0.33	2	0.3	Compare to reference site
	Lithium	7439-93-2	SW846-6010 or SW846-6020	mg/kg	0.4	2	1	Compare to reference site
	Magnesium	7439-95-4	SW846-6010 or SW846-6020	mg/kg	8.5	30	5	Compare to reference site
	Manganese	7439-96-5	SW846-6010 or SW846-6020	mg/kg	0.2	1	0.5	Compare to reference site
	Molybdenum	7439-98-7	SW846-6010 or SW846-6020	mg/kg	0.2	1	1	Compare to reference site
	Nickel	7440-02-0	SW846-6010 or SW846-6020	mg/kg	0.15	0.5	1	Compare to reference site
	Potassium	7440-09-7	SW846-6010 or SW846-6020	mg/kg	6.4	25	5	Compare to reference site
	Selenium	7782-49-2	SW846-6010 or SW846-6020	mg/kg	0.36	1	0.5	Compare to reference site
	Silver	7440-22-4	SW846-6010 or SW846-6020	mg/kg	0.1	0.5	0.5	Compare to reference site
	Sodium	7440-23-5	SW846-6010 or SW846-6020	mg/kg	7	25	5	Compare to reference site
	Thallium	7440-28-0	SW846-6010 or SW846-6020	mg/kg	0.14	0.4	0.2	Compare to reference site
	Uranium	7440-61-1	SW846-6010 or SW846-6020	mg/kg	0.0132	0.04	5	Compare to reference site

Table B.3. Analytes, RRLs, and screening levels for sediment and soil for the BCV mercury sources RSE (cont.)

WRRP QAPP <sup>a</sup> parameter group	Analyte	CAS number	Analytical method <sup>b</sup>	Units	Laboratory MDL	Laboratory PQL	$RRL^c$	Screening level
	Vanadium	7440-62-2	SW846-6010 or SW846-6020	mg/kg	0.1	0.5	1	Compare to reference site
	Zinc	7440-66-6	SW846-6010 or SW846-6020	mg/kg	0.4	2	0.5	Compare to reference site
S-BCVRSE		•	Miscellar	eous param	eters			
(cont)	Total organic carbon	E701250	SW846-9060	mg/kg	200	500	1	
	Particle size analysis	NA	ASTM-D6913		   			†
		-		Anions				-
	Chloride	16887-00-6	SW846-9056	mg/kg	0.72	2	0.72	
	Fluoride	16984-48-8	SW846-9056	mg/kg	0.34	1	0.34	†
	Nitrate	14797-55-8	SW846-9056	mg/kg	0.33	1	0.33	
	Nitrite	14797-65-0	SW846-9056	mg/kg	0.33	1	0.33	†
	Sulfate	14808-79-8	SW846-9056	mg/kg	1.33	4	1.33	
	Sulfide	18496-25-8	SW846-9030/9034	mg/kg	9	25	9	
HGSEQ	Mercury (F0)	NA	SW846-3200 (modified)	ug/kg	3.6	21	3.6	
	Mercury (F1)	NA	SW846-3200 (modified)	ug/kg	240	740	240	†
	Mercury (F2)	NA	SW846-3200 (modified)	ug/kg	240	740	240	T
	Mercury (F3)	NA	SW846-3200 (modified)	ug/kg	240	740	240	
	Mercury (F4)	NA	SW846-3200 (modified)	ug/kg	240	740	240	
	Mercury (F5)	NA	SW846-3200 (modified)	ug/kg	41	370	41	 
	Mercury (F6)	NA	SW846-3200 (modified)	ug/kg	0.11	1.2	0.11	

<sup>&</sup>lt;sup>a</sup> UCOR-4049. Quality Assurance Project Plan for the Water Resources Restoration Program, U.S. Department of Energy, Oak Ridge Reservation, Oak Ridge, Tennessee, latest revision, United Cleanup Oak Ridge, TN.

ASTM = American Standard Test Method

BCV = Bear Creek Valley

CAS = Chemical Abstracts Service

EPA = U.S. Environmental Protection Agency

MDL = method detection limit

<sup>&</sup>lt;sup>b</sup> An alternative method or alternate technique may be used to achieve the RRLs.

<sup>&</sup>lt;sup>c</sup> RRLs are defined so that the data obtained meet program/project requirements for reporting quantitative data. For this parameter group, the laboratory is being requested to report detections with respect to the MDLs, which are generally lower than the RRLs.

<sup>-- =</sup> not available or not applicable

#### Table B.3. Analytes, RRLs, and screening levels for sediment and soil for the BCV mercury sources RSE (cont.)

Mercury (F0) = volatile elemental mercury

Mercury (F1) = water soluble mercury

Mercury (F2) = pH2 soluble mercury

Mercury (F3) = 1N potassium hydroxide extractable mercury

Mercury (F4) = 12N nitric acid soluble mercury

Mercury (F5) = aqua regia soluble mercury residue

Mercury (F6) = mineral-bound mercury

NA = not available

PQL = practical quantitation limit

QAPP = Quality Assurance Project Plan

RRL = requested reporting limit

RSE = remedial site evaluation

SW846 = EPA test methods for evaluating solid waste, physical/chemical methods

WRRP = Water Resources Restoration Program

This page intentionally left blank.

#### DOE/OR/01-2958&D2

### RECORD COPY DISTRIBUTION

File—DMC—RC

# U.S. ENVIRONMENTAL PROTECTION AGENCY COMMENT RESPONSE FORM



Document Number: DOE/OR/01-2958&D1	Document Title: Bear Creek Valley Mercury Sources Remedial Site Evaluation Sampling and Analysis Pla	n, Oak Ridge, Tennessee
Name of Reviewer:	Organization:	Date Comments
Jana Dawson	U.S. Environmental Protection Agency	Transmitted:
		November 15, 2023

Comment No.	Sect./Page	Comment	Response
1	Section 2.2.2/Page 3	Section 2.2.2 (Summary of Mercury Source Areas) page 3 states "The baseline risk assessment (BRA) in the BCV RI stated "the sources of mercury and PCBs to the BCV fish are currently unknown." While it was noted in the meeting minutes that DOE does not believe PCB discharge from the EMDF will be an issue with regards to attainment of Ambient Water Quality Criteria (AWQC) in Bear Creek for PCBs, it would seem appropriate and cost effective to also conduct PCB analyses for the surface water and fish tissue samples that will be collected as part of this mercury source investigation in order to identify the source of both mercury and PCBs into Bear Creek. EPA strongly recommends adding the PCB analyses to this sampling effort.	Clarification. Analysis of PCBs in surface water and fish is already conducted in Bear Creek as part of the annual routine and Five-Year Review (FYR) sampling, as outlined in the BCV RAR CMP and as reported in the annual Remediation Effectiveness Report and FYR.  The EMDF ROD mercury-management approach is the catalyst to this focused RSE to recognize if there are non-point source reductions that could offset the point source discharge at EMDF.  As stated in Section 3.1 (Data Quality Objective Step 1: State the Problem), there are insufficient data along Bear Creek and its tributaries to determine if there are potential sources of mercury and methylmercury in channel sediment and creek bank and floodplain soils that may be contributing to exceedances of fish tissue criterion in prior years.  As stated in Section 3.2 (Data Quality Objective Step 2: Identify the Goals of the Study), the purposes of this RSE are to:  1. Determine if there are areas (channel sediment and creek bank and floodplain soils) along Bear Creek and its tributaries that are potential sources of mercury and methylmercury that may affect fish.  2. Obtain data from various hydrologic settings (pools, beaver ponds, etc.) that may contribute to mercury methylation and its bioaccumulation in the environment of Bear Creek and a reference location (e.g., Hinds Creek).



Comment No.	Sect./Page	Comment	Response
			No change to the document proposed.
2	Section 2.2.2/Page 3	Section 2.2.2 (Summary of Mercury Source Areas) page 3 states "The BCV OU2 RI indicated mercury concentrations were elevated at the SY-200 Yard but were generally within an order of magnitude of background; however, free mercury was seen in some of the borings during the BCV OU2 RI. The BCV OU2 ROD identified the SY-200 Yard as the area with mercury." Please describe how much certainty there is that the location of the observed free mercury in soil borings was from soil bores collected at the SY-200 Yard since the SAP indicates there was initially some uncertainty about where the observations of free mercury were identified.	Clarification. The second paragraph in Section 2.2.2 was revised as follows: "The BCV OU2 RI indicated mercury concentrations were elevated at the SY-200 Yard but were generally within an order of magnitude of background; however, free mercury was seen in some of the borings at the SY-200 Yard during the BCV OU2 RI."
3	Section 4	Section 4 (Sampling and Analysis Plan – Sample Location Selection) does not propose any soil samples near suspected or known mercury source areas. By only sampling Bear Creek transects and associated bank/floodplain soils immediately next to transect location, how will it be determined which of the potential source areas are contributing mercury to Bear Creek. For example, a transect is proposed at BCT-14, which is stated to be downstream of the SY-200 Yard, Spoil Area 1, and S-3 Ponds Site. If elevated mercury is identified at this transect, how will it be determined which of the areas - SY-200 Yard area, the Soil Area 1, and/or S-3 Ponds are the main contributors of mercury to Bear Creek? The SAP does not appear to include sufficient samples to identify specific source areas of mercury. Please provide a response and/or SAP edits to state how specific source areas will be identified, and further, how such areas may be delineated to identify the extent of mercury migrating to Bear Creek, if elevated mercury is detected at any of the transects/ mercury-source investigation samples.	Clarification. As stated in Section 3.2 (Data Quality Objective Step 2: Identify the Goals of the Study), the purposes of this RSE are to:  1. Determine if there are areas (channel sediment and creek bank and floodplain soils) along Bear Creek and its tributaries that are potential sources of mercury and methylmercury that may affect fish.  2. Obtain data from various hydrologic settings (pools, beaver ponds, etc.) that may contribute to mercury methylation and its bioaccumulation in the environment of Bear Creek and a reference location (e.g., Hinds Creek).  As stated in Section 3.1 (Data Quality Objective Step 1: State the Problem), there are insufficient data along Bear Creek and its tributaries to determine if there are potential sources of mercury and methylmercury in channel sediment and creek bank and floodplain soils that may be contributing to exceedances of fish tissue criterion in prior years.  Data collected as part of this RSE will determine if there is any evidence these potential source areas (SY-200 Yard, Spoil Area 1, and S-3 Ponds Site) are contributing mercury to Bear Creek. If elevated mercury is detected in sediment or soil at transect



Comment No.	Sect./Page	Comment	Response
			BCT-14, downgradient of these source areas, then additional evaluation may be required, as discussed in the mercury-management approach in the EMDF ROD.
			During field reconnaissance for BCV Mercury Sources RSE sample transect locations, an additional transect was included in the planned work. Sample transect BCT-12 was split into BCT-12A (upstream of NT-3) and BCT-12B (downstream of NT-3) to further refine data resolution at a formerly remediated area (BYBY). Revised Figure 4.1 is attached at the end of these comment responses. Revisions to the BCV Mercury Sources RSE SAP to reflect this change have been made, as applicable.



### Attachment






Figure 4.1. BCV transect sampling locations. (Revised based on EPA comment 3 and TDEC general comment 4)





Document Number:	Document Title:	
DOE/OR/01-2958&D1	Bear Creek Valley Mercury Sources Remedial Site Evaluation Sampling and Analysis Plan, Oak Ridge, Tennessee	
Name of Reviewer:	Organization:	Date Comments
Dana Casey/Cody Juneau	Tennessee Department of Environment and Conservation	Transmitted:
		December 5, 2023

Comment No.	Sect./Page	Comment	Response	
	General			
1		Section 1 or Section 2 should introduce the point that this Remedial Site Evaluation (RSE) is driven by the Environmental Management Disposal Facility (EMDF) Record of Decision (ROD). This important context is not presented until Section 7.1. As described in the EMDF ROD, the effective implementation of this RSE is required to identify mercury methylation areas in Bear Creek which currently prevent the creek from being in compliance with applicable water quality standards.	Agree. Text in the first paragraph in Section 1.1 was revised as follows: "The objective of the Bear Creek Valley (BCV) mercury sources remedial site evaluation (RSE) is to evaluate potential sources of mercury and methylmercury within the BCV Watershed, as outlined in the mercury-management approach for Bear Creek in the Record of Decision for Comprehensive Environmental Response, Compensation, and Liability Act Oak Ridge Reservation Waste Disposal at the Environmental Management Disposal Facility, Oak Ridge, Tennessee (DOE/OR/01-2794&D2/R2; Environmental Management Disposal Facility [EMDF] Record of Decision [ROD]). The BCV Watershed is located in"	
2		Please clarify the objective. Section 1.1 states the objective "is to evaluate potential sources of mercury and methylmercury within the Bear Creek Valley (BCV) Watershed." However, the EMDF ROD states the RSE will "evaluate mercury methylation in Bear Creek "These objectives do not directly align as the first implies finding any mercury within the watershed while the second implies evaluating only those sources contributing to the water quality of Bear Creek. The proposed sampling aligns with the objective stated in the ROD (e.g., only sampling to a depth of 0.5 ft. when there is a potential for mercury at greater soil depths), not the watershed-wide objective stated in this SAP. The FFA parties should discuss and be in agreement on the objective of this effort.	Clarification. The Bear Creek Valley Watershed is the administrative name that is used to refer to projects within this watershed. As discussed in the mercury-management approach outlined in the EMDF ROD, the plan for reducing mercury loading and restoring Bear Creek to meet recreational use designations (as measured in fish tissue concentrations) may be a phased approach. The approach may recognize non-point source reductions to offset the point source discharge at EMDF, following treatment or other measures, to permanently reduce loading and potentially reduce the rate of mercury methylation. As stated in Section 3.2 (Data Quality Objective Step 2: Identify the Goals of the Study) and as agreed upon during the DQO meeting held on June 29, 2023, the purposes of this RSE are to:	
			Determine if there are areas (channel sediment and creek bank and floodplain soils) along Bear Creek and	



Comment No.	Sect./Page	Comment	Response
			its tributaries that are potential sources of mercury and methylmercury that may affect fish.
			Obtain data from various hydrologic settings (pools, beaver ponds, etc.) that may contribute to mercury methylation and its bioaccumulation in the environment of Bear Creek and a reference location (e.g., Hinds Creek).
			As stated in Section 3.1 (Data Quality Objective Step 1: State the Problem), there are insufficient data along Bear Creek and its tributaries to determine if there are potential sources of mercury and methylmercury in channel sediment and creek bank and floodplain soils that may be contributing to exceedances of fish tissue criterion in prior years. Based on this statement of the problem, this RSE SAP includes collection of sediment, creek bank soil, and floodplain soil data, as well as surface water data. In addition, as part of the RSE, additional parameters will be collected that may contribute to a better understanding of mercury methylation in Bear Creek. No change to the document proposed.
3		The sampling plan seems to ignore bank soils along the sides of the stream channel that are not within the top 0"-6". Sampling bank soils midway between the stream bottom and the floodplain top may shed some light on the potential for mercury/methylmercury to be migrating/leaching out of the soils and into the stream. Additionally, sampling of pore water within the shallow soils or midstream bank should be considered as part of a full effort to characterize mercury within the watershed.	Clarification. See responses to specific comments 4 and 6.
4		While there are sampling locations identified in upper Bear Creek that are both collocated and independent of beaver dams, that does not appear to be the case in lower Bear Creek. Consider adding some sampling locations in lower Bear Creek that are not colocated with beaver dams. This was discussed as part of the SAP Data Quality Objectives (DQO) meeting.	Clarification. It should be noted beaver activity has been increasing in the lower portions of Bear Creek. However, during field reconnaissance conducted in November 2023, the beaver dam located near transect locations BCT1 and BCT2 was determined to no longer be present.  As stated in Section 4.5 of the BCV Mercury Sources RSE SAP, actual field locations may be adjusted based on field conditions and sampling viability. Additional field reconnaissance in the vicinity of BCT1 and BCT2 was



Comment No.	Sect./Page	Comment	Response
			conducted on December 7, 2023, and it was determined that location BCT1 will be moved slightly further downstream since the beaver dam is gone. Figure 4.1 was revised to show the new location and is attached at the end of these comment responses.
			Note that transects BCT10 and BCT11 are also located in the lower reaches of Bear Creek (near the EMDF and the Bear Creek Burial Grounds, respectively) and are not associated with beaver dams.
	1	Specific	
1	Page 3, Section 2.2.1, last sentence	Revise the text to indicate when the BCV Phase I ROD addendum is projected to be submitted.	Clarification. The date associated with this change to the BCV Phase I ROD is anticipated to be sometime in FY 2024; however, it is suggested this detail not be added to the RSE SAP because this detail is not germane to implementing the field event.
			The last sentence of Section 2.2.1 was revised as follows: "Based on the EMDF ROD, the end use for Zone 1 and Zone 2 will be revised to restricted recreational and controlled industrial, respectively, which will be codified in an upcoming modification to the BCV Phase I ROD."
2	Page 7, second paragraph, last sentence	Data provided in Table 4.13 of the 2023 Remediation Effectiveness Report (RER) do not align with this statement. Even the averages often exceed the fish tissue criterion. Delete this sentence and instead provide the range of mercury levels in rock bass and redbreast sunfish sampled in Bear Creek.	Clarification. The last sentence in the second paragraph in Section 2.3 was revised as follows: "Long-term trends of monitored sunfish show that, despite significant fluctuation, mean mercury concentrations in fish (Figure 2.3) have been generally declining in the last 10 years and are below or just above the fish tissue criterion (0.3 µg/g) as of 2022.
			Note that a typographical error on Figure 2.3 was also fixed. Revised Figure 2.3 is attached at the end of these comment responses.
3	Page 12, Section 3.4, last bullet and Page 27,	As appropriate, revise the text to update the sampling and data evaluation/reporting schedules.	Agree. The third bullet in Section 3.4 was revised as follows: "Temporal – samples anticipated to be collected in December 2023 to meet the RSE milestone of September 2024."
	Section 7.2		In Section 7.2, the first sentence was revised as follows: "Fieldwork described in this BCV Mercury Sources RSE



Comment No.	Sect./Page	Comment	Response
			SAP is anticipated to be conducted in December 2023.  Data evaluation…"
	Page 15, Section 4.2, first bullet &sub-bullets	Revise the text to define upper and lower. Based on Figure 4.2, it appears these intervals are 0 to 3 inches and 3-6 inches (0 to 0.25 ft and 0.25 to 0.5 ft, respectively).	Clarification. As stated in the first bullet, the creek bank soils will be divided in half into upper and lower sections. This division of the upper and lower intervals will be defined based on the in-field height of the creek bank soil. Creek bank soil samples for each upper and lower interval will be collected from 0 to 0.5 ft within each interval.
			For additional clarification, text in the first bullet was revised as follows:
			"Creek bank soils will be divided in half into upper and lower sections based on in-field conditions as follows (Figure 4.2):"
			Text in the first sub-bullet was revised as follows:
			"For the upper section of the creek bank soils, samples will be collected by removing the surface soil on the upper half of each side of the bank. The upper"
			The text in the second sub-bullet was revised as follows:
			"For the lower section of the creek bank soils, samples will be collected by removing the surface soil on the lower half (above the creek level) of each side of the bank. The lower"
			Note that Figure 4.2 (transect diagram) was also replaced and is attached at the end of these comment responses.
5	Page 22, Section 5.3, second paragraph, last sentence	Revise the text as follows: "For surface water samples collected, analytical results will be compared with Ambient Water Quality Criteria (AWQC)." The text encourages confusion by implying AWQC are not required screening levels when they will almost certainly be Applicable or Relevant & Appropriate Requirements (ARARs) in the forthcoming BCV ROD.	Disagree. The final BCV ROD (FFA Appendix J date 2039) will address any AWQC exceedances in Bear Creek. The RSE SAP compares available surface water data to AWQC for comparison purposes only. No change to the document proposed.
6	Page A-6 [DQO Meeting Minutes], Transect	As stated in a follow-up note in the meeting minutes, TDEC recommends including hyporheic zone/shallow groundwater sampling in the RSE scope. There are two reasons for this recommendation.	Clarification. While hyporheic zone/shallow groundwater sampling in Bear Creek could be informative, the hyporheic push-point sampling that was conducted within the sediment bars in the East Fork Poplar Creek channel



Comment No.	Sect./Page	Comment	Response
	Sampling (Slide 38), Bullet 3	First, it is important to develop a more complete conceptual site model of mercury in Bear Creek to identify sources of mercury or methyl mercury. U.S. Department of Energy (DOE) research in a similar setting on the ORR finds that although the observed increase in methylmercury concentration and flux from upstream to downstream is related to instream methylation by periphyton and other biological activity (Watson, et al., 2016), this is not the only source of mercury and methylation in the stream. The researchers hypothesize methylmercury may be produced in hyporheic zones where anoxic, reducing geochemical environments may be conducive to methylation, resulting in dissolved methyl mercury concentrations in groundwater up to 10x greater than in surface water (Watson et al. 2016). More recent research supports that hypothesis, finding "additional sources of dissolved mercury inputs to East Fork Poplar Creek (EFPC) at baseflow during this study predominantly arise from the hyporheic zone," with up to 1/3 of downstream mercury discharging from hyporheic zone shallow groundwater and riparian inputs (Demers et al. 2018). The shallow/hyporheic zone groundwater "shows a strong, positive correlation between dissolved Mercury (Hg) and dissolved Methylmercury (MeHg)," whereas, historically, there has been a poor correlation between the two in surface water (Watson et al. 2016).  Second, the purpose of the mercury management strategy in the EMDF ROD that drives this RSE is to support the potential need to offset future mercury discharges from EMDF. Failure to identify and remove or treat sources of mercury or methyl mercury to offset EMDF discharges carries a risk the parties will be unable to develop the substantive equivalents to load allocations and waste load allocations, as envisioned in the EMDF ROD. This has the potential to impact TDEC's ability to approve mercury discharge limits for mercury that meet DOE's waste disposal and wastewater treatment needs while preventing discharges that do not "cause or contribute"	does not translate well to the shallow bedrock environment at Bear Creek where there are limited areas of gravely sediment accumulation. Much or most of the bed in Bear Creek is exposed bedrock that does not allow for "textbook" hyporheic exchange. Where some unconsolidated sediments accumulate, they are often large-enough grain size that there is likely little methylmercury production (not an anoxic environment and low water residence time). Lateral exchange with shallow saturated zones adjacent to the channel may occur at certain times of the year, but there is no indication of broader mercury contamination in those zones. In addition, the bank soils along Bear Creek are comprised primarily of tight clays that are not conducive to groundwater transmission.  Please note that, historically, ESD has conducted special studies in Bear Creek. One of the special studies looked into conducting porewater sampling in Bear Creek similar to the work completed in East Fork Poplar Creek. During that special studies investigation, it was impossible to conduct the sampling due to no (or a very thin veneer of) unconsolidated sediment, which did not allow for the advancement of push points, and ESD collected sediment samples as a contingency.  As discussed in the mercury-management approach in the EMDF ROD, additional evaluation may be required if non-point source mercury contributors are identified in the RSE. No change to the document proposed.



Comment No.	Sect./Page	Comment	Response
		Impacts, 20(4), p. 686-707; https://pubs.rsc.org/en/content/articlelanding/2018/em/c7em00538e	



### **Attachments**






Figure 2.3. Average concentrations of mercury in Bear Creek fish. (Revised based on TDEC specific comment 2)






Figure 4.1. BCV transect sampling locations. (Revised based on TDEC general comment 4 and EPA comment 3)






Figure 4.2. BCV RSE transect diagram. (Revised based on TDEC specific comment 4)

## APPENDIX B. PHOTOGRAPHS OF TRANSECT LOCATIONS



Figure B.1. Transect BCT1.



Figure B.3. Transect BCT2 – downstream.



Figure B.2. Transect BCT2 – upstream.



Figure B.4. Transect BCT2 – substrate.



Figure B.5. Transect BCT3 – downstream.



Figure B.7. Transect BCT4 – upstream.



Figure B.6. Transect BCT3 – upstream.



Figure B.8. Transect BCT4 – substrate.



Figure B.9. Transect BCT5 (downstream of dam) and transect BCT6 (upstream of dam); beaver dam was destroyed by a rain event.




Figure B.10. Transect BCT6.



Figure B.11. Transects BCT5 and BCT6 – substrate.



Figure B.12. Transect BCT7.




Figure B.13. Transect BCT8.



Figure B.14. Transect BCT9 – downstream.



Figure B.15. Transect BCT9 – upstream.



Figure B.16. Transect BCT10 – upstream.



Figure B.17. Transect BCT10 – downstream.



Figure B.18. Transect BCT10 – substrate.



Figure B.19. Transect BCT11 – downstream.



Figure B.20. Transect BCT11 – upstream.



Figure B.21. Transect BCT11 – substrate.



Figure B.22. Transect BCT12B – downstream.



Figure B.23. Transect BCT12B – upstream.



Figure B.24. Transect BCT12B – substrate.



Figure B.25. Transect BCT12A – downstream.



Figure B.26. Transect BCT12A – upstream.



Figure B.27. Transect BCT12A – substrate.



Figure B.28. Transect BCT13 – upstream.



Figure B.29. Transect BCT13 – downstream.



Figure B.30. Transect BCT14 – downstream.



Figure B.31. Transect BCT14 – upstream.



Figure B.32. Transect BCT15 – downstream.



Figure B.33. Transect BCT15 – upstream.



Figure B.34. Transect BCT15 – substrate.



Figure B.35. Transect HCTREF.

Figure B.36. Transect HCTREF – substrate.

## APPENDIX C. ANALYTICAL DATA SUMMARY TABLES BY MEDIA

Table C.1. BCV RSE unfiltered surface water data for the initial sampling event

Chemical	CAS	Units	BCT1-SW	BCT2-SW	BCT3-SW	BCT4-SW	BCT5-SW	BCT6-SW	BCT7-SW	BCT8-SW	BCT9-SW	BCT10-SW	BCT11-SW	BCT12B-SW	BCT12B-SW-D	BCT12A-SW	BCT13-SW	BCT14-SW	BCT14-SW-D	BCT15-SW	HCTREF-SW
									•		Physical	,	•	•			·		•		
Conductivity	N237	umho/cm	355.6	268	239	230	143	194	338	308	263	332	323	402		440	483	534		669	269
Dissolved oxygen	N328	ppm	10.56	11.88	11.76	10.61	9.87	10.54	11.75	11.14	9.9	10.66	10.06	9.79	 	10.16	11.01	12.11		13.36	10.94
Redox	NS215	mV	161	170.5	179	149	133	118	92	114	139	161	140	108	i 	140	149	154	i   	182.8	96
Temperature	N908	deg C	8.2	6	6.1	7.6	7.4	7.4	5.7	8.9	9.9	9.8	7.8	8.1	 	8.8	7.8	9.6	<u></u>	4.2	5.9
pН	N704	Std Unit	8.22	7.73	7.74	8.29	7.78	8.14	8.29	8.01	8.11	7.81	8.12	7.58		7.85	8	7.22		6.96	8.07
											Anions										
Chloride	16887006	mg/L	8.09	7.64	8.56	8.73	18	17.6	22.5	26.6	26.7	35.4	47.3	58.9	59.3	87.1	88.7	32.2	32.2	18.2	8
Fluoride	16984488	mg/L	0.105 J	0.121	0.131	0.13	0.117 J	0.116 J	0.15	0.16	0.156	0.164 J	0.207 J	0.232 J	0.233 J	0.255 J	0.318 J	0.402	0.395	0.467	0.101 J
Nitrate/Nitrite as nitrogen	N2788	mg/L	0.555	0.59	0.994	1.05	0.695	0.735	1.59	1.91	2	1.83	3.48	3.56	3.53	5.35	11.3	19.2	19.4	84.8	0.85
Sulfate	14808798	mg/L	30.7	29.2	30.7	31.6	9.6	9.62	16.6	19.3	18.8	15.9	20.6	20.8	20.9	21.7	26.3	33.4	33.4	31.5	8.04
											Metals										
Aluminum	7429905	mg/L	0.083	0.0985	0.159	0.112	0.228	0.978	0.517	0.181	0.177	0.189	0.142	0.317	0.32	0.22	0.198	0.0873	0.0838	0.206	0.106
Antimony	7440360	mg/L	0.001 U	0.001 U	0.001 U	0.001 U	0.001 U	0.001 U	0.001 U	0.001 U	0.001 U	0.001 U	0.001 U	0.001 U	0.001 U	0.001 U	0.001 U	0.001 U	0.001 U	0.001 U	0.001 U
Arsenic	7440382	mg/L	0.005 U	0.005 U	0.005 U	0.005 U	0.005 U	0.005 U	0.005 U	0.005 U	0.005 U	0.005 U	0.005 U	0.005 U	0.005 U	0.005 U	0.005 U	0.005 U	0.005 U	0.005 U	0.005 U
Barium	7440393	mg/L	0.0613	0.059	0.0585	0.0557	0.0521	0.0494	0.0707	0.0711	0.0674	0.0718	0.0823	0.0788	0.0882	0.104	0.117	0.116	0.119	0.209 J	0.057
Beryllium	7440417	mg/L	2.0E-04 U	2.0E-04 U	2.0E-04 U	2.0E-04 U	2.0E-04 U	2.0E-04 U	2.0E-04 U	2.0E-04 U	2.0E-04 U	2.0E-04 U	2.0E-04 U	2.0E-04 U	2.0E-04 U	2.0E-04 U	2.0E-04 U	2.0E-04 U	2.0E-04 U	2.0E-04 U	2.0E-04 U
Boron	7440428	mg/L	0.029	0.0253	0.0272	0.0247	0.0189	0.0186	0.0383	0.0372	0.0391	0.0391	0.0394	0.0211 J	0.0164 J	0.0201 J	0.0272	0.0533	0.0558	0.0653	0.00877 J
Cadmium	7440439	mg/L	3.0E-04 U	3.0E-04 U	3.0E-04 U	3.0E-04 U	3.0E-04 U	3.0E-04 U	3.0E-04 U	3.0E-04 U	3.0E-04 U	3.0E-04 U	3.0E-04 U	3.0E-04 U	3.0E-04 U	3.0E-04 U	3.0E-04 U	0.00104	0.00104	0.00836	3.0E-04 U
Calcium	7440702	mg/L	50	48.1	49.6	48.1	40.1	39.4	53.9	56.7	53.2	54.6	64	55.9	61.4	73.6	93.9	99	100	148	46.6
Chromium	7440473	mg/L	0.003 U	0.003 U	0.003 U	0.003 U	0.003 U	0.003 U	0.003 U	0.003 U	0.003 U	0.003 U	0.003 U	0.003 U	0.003 U	0.003 U	0.003 U	0.003 U	0.003 U	0.003 U	0.003 U
Cobalt	7440484	mg/L	3.0E-04 U	3.0E-04 U	3.0E-04 U	3.0E-04 U	3.0E-04 U	3.0E-04 U	4.3E-04 J	3.0E-04 U	3.0E-04 U	3.0E-04 U	3.0E-04 U	3.0E-04 U	3.0E-04 U	3.0E-04 U	3.0E-04 U	3.8E-04 J	3.5E-04 J	0.00235	3.0E-04 U
Copper	7440508	mg/L	3.3E-04 J	3.0E-04 U	3.1E-04 J	3.0E-04 U	6.5E-04 J	3.5E-04 J	5.5E-04 J	4.0E-04 J	3.5E-04 J	3.4E-04 J	5.9E-04 J	6.3E-04 J	6.8E-04 J	5.6E-04 J	5.3E-04 J	6.0E-04 J	0.00113 J	5.5E-04 J	3.3E-04 J
Iron	7439896	mg/L	0.154	0.164	0.236	0.197	0.219	0.196	0.556	0.229	0.224	0.174	0.216	0.411	0.42	0.322	0.239	0.223	0.22	0.204	0.17
Lead	7439921	mg/L	5.0E-04 U	5.0E-04 U	5.0E-04 U	5.0E-04 U	5.0E-04 U	5.0E-04 U	5.0E-04 U	5.0E-04 U	5.0E-04 U	5.0E-04 U	5.0E-04 U	5.0E-04 U	5.0E-04 U	5.0E-04 U	5.0E-04 U	5.0E-04 U	5.0E-04 U	5.0E-04 U	5.0E-04 U
Lithium	7439932	mg/L	0.00452 J	0.00534 J	0.00539 J	0.00491 J	0.00527 J	0.00551 J	0.00879 J	0.0102	0.0109	0.0126	0.00976 J	0.00554 J	0.00559 J	0.00738 J	0.00956 J	0.00941 J	0.0099 J	0.00888 J	0.003 U
Magnesium	7439954	mg/L	14.3	13.7	13.6	13.3	10.3	9.53	12.3	12.9	12.2	12.3	13	10.7	11.7	13.3	14.2	14.5	14.6	24.4	13.9
Manganese	7439965	mg/L	0.0216	0.0146	0.0328	0.0373	0.018	0.0171	0.161	0.034	0.0204	0.0095	0.0157	0.039	0.0413	0.0533	0.0768	0.278	0.28	2.1	0.0231
Mercury	7439976	mg/L	1.1E-06 =	9.9E-07 =	1.2E-06 =	1.3E-06 =	2.5E-06 =	2.2E-06 =	4.1E-06 J	2.7E-06 =	2.3E-06 =	4.5E-06 =	6.7E-06 =	1.3E-05 =	1.3E-05 =	1.3E-05 =	2.2E-05 =	6.9E-06 =	6.6E-06 =	1.4E-06 =	9.1E-07 =
Molybdenum	7439987	mg/L	4.0E-04 J	5.1E-04 J	5.5E-04 J	4.0E-04 J	2.0E-04 U	2.0E-04 U	2.6E-04 J	3.1E-04 J	2.9E-04 J	2.5E-04 J	3.0E-04 J	2.8E-04 J	3.0E-04 J	3.8E-04 J	3.5E-04 J	6.3E-04 J	5.1E-04 J	3.2E-04 J	2.0E-04 U
Nickel	7440020	mg/L	6.0E-04 U	6.0E-04 U	6.3E-04 J	6.0E-04 U	7.5E-04 J	6.0E-04 U	6.0E-04 U	6.1E-04 J	6.7E-04 J	6.0E-04 U	8.0E-04 J	9.8E-04 J	0.00101 J	0.00109 J	0.00139 J	0.00377	0.00362	0.0228	6.0E-04 U
Phosphorous	7723140	mg/L	0.018 U	0.018 U	0.018 U	0.018 U	0.018 U	0.018 U	0.018 U	0.018 U	0.018 U	0.018 U	0.018 U	0.018 U	0.037 J	0.018 U	0.018 U	0.018 U	0.018 U	0.018 U	0.018 U
Potassium	7440097	mg/L	1.77	1.55	1.59	1.49	1.22	1.13	1.55	1.56	1.43	1.52	1.77	1.75	1.89	2.05	2.3	2.53	2.56	3.1 J	1.82
Selenium	7782492	mg/L	0.006 U	0.00796 J	0.0117 J	0.0111 J	0.006 U	0.0101 J	0.006 U	0.0137 J	0.006 U	0.0105 J	0.0129 J	0.00752 J	0.0111 J	0.006 U	0.00741 J	0.006 U	0.00896 J	0.00982 J	0.00631 J
Silicon	7440213	mg/L	3.83	4.23	4.1	3.9	3.5	3.35	3.82	3.51	3.3	3.5	3.45	3.75	4.09	3.59	3.53	3.28	3.34	4.33	3.56
Silver	7440224	mg/L	3.0E-04 U	3.0E-04 U	3.0E-04 U	3.0E-04 U	3.0E-04 U	3.0E-04 U	3.0E-04 U	3.0E-04 U	3.0E-04 U	3.0E-04 U	3.0E-04 U	3.0E-04 U	3.0E-04 U	3.0E-04 U	3.0E-04 U	3.0E-04 U	3.0E-04 U	3.0E-04 U	3.0E-04 U
Sodium	7440235	mg/L	5.62	5.74	6.21	5.99	8.85	8.69	10.9	13	12	16.6	21.3	25.9 J	28.2 J	36.2 J	39.6	24	24.4	21.6	4.25
Strontium	7440246	mg/L	0.156	0.156	0.169	0.165	0.0669	0.0695	0.117	0.109	0.114	0.106	0.149	0.135	0.139	0.19	0.252	0.291	0.297	0.432	0.067
Thallium	7440280	mg/L	6.0E-04 U		6.0E-04 U		6.0E-04 U	6.0E-04 U	6.0E-04 U		6.0E-04 U	6.0E-04 U	6.0E-04 U	6.0E-04 U	6.0E-04 U	6.0E-04 U	6.0E-04 U	6.0E-04 U	6.0E-04 U	6.0E-04 U	6.0E-04 U
Uranium	7440611	mg/L	0.0112	0.0139	0.0153	0.0152	0.0168	0.0174	0.0334	0.0403	0.0408	0.0452	0.0378	0.0517	0.051	0.0727	0.089	0.134	0.137	0.0135	1.6E-04 J
Vanadium	7440622	mg/L	0.00101 J	0.001 U	0.001 U	0.001 U	0.00108 J	0.001 U	0.001 U	0.00139 J	0.0011 J	0.001 U	0.001 U	0.00122 J	0.00126 J	0.001 U	0.001 U	0.001 U	0.001 U	0.001 U	0.001 U
Zinc	7440666	mg/L	0.0033 U	0.0033 U	0.0052 J	0.00377 J	0.0033 U	0.0033 U	0.00519 J	0.0033 U	0.0033 U	0.0033 U	0.0033 U	0.00358 J	0.0033 U	0.0033 U	0.0033 U	0.0033 U	0.0033 U	0.00804 J	0.0033 U
			•	•		•	•	•	•		Other	•	•				•	•			
Methylmercury	22967926	mg/L	3.6E-08 J	5.7E-08 J	7.1E-08 J	7.9E-08 J	2.2E-08 UJ	2.2E-08 UJ	1.4E-07 =	7.6E-08 J	8.4E-08 =	5.1E-08 J	6.3E-08 J	9.8E-08 =	8.4E-08 =	6.4E-08 J	8.8E-08 =	1.7E-07 =	1.3E-07 =	6.6E-08 J	2.3E-08 J
										We	t chemistry										
Dissolved solids	N340	mg/L	196	200	209	186	136	150	221	226	230	247	284	287	286	373	485	428	413	676	167
Sulfide	18496258	mg/L	0.033 U	0.033 U	0.033 U	0.033 U	0.033 U	0.033 U	0.033 U	0.033 U	0.033 U	0.033 U	0.033 U	0.033 U	0.033 U	0.033 U	0.033 U	0.033 U	0.033 U	0.033 U	0.033 U
Suspended solids	N873	mg/L	1.01 J	0.57 U	4.89	0.7 J	1.47 J	3.33	15.9	2.28 U	2.28 U	3.75	2.45 J	2.3 J	3.46 J	0.57 U	2.86 J	3.1	4.6	15.4	8 J
Total organic carbon average	NS2302	mg/L	1.79	1.46	1.64	1.61	1.24	1.26	1.4 J	1.46	1.81	1.49	1.79	2.38	2.59	2.34	2.27	2.51	2.62	2.67	1.15
Turbidity	N1036	NTU	3	4	4	5	7	6	13	6	5	9	6	9		7	6	6		2	6
J = The analyte was positively ident	tified: the associ	iated numeric	cal value is the	e approximate c	concentration o	f the analyte ir	the sample.	•	•	•		•	•		•	•	•	•	•		

J = The analyte was positively identified; the associated numerical value is the approximate concentration of the analyte in the sample.

U = The analyte was analyzed for, but was not detected above the reported sample quantitation limit.

UJ = Analyte or compound was not detected above the reported detection limit, and the reported detection limit is approximated due to quality deficiency.

<sup>=</sup> Validated result, which is detected and unqualified.

<sup>-- =</sup> Analyte or compound not sampled for.

BCV = Bear Creek Valley CAS = Chemical Abstract Service RSE = remedial site evaluation

Table C.2. BCV RSE filtered surface water data for the initial sampling event

Chemical	CAS	Units	BCT1-SW	BCT2-SW	BCT3-SW	BCT4-SW	BCT5-SW	BCT6-SW	BCT7-SW	BCT8-SW	BCT9-SW	BCT10-SW	BCT11-SW	BCT12B-SW	BCT12B-SW-D	BCT12A-SW	BCT13-SW	BCT14-SW	BCT14-SW-D	BCT15-SW	HCTREF-SW
									1		Metals		•						'		•
Aluminum	7429905	mg/L	0.0193 U	0.0193 U	0.0193 U	0.0193 U	0.0202 J	0.0778	0.0193 U												
Antimony	7440360	mg/L	0.001 U	0.001 U	0.001 U	0.001 U	0.001 U	0.001 U	0.001 U												
Arsenic	7440382	mg/L	0.005 U	0.005 U	0.005 U	0.005 U	0.005 U	0.005 U	0.005 U												
Barium	7440393	mg/L	0.0612	0.0568	0.0559	0.055	0.0481	0.0481	0.0654	0.0672	0.0697	0.0697	0.0815	0.0833	0.0798	0.103	0.116	0.114	0.122	0.212	0.0552
Beryllium	7440417	mg/L	2.0E-04 U	2.0E-04 U	2.0E-04 U	2.0E-04 U	2.0E-04 U	2.0E-04 U	2.0E-04 U												
Boron	7440428	mg/L	0.0267	0.0256	0.0256	0.0261	0.0187	0.0177	0.036	0.039	0.0381	0.0422	0.0383	0.0177 J	0.0164 J	0.0198 J	0.0263	0.0548	0.052	0.066	0.00938 J
Cadmium	7440439		3.0E-04 U	3.0E-04 U	3.0E-04 U	8.6E-04 J	9.5E-04 J	0.0085	3.0E-04 U												
Calcium	7440702	mg/L	50.2	49.1	48.6	48.5	38.6	39.3	51.9	54.3	55.3	54.6	65.1	59.6	59.6	74.8	93.7	99.3	100	150	45.4
Chromium	7440473	mg/L	0.003 U	0.003 U	0.003 U	0.003 U	0.003 U	0.003 U	0.003 U												
Cobalt	7440484	mg/L	3.0E-04 U	3.0E-04 U	3.0E-04 U	3.0E-04 U	3.0E-04 U	0.00233	3.0E-04 U												
Copper	7440508		3.1E-04 J	3.0E-04 U	3.1E-04 J	3.0E-04 U	3.0E-04 U	3.5E-04 J	4.8E-04 J	4.1E-04 J	4.1E-04 J	3.6E-04 J	5.2E-04 J	5.2E-04 J	5.7E-04 J	3.0E-04 U					
Iron	7439896	mg/L	0.0462 J	0.0466 J	0.0443 J	0.041 J	0.033 U	0.033 U	0.0407 J	0.0335 J	0.033 U	0.033 U	0.0333 J	0.0599 J	0.0535 J	0.0566 J	0.0431 J	0.0719 J	0.033 U	0.0906 J	0.033 U
Lead	7439921	mg/L	5.0E-04 U	5.0E-04 U	5.0E-04 U	5.0E-04 U	5.0E-04 U	5.0E-04 U	5.0E-04 U												
Lithium	7439932	mg/L	0.00454 J	0.00502 J	0.0047 J	0.00478 J	0.00526 J	0.00509 J	0.00861 J	0.0102	0.00998 J	0.0132	0.00988 J	0.00541 J	0.00544 J	0.00713 J	0.00908 J	0.00948 J	0.01	0.00901 J	0.003 U
Magnesium	7439954	mg/L	14.3	13.4	13.3	13.2	9.71	9.9	11.4	12.1	12.2	12.3	13	11.4	11.3	13.4	14.2	14.6	14.8	23.6	13.6
Manganese	7439965	mg/L	0.0195	0.0131	0.0221	0.035	0.0143	0.0134	0.104	0.029	0.015	0.00578	0.00697	0.0317	0.032	0.0474	0.0696	0.252	0.25	2.04	0.0196
Mercury	7439976		5.7E-07 =	4.7E-07 J	5.2E-07 =	6.5E-07 =	8.0E-07 =	9.6E-07 =	1.1E-06 J	9.4E-07 =	8.1E-07 =	1.8E-06 =	1.9E-06 =	4.3E-06 =	3.9E-06 =	4.3E-06 =	3.8E-06 =	3.2E-06 =	3.2E-06 =	6.4E-07 =	3.9E-07 J
Molybdenum	7439987			4.7E-04 J	4.5E-04 J	3.7E-04 J	2.0E-04 U	2.0E-04 U	2.4E-04 J	3.4E-04 J	3.4E-04 J	2.3E-04 J	2.9E-04 J	2.9E-04 J	3.0E-04 J	3.6E-04 J	3.6E-04 J	5.0E-04 J	4.9E-04 J	3.6E-04 J	2.0E-04 U
Nickel	7440020	mg/L	6.0E-04 U	0.00158 J	6.0E-04 U	6.0E-04 U	6.0E-04 U	6.3E-04 J	6.9E-04 J	6.8E-04 J	9.2E-04 J	0.00126 J	0.0035	0.00316	0.0237	6.0E-04 U					
Phosphorous	7723140	mg/L	0.018 U	0.018 U	0.018 U	0.018 U	0.018 U	0.018 U	0.018 U												
Potassium	7440097	mg/L	1.73	1.57	1.48	1.47	1.11	1.1	1.45	1.51	1.47	1.49	1.77	1.81	1.85	1.98	2.25	2.48	2.56	3.15	1.69
Selenium	7782492	mg/L	0.006 U	0.006 U	0.0133 J	0.00643 J	0.006 U	0.00754 J	0.0141 J	0.0228 J	0.0108 J	0.006 U	0.006 U	0.006 U							
Silicon	7440213	mg/L	3.77	4.04	3.85	3.78	3.12	3.18	3.22	3.13	3.24	3.31	3.37	3.59	3.58	3.37	3.35	3.16	3.31	4.15	3.32
Silver	7440224	mg/L	3.0E-04 U	3.0E-04 U	3.0E-04 U	3.0E-04 U	3.0E-04 U	3.0E-04 U	3.0E-04 U												
Sodium	7440235		5.6	5.78	6.18	6.01	8.54	8.62	10.6	12.5	12.6	16.4	21.7	27.4 J	27.5 J	36.8 J	39.9	24.2	24.7	22.5	4.16
Strontium	7440246	mg/L		0.165	0.166	0.165	0.0702	0.0676	0.115	0.113	0.112	0.108	0.148	0.144	0.145	0.187	0.247	0.289	0.284	0.444	0.068
Thallium	7440280	mg/L	6.0E-04 U	6.0E-04 U			6.0E-04 U	6.0E-04 U		6.0E-04 U	6.0E-04 U	6.0E-04 U	6.0E-04 U	6.0E-04 U	6.0E-04 U	6.0E-04 U					
Uranium	7440611	mg/L	:	0.0124	0.0149	0.0149	0.0169	0.0165	0.0313	0.0413	0.0401	0.0465	0.0369	0.0553	0.0557	0.0724	0.0887	0.134	0.138 J	0.0136	1.7E-04 J
Vanadium	7440622		1	0.001 U	0.001 U	0.001 U	0.001 U	0.001 U	0.001 U	0.001 U											
Zinc	7440666	mg/L	0.0033 U	0.00845 J	0.0034 J	0.00347 J	0.0033 U	0.0033 U	0.00331 J	0.0033 U	0.0033 U	0.0033 U	0.0033 U	0.0147 J	0.0146 J	0.0033 U					
											Other										
Methylmercury	22967926	mg/L	4.4E-08 J	6.0E-08 J	4.8E-08 J	6.4E-08 J	2.3E-08 J	3.1E-08 J	9.0E-08 =	9.0E-08 =	7.3E-08 J	6.4E-08 J	4.3E-08 J	6.3E-08 J	7.5E-08 J	7.2E-08 J	7.4E-08 J	7.4E-08 J	7.6E-08 J	5.0E-08 J	2.5E-08 J
											Wet chemi	stry									
Dissolved organic carbon	NS248	mg/L	1.85	1.48	1.57	1.64	1.34	1.3	1.5	1.61	1.57	1.56	1.86	2.21	2.76	2.36	2.04	2.7	2.57	2.62	1.42
I - The englyte was positively								_													

J = The analyte was positively identified; the associated numerical value is the approximate concentration of the analyte in the sample.

BCV = Bear Creek Valley CAS = Chemical Abstract Service RSE = remedial site evaluation

U = The analyte was analyzed for, but was not detected above the reported sample quantitation limit.

<sup>=</sup> Validated result, which is detected and unqualified.

<sup>-- =</sup> Analyte or compound not sampled for.

Table C.3. BCV RSE unfiltered surface water data from April 2024

Chemical	CAS	Units	BCT1-SW	BCT2-SW	BCT3-SW	BCT4-SW	BCT5-SW	BCT6-SW	BCT7-SW	BCT8-SW	BCT9-SW	BCT10-SW	BCT11-SW	BCT12B-SW	BCT12B-SW-D	BCT12A-SW	BCT13-SW	BCT14-SW	BCT14-SW-D	BCT15-SW	HCTREF-SW
					•			•	•		Phys	sical		•			•	•		•	
Conductivity	N237	umho/cm	298	240	278	232	240	239	283	292	278	295	333	349		393	460	684		556	289
Dissolved oxygen	N328	ppm	10.96	8.18	8.1	8.48	7.6	8.65	6.59	9.83	7.97	9.59	9.41	9.36		9.49	9.23	8.8		9.11	9.5
Redox	NS215	mV	-7.9	-1.7	14.5	13.2	19.9	28.6	24.1	15.4	229.5	219	222.6	211.1		199.1	206.5	235.3		230.6	244.1
Temperature	N908	deg C	19	19.2	16.6	16.8	16.4	16.4	18.1	16.8	16.7	15.4	15.9	15.6		15.7	15.9	16.2		17.3	18.2
pН	N704	Std Unit	8.02	7.95	7.66	7.84	7.71	7.69	7.64	7.74	7.56	7.82	7.96	7.91		7.96	7.82	7.46		7.14	8.1
											Met	tals									
Mercury	7439976	mg/L	1.6E-06 =	1.8E-06 =	1.8E-06 =	1.9E-06 =	1.5E-06 =	1.6E-06 =	4.2E-06 =	2.1E-06 =	2.1E-06 =	2.6E-06 =	4.5E-06 =	7.9E-06 =	7.7E-06 =	4.2E-06 =	9.8E-06 =	8.4E-06 =	8.1E-06 =	5.1E-06 =	1.0E-06 =
											Otl	ier									
Methylmercury	22967926	mg/L	1.1E-07 J	9.6E-08 J	9.8E-08 J	8.3E-08 J	8.1E-08 J	7.8E-08 J	1.7E-07 J	1.3E-07 J	8.9E-08 J	5.1E-08 J	4.7E-08 J	1.6E-07 J	1.6E-07 J	1.2E-07 J	2.7E-07 J	8.5E-08 J	7.8E-08 J	6.6E-08 J	4.2E-08 J
		•		•			•	•		•	Wet che	emistry	•								
Turbidity	N1036	NTU	5	5	1	6	6	4	9	6	6	5	6	7		4	7	5		9	5

J = The analyte was positively identified; the associated numerical value is the approximate concentration of the analyte in the sample.

BCV = Bear Creek Valley CAS = Chemical Abstract Service RSE = remedial site evaluation

 $<sup>=</sup> Validated \ result, \ which \ is \ detected \ and \ unqualified.$ 

<sup>-- =</sup> Analyte or compound not sampled for.

Table C.4. BCV RSE filtered surface water data from April 2024

CAS	Units	BCT1-SW	BCT2-SW	BCT3-SW	BCT4-SW	RCT5-SW	BCT6-SW	BCT7-SW	RCT8-SW	RCT9-SW	RCT10-SW	RCT11-SW	BCT12R-SW	RCT12R-SW-D	RCT12A-SW	BCT13-SW	BCT14-SW	RCT14-SW-D	RCT15-SW	HCTREF-SW
CITO	Cints	Dellow	BC12 SW	Bert St.	Berrow	Berssy	Berest	Berrow	Belosii			Bellion	DCT12B SW	Delliz Sw B	Belling	Derie Sii	Bellion	Dell's w D	De 113 5 W	I HE THE I S W
7439976	mg/L	5.4E-07 =	4.4E-07 J	4.7E-07 J	5.1E-07 =	4.4E-07 J	4.2E-07 J	8.7E-07 =	7.0E-07 =	7.4E-07 =	8.7E-07 =	1.4E-06 =	1.8E-06 =	1.8E-06 =	1.2E-06 =	1.8E-06 =	1.8E-06 =	1.9E-06 =	1.0E-06 =	3.3E-07 J
										0	ther		-			•			•	
22967926	mg/L	8.3E-08 J	7.7E-08 J	5.2E-08 J	5.4E-08 J	3.2E-08 J	5.9E-08 J	8.4E-08 J	8.5E-08 J	3.7E-08 J	2.2E-08 UJ	3.4E-08 J	1.0E-07 J	1.2E-07 J	2.2E-08 UJ	1.5E-07 J	5.8E-08 J	4.2E-08 J	6.9E-08 J	5.3E-08 J
	7439976	7439976 mg/L	7439976   mg/L   5.4E-07 =	7439976   mg/L   5.4E-07 =   4.4E-07 J	7439976   mg/L   5.4E-07 =   4.4E-07 J   4.7E-07 J	7439976   mg/L   5.4E-07 =   4.4E-07 J   4.7E-07 J   5.1E-07 =	7439976   mg/L   5.4E-07 =   4.4E-07 J   4.7E-07 J   5.1E-07 =   4.4E-07 J	7439976   mg/L   5.4E-07 =   4.4E-07 J   4.7E-07 J   5.1E-07 =   4.4E-07 J   4.2E-07 J	7439976   mg/L   5.4E-07 =   4.4E-07 J   4.7E-07 J   5.1E-07 =   4.4E-07 J   4.2E-07 J   8.7E-07 =	7439976   mg/L   5.4E-07 =   4.4E-07 J   4.7E-07 J   5.1E-07 =   4.4E-07 J   4.2E-07 J   8.7E-07 =   7.0E-07 =	M   7439976   mg/L   5.4E-07 =   4.4E-07 J   4.7E-07 J   5.1E-07 =   4.4E-07 J   4.2E-07 J   8.7E-07 =   7.0E-07 =   7.4E-07 =   0.0E-07 =	T439976   mg/L   5.4E-07 =   4.4E-07 J   4.7E-07 J   5.1E-07 =   4.4E-07 J   4.2E-07 J   8.7E-07 =   7.0E-07 =   7.4E-07 =   8.7E-07 =   Other	7439976   mg/L   5.4E-07 =   4.4E-07 J   4.7E-07 J   5.1E-07 =   4.4E-07 J   4.2E-07 J   8.7E-07 =   7.0E-07 =   7.4E-07 =   8.7E-07 =   1.4E-06 =     Other	Metals   7439976   mg/L   5.4E-07 =   4.4E-07 J   4.7E-07 J   5.1E-07 =   4.4E-07 J   4.2E-07 J   8.7E-07 =   7.0E-07 =   7.4E-07 =   8.7E-07 =   1.4E-06 =   1.8E-06 =     Other	7439976   mg/L   5.4E-07 =   4.4E-07 J   4.7E-07 J   5.1E-07 =   4.4E-07 J   4.2E-07 J   8.7E-07 =   7.0E-07 =   7.4E-07 =   8.7E-07 =   1.4E-06 =   1.8E-06 =     Other	7439976   mg/L   5.4E-07 =   4.4E-07 J   4.7E-07 J   5.1E-07 =   4.4E-07 J   4.2E-07 J   8.7E-07 =   7.0E-07 =   7.4E-07 =   8.7E-07 =   1.4E-06 =   1.8E-06 =   1.8E-06 =   1.2E-06 =	7439976   mg/L   5.4E-07   4.4E-07 J   4.7E-07 J   5.1E-07   4.4E-07 J   4.2E-07 J   8.7E-07   7.0E-07   7.4E-07   8.7E-07   1.4E-06   1.8E-06   1.8E-06   1.2E-06   1.8E-06   1.8E-06	T439976   mg/L   5.4E-07   4.4E-07 J   4.7E-07 J   5.1E-07   4.4E-07 J   4.2E-07 J   8.7E-07   7.0E-07   7.4E-07   8.7E-07   1.4E-06   1.8E-06   1.8E-06   1.8E-06   1.2E-06   1.8E-06   1.8E-06	Metals   1/439976   mg/L   5.4E-07   4.4E-07 J   4.7E-07 J   5.1E-07   4.4E-07 J   4.2E-07 J   4.2E-07 J   8.7E-07   7.0E-07   7.4E-07   8.7E-07   1.4E-06   1.8E-06   1.8E-06   1.2E-06   1.8E-06   1.8E-06	Telephone   Tele

J = The analyte was positively identified; the associated numerical value is the approximate concentration of the analyte in the sample.

UJ = Analyte or compound was not detected above the reported detection limit, and the reported detection limit is approximated due to quality deficiency.

<sup>=</sup> Validated result, which is detected and unqualified.

BCV = Bear Creek Valley CAS = Chemical Abstract Service RSE = remedial site evaluation

Table C.5. BCV RSE floodplain soil data

Chemical	CAS Units	BCT1-FP	BCT2-FP	BCT3-FP	BCT4-FP	BCT5-FP	BCT6-FP	BCT7-FP	BCT8-FP	BCT9-FP	BCT10-FP	BCT11-FP	BCT12B-FP	BCT12B-FP-D	BCT12A-FP	BCT13-FP	BCT14-FP	BCT14-FP-D	BCT15-FP	HCTREF-FP
all "I	1,007000 //	2.41.7		1 410	1 22		1 246		nions	7.02.1			2.67	1 2.42	1.72	2.06	1 210		1 0.04	2.65
Chloride		2.41 J	2.1 J	4.18	3.2	5.47	2.46	6.31 J	6.85 J	7.93 J	2.53 J	5.82	2.67	2.43	4.73	3.06	3.19	5.48	8.94	2.65
Fluoride	16984488 mg/kg		1.31	2.08	1.05 J	1.7 J	2.23 J	2.32	2.04	3.29	1.48	3.68 J	6.54	4.18	4.37	13.1	2.36	2.44	5.75	1.08 J
Nitrate	14797558 mg/kg		1.76	0.938 J	2.09	4.47	1.13 J	2.79	2.59	3.99	1.56	2.69	3.78	3.04	2.57	1.5	2.72	2.82	15.8	0.841 J
Nitrite	14797650 mg/kg		0.39 U	0.402 U	0.434 U	0.832 J	0.395 U	0.479 U	0.488 U	0.441 U	0.423 U	0.421 U	0.422 U	0.389 U	0.429 U	0.446 U	0.448 U	0.446 U	0.445 U	0.41 U
Sulfate	14808798   mg/kg	4.09 J	6.99	6.13	6.12	7.66	4.42 J	10.7	14.2 etals	10.6	3.08 J	3.85 J	3.65 J	3.08 J	<u> 6</u>	21.4 J	5.67	6.64	16.1	4.7 J
Aluminum	7429905 mg/kg	5180	5000	5840	4920	5250	6920	5280	7470 J	8010 J	4950	5600	7430	6910	12,500	7000	8720	8620	10,200	5580
Antimony	<del>-</del>	0.412 U	0.375 U	0.399 U	0.372 U	0.425 U	0.387 U	0.47 U	0.487 U	0.452 U	0.375 U	0.389 U	1 J	0.691 J	1.19 J	0.835 J	0.453 U	1.16 J	0.421 U	0.407 U
Arsenic	7440382 mg/kg	2.51	3	3.36	3.48	4.79	4.27	3.27	4.22	8.89	4.22	4.27	5.75	4.85	7.3	5.37	7.14	7.93	4.35	3.56
Barium	7440393 mg/kg	79.6	104	60.7	83.5	87.3	74.8	93.3	106	75	78.8	89.9	84.8	78.5	177	161	88.3	85	283	74.4
Beryllium	7440417 mg/kg		0.56 J	0.414 J	0.584	0.499 J	0.481 J	0.649 J	0.894	0.783	0.252 J	0.624	0.251 J	0.205 J	0.445 J	0.379 J	0.137 U	0.129 U	1.01	0.45 J
Boron	7440428 mg/kg		1.14 U	6.06	5.52 J	1.29 U	1.17 U	1.86 J	9.74	8.59	1.68 J	3.42 J	2.49 J	1.1 U	3.81 J	1.37 U	1.6 J	1.75 J	4.35 J	1.23 U
Cadmium	7440428 mg/kg		0.464	0.238	0.479	0.926	0.139 J	1.31	1.56	1.6	2.65	1.04	6.04	7.86	6.95	6.93	0.197 J	0.201 J	10.3	0.0799 J
Calcium		1780	2030	1260	2300	3230	1480	5630	8190 E	2850 E	1770	1870	2330	1690	2240	27,300	3200	3000	5680	1890
Chromium		10.7	12.6	11.2	10.7	18	18	13.9	23.8	19.3	13.6	13.2	14.9	14.3	20.1	19.1	23.8	19.3	23.2	15.5
Cobalt	7440473 mg/kg 7440484 mg/kg	9.1	10	6.48	9.58	10.6	12	10.2	12.4	13.3	11.1	13.5	12.6	12	18.3	17.4	9.31	8.51	25.5	10.9
	7440508 mg/kg		8.3	6.34	7.27	9.96	9.03	8.65	11.8	16	9.05	12.2	28.7	19	29.3	19.2	15.1	14.5	14	9.87
Copper Iron	7439896 mg/kg		12,600	11,500	12,000	13,600	18,800	12,500	19,400 J	18,800 J	11,600	14,200	13,400	14,300	19,900	20,500	20,000	17,900	22,400	14,200
Lead	7439921 mg/kg		12.9	9.62	10.4	15 J	15.6 J	11.9	13.5 J	16.4 J	17.4	22.6	32.8	31.3	43.6	21.6	21.9	21	16.4	15.3 J
Lithium	7439932   mg/kg	12.2	12.6	12.7	12	13.7	10.1	15.8	17.9	19.4	19.2	12.2	23.1	23.4	24.9	24.7	11.5	13	16.9	11.9
Magnesium	7439954 mg/kg	1180	1380	1240	1210	1520	1010	2540	3670	1340	795	863	971	913	1260	2040	861	843	5370	1270
Manganese	7439965 mg/kg		975	448	836	905	1120	922	994	682	1140	1170	964	900	1230	2460	1150	1070	5180	728
Mercury	7439976 mg/kg		0.11 =	0.086 =	0.18 =	0.16 =	0.062 =	0.22 =	0.27 =	0.38 =	0.64 =	0.81 J	2.8 =	3 =	3.5 =	3.3 =	0.74 =	0.72 =	0.47 =	0.04 =
Molybdenum	7439987 mg/kg		0.295 J	0.348 J	0.249 J	0.257 U	0.234 U	0.285 U	0.295 U	0.274 U	0.578 J	0.672 J	0.982 J	0.901 J	0.971 J	0.87 J	0.496 J	0.441 J	0.29 J	0.246 U
Nickel	7440020   mg/kg	13.9	13.3	9.77 J	13.9 J	12.3	9.17	13.2	17 E	17.1 E	18.3 J	14	31.5 J	18.9 J	42.1 J	34.5 J	14.8	14.4	75.7	10.7
Phosphorous	7723140 mg/kg	162	206	162	200	207	158	228	250	227	213	223	244	248	291	244	274	296	270	220
Potassium	7440097 mg/kg	641	661	711	617	616	640	564	821	728	394	550 J	676	521	848	572	583	599	1300	699
Selenium	7782492 mg/kg	0.86 J	0.952 J	0.798 J	1.02 J	1.49	1.27	0.941 J	0.874 J	1.02 J	1.36	1.48	1.47	1.48	2.02	1.67	1.23 J	1.24 J	2.34	1.21
Silicon	7440213 mg/kg	2550	2470	2080	1880	557	552	1460	1780	1940	1800	2970	2690	2590	3290	2670	2550	2460	1040	496
Silver	7440224   mg/kg	0.135 J	0.168 J	0.121 U	0.113 U	0.643 U	0.586 U	0.142 U	0.148 U	0.685 U	0.114 U	0.118 U	0.235 J	0.11 U	0.121 U	0.298 J	0.137 U	0.129 U	1.19	0.616 U
Sodium	7440235 mg/kg	8.73 U	7.96 U	8.46 U	7.89 U	9.43 J	12.5 J	25.2 J	34.2 J	30.8 J	7.96 U	16 J	8.35 U	7.73 U	10.3 J	15.9 J	11.5 J	11.6 J	78.6	9.58 J
Strontium	7440246 mg/kg	4.33	5.77	4.6	7.21	6.42	5.47	8.84	9.74	6.36	4.95	6.79	8.27	8.92	9.44	15	10.9	11.7	17.1	5.53
Thallium	7440280 mg/kg	0.159 U	0.167 U	0.152 U	0.172 U	0.181 U	0.176 U	0.195 U	0.204 U	0.19 J	0.157 U	0.179 U	0.163 U	0.173 U	0.199 J	0.167 U	0.199 J	0.201 J	0.186 U	0.155 U
Uranium	7440611 mg/kg	11.6	7.17	4.59	10.1	16.1	3.3	16.9	16.8	30.1	41.7	19.9	63.6	78	106	16.5	4.61	5.48	7.08	0.35
Vanadium	7440622 mg/kg	13	13.9	14.9	13.5	18.4	23.6	13.3	21.8	29.9	15.6	18.2	19.8	25.3	31.5	22.8	38.4	32.1	16.4	18
Zinc	7440666 mg/kg	28.1	29.9	20.6	29.1	36.1	23.6	34.9 J	46.3	56.8	32.2	31.2 J	64.7	51.2	74.9	68.9	53.6	52.1	64.7	35.5
								0	ther											
Methylmercury	22967926 mg/kg	2.8E-04 =	4.9E-04 =	1.1E-04 =	3.7E-04 =	2.2E-04 =	1.3E-04 =	6.0E-04 =		4.0E-04 =	1.6E-04 J	1.7E-04 =	0.001 =	0.0017 =	7.0E-04 =	5.1E-04 J	4.8E-04 =	4.2E-04 =	5.7E-04 =	1.0E-04 =
Mercury (F0) - volatile elemental mercury	NS1021 mg/kg		<del> </del>	1	<u> </u>	1.7E-04 UJ	0.0014 J	Sequentia 2.0E-04 UJ	extraction	5.6E-04 J	ŀ	1.7E-04 UJ	1.8E-04 UJ	1	<u> </u>	<u> </u>	1.8E-04 UJ		1	2.2E-04 J
Mercury (F1) - volatile elemental mercury  Mercury (F1) - water soluble mercury	NS1021 mg/kg		 	<u> </u>	<del> </del>	0.0032 J	0.0014 J 0.0025 J	0.0034 J	0.0039 J	0.013 =	  	0.025 =	0.054 J		 ! 		0.02 J		<u> </u>	0.0011 UJ
Mercury (F2) - pH 2 soluble mercury	NS1022 mg/kg			<del> </del>	ļ	0.0032 J	0.0023 J	0.0016 J	0.0037 J	0.0033 J		0.023 =	0.034 J				0.002 J	<del> </del>	<u></u>	0.0011 UJ
	NS1023 mg/kg				ļ <u>-</u>	0.002 3	0.0012 03	0.00163	0.00193	0.00333	i	0.55 =	0.04 J 0.85 J				0.0097 J	ļ	<del> </del>	0.011 UJ
Mercury (F3) - 1N potassium hydroxide extractable mercury  Mercury (F4) - 12N nitric acid soluble mercury	NS1024 Ilig/kg		 		<del>-</del>	0.0062 J	0.026 – 0.0059 UJ	0.11 – 0.015 J	0.099 – 0.018 J	0.10 -		0.33 =	0.83 J 0.56 J				0.27 J	<del>-</del>		0.0055 UJ
Mercury (F4) - 1218 miric acid soluble mercury  Mercury (F5) - aqua regia soluble mercury	NS1023 mg/kg			<del> </del>	<del> </del>	0.0035 J	0.0039 UJ	0.013 J	0.018 J	0.031 =		0.045 =	0.053 J		<del> </del>	<del> </del>	0.037 J	<del> </del>	<del> </del>	÷
Mercury (F6) - mineral-bound mercury				<del> </del>	ļ <del>-</del>	7.6E-04 J	4.2E-04 J	4.7E-04 J	4.0E-04 J	0.011 – 0.001 J		7.3E-04 J	0.0026 J				0.0047 J		<u> </u>	0.0012 J 0.01 J
Mercury (F6) - total mercury by summation	NS1027 mg/kg NS1028 mg/kg			<del> </del>	<del> </del>	0.099 =	0.031 =	0.13 =	0.13 =	0.001 3		0.75 =	1.6 J		<del> </del>		0.0017J		<u>                                     </u>	0.01 J 0.029 J
vicion y (1 5) - total incicuty by Summation	i 1101020   Hig/Kg		<u></u>	i	i	i 0.033 -	i 0.031 -	•	hemistry	i 0.22 -	i	i 0.75 -	i 1.0 J	i	<u></u>	<u> </u>	i 0.343	<u></u>	<u> </u>	0.029 J
Moisture	N544 %	22	19.9	20.1	25.2	23.9	21.8	36.4	36.1	28.1	24	22.7	23.5	20	25.3	27.3	28.6	28.7	33.3	20.9
Sulfide	18496258 mg/kg		21.6 J	30.3	11.6 U	22.9 J	40	15.4 J	59.1	40.9	30.7 J	30.4 J	23.4 J	18.6 J	32.6 J	22.2 J	13.3 J	15.2 J	19.2 J	24.4 J
Total organic carbon average		14,600	17,000	9760	19,400	25,200	8810	25,800	24,200	18,900	20,200	15,800	24,100	31,900	33,600	25,500	39,200	38,800	26,600	17,400
E = Estimated, matrix interference.		¥	. , , , , , , ,		. , ,	. /	:	. ,	. ,		. ,	. /	. ,	. 27.7.7	. ,		: /	. /	. , , , , , ,	

J = The analyte was positively identified; the associated numerical value is the approximate concentration of the analyte in the sample.

U = The analyte was analyzed for, but was not detected above the reported sample quantitation limit.

UJ = Analyte or compound was not detected above the reported detection limit, and the reported detection limit is approximated due to quality deficiency.

<sup>=</sup> Validated result, which is detected and unqualified.

<sup>-- =</sup> Analyte or compound not sampled for.

BCV = Bear Creek Valley CAS = Chemical Abstract Service RSE = remedial site evaluation

Table C.6. BCV RSE creek bank soil data

Chemical	CAS	Units	BCT1-BSL	BCT1-BSU	BCT2-BSL	BCT2-BSU	BCT3-BSL	BCT3-BSU	BCT4-BSL	BCT4-BSU	BCT5-BS	BCT5-BSL	BCT5-BSU	BCT6-BS	BCT6-BSL	BCT6-BSU	BCT7-BS	BCT8-BS	BCT9-BS	BCT10-BSL	BCT10-BSU	BCT11-BS	BCT11-BSL	BCT11-BSU
	•					•	•	•	•		Anions	•	•	•	•		•	•		•	•	•		
Chloride	16887006	mg/kg	2.29 J	2.32 J	2.9	2.65	7.5	3.5	3.91	2.65		13	2.42		4.83	2.96	14.1 J	9.9 J	11.8 J	3.3	1.95 J		13.8	9.28
Fluoride	16984488	mg/kg	1.89	0.858 J	1.91	2.64	5.34	4.11	1.99	2.17		3.37 J	2.66 J		2.18 J	2.25 J	3.02	1.82	2.68	3.65	2.64		4.1 J	3.83 J
Nitrate	14797558	mg/kg	2.11	1.96	1.09 J	1.43	0.41 U	1.09 J	1.81	1.42		0.851 J	0.969 J		0.856 J	1.04 J	1.38	0.889 J	4.4	1.28	0.403 U		2.18	2.03
Nitrite	14797650	mg/kg	0.408 U	0.423 U	0.405 U	0.404 U	0.41 U	0.4 U	0.45 U	0.404 U		0.411 U	0.391 U		0.425 U	0.394 U	0.449 U	0.442 U	0.429 U	0.417 U	0.403 U		0.408 U	0.402 U
Sulfate	14808798	mg/kg	6.88	3.98 J	20.8	6.99	16.1	4.05 J	11.6	7.66		6.28	2.8 J	ļ	10.1	3.46 J	11.9	9.28	9.36	6.35 J	2.48 J		7.87	4.29 J
										]	Metals													
Aluminum	7429905	mg/kg	4370	5430	7570	6560	5900	7330	4930	5280		5670	5100		7030	6340	5160	7320 J	7900 J	5790	3530		5520	5410
Antimony	7440360	mg/kg	0.432 U	0.421 U	0.418 U	0.362 U	0.401 U	0.655 J	0.41 U	0.405 U		0.942 J	0.356 U		0.367 U	0.351 U	0.47 U	0.419 U	0.435 U	0.376 U	0.672 J		0.406 J	0.365 U
Arsenic	7440382	mg/kg	3.3	2.58	7.59	3.12	2.95	3.13	3.6	3.5		3.04	4.58		3.91	4.79	2.79	5.99	5.39	4.05	2.59		3.41	3.15
Barium	7440393	mg/kg	62.7	89.5	92.5	116	68	62.1	92.1	77		75.6	58.3		138	103	87.7	68.6	95.1	79.8	42.1		90.2	86.7
Beryllium	7440417	mg/kg	0.428 J	0.586 J	0.717	0.609	0.463 J	0.446 J	0.55 J	0.56 J		0.431 J	0.398 J		0.721	0.563	0.534 J	1.01	1.04	0.145 J	0.104 U		0.659	0.601
Boron	7440428	mg/kg	3.49 J	1.28 U	1.27 U	1.1 U	4.13 J	4.26 J	5.23 J	4.11 J		1.24 U	1.08 U		1.11 U	1.07 U	1.42 U	13.8	8.14	3.01 J	1.04 U		3.56 J	3.12 J
Cadmium	7440439	mg/kg	1.11	0.623	0.129 J	0.278	0.0384 J	0.0484 J	0.301	0.247		0.094 J	0.575		0.0801 J	0.092 J	0.621	1.53	2.24	0.872	0.106 J		0.29	0.378
Calcium	7440702	mg/kg	2760	1680	1540	1820	1260	4440	3370	1520		1360	1550		1670	1470	3440	3530 E	2790 E	3060	633		1870	1890
Chromium	7440473	mg/kg	9.36	11.7	20.2	11.2	9.76	10.3	10.4	11.6		14	14.3	]	24.4	18.1	10.9	34.8	22.4	15.3	8.49		16.7	12.4
Cobalt	7440484	mg/kg	6.74	9.58	14.1	12.8	8.28	8.5	9.13	9.72		11.7	8.8	]	12.5	15.5	9.24	13.4	15.1	11.3	6.62		14.6	13.9
Copper	7440508	mg/kg	8.21	8.04	8.06	8.52	5.63	8.19	11.8	6.9		8.12	5.96	]	7.69	8.84	6.17	9.33	11.2	8.91	6.84		7.2	8.44
Iron	7439896	mg/kg	9380	12,500	20,500	14,600	11,200	12,000	11,200	12,000		12,600	14,000		21,500	16,400	10,400	25,400 J	18,400 J	13,900	8010		14,700	13,400
Lead	7439921	mg/kg	10.2	12.3	14.8	16.6	10.4	12.5	10	11.8		14.4 J	15.8 J		15.1 J	17.1 J	12.8	15.7 J	17.9 J	14.5	10.1		15.5	15
Lithium	7439932	mg/kg	14.5	12.2	9.4	11.1	10.2	14	10.3	9.98		9.78	12.8		12.8	11.5	12.3	16.6	18.7	13.4	7.71		10.3	9.48
Magnesium	7439954	mg/kg	1030	1240	1520	1420	989	1500	1560	1070		875	692		1250	1030	1540	1920	1400	1610	459		1070	1060
Manganese	7439965	mg/kg	635	842	767	1360	547	505	745	688		886	567		1760	1350	1210	583	951	1030	401	<u> </u>	1380	1250
Mercury	7439976	mg/kg	0.18 =	0.15 =	0.041 =	0.12 =	0.024 J	0.032 =	0.066 =	0.1 =		0.042 =	0.2 =	ļ	0.033 =	0.034 =	0.2 =	0.18 =	0.55 =	0.15 =	0.075 =	ļ	0.12 J	0.23 J
Molybdenum	7439987	mg/kg	0.355 J	0.345 J	0.495 J	0.421 J	0.445 J	0.312 J	0.257 J	0.245 U		0.248 U	0.216 U	<u> </u>	0.222 U	0.213 U	0.327 J	0.254 U	0.264 U	0.418 J	0.315 J		0.527 J	0.555 J
Nickel	7440020	mg/kg	11.7	13.8	10.8	15.8	8.33 J	10.1 J	12.3 J	12.3 J		8.44	9.01		11.8	10.3	8.23	17.5 E	19.7 E	12.5 J	5.39 J		9.63	9.1
Phosphorous	7723140	mg/kg	176	179	159	136	144	158	181	154		144	146		133	145	189	241	242	187	114		174	174
Potassium	7440097	mg/kg	642	718	773	661	576	797	598	606	<u></u>	562	471		595	663	420	826	841	451	246	<u> </u>	624 J	612 J
Selenium	+	mg/kg	0.873 J	0.832 J	1.08 J	0.801 J	1.04 J	0.924 J	0.971 J	0.949 J		1.33	1.14 J	ļ	1.84	1.34	1.04 J	0.883 J	1.04 J	1.2	0.842 J	ļ	1.34	1.22 J
Silicon	+	mg/kg	1910	2530	3050	2870	2050	2040	2080	2100		517	443	<u> </u>	460	508	1490	1850	1960	1820	1240	ļ	2640	2450
Silver	7440224	mg/kg	0.131 U	0.128 J	0.267 J	0.326 J	0.121 U	0.124 J	0.124 U	0.123 U		0.621 U	0.54 U	<u> </u>	0.556 U	0.533 U	0.194 J	0.635 U	0.906 J	0.114 U	0.104 U	ļ	0.11 U	0.111 U
Sodium	7440235	mg/kg	9.16 U	8.93 U	8.86 U	7.68 U	22 J	7.14 U	8.7 U	8.59 U		33.6	9.79 J	ļ	27.3 J	14.4 J	36 J	34.9 J	41.5 J	10.4 J	7.26 U	ļ	28 J	23.4 J
Strontium	7440246	mg/kg	6.27	5.33	5.39	6.11	5.68	14.4	7.23	5.95		5.92	5.01	<del> </del>	6.96	5.37	8.23	8.32	5.81	6.2	2.72	ļ	6	5.84
Thallium	7440280	<u></u> +-	0.179 U	0.162 U	0.173 U	0.169 U	0.167 U	0.174 J	0.167 U	0.154 U		0.152 U	0.174 U	<del> </del>	0.167 U	0.149 U	0.18 U	0.184 U	0.18 U	0.166 U	0.168 U	ļ	0.172 U	0.172 U
Uranium	7440611	mg/kg	17.4	9.91	1.35	2.98	0.864	1.08	6.61	4.88		1.43	14.6	ļ	0.658	3.18	11.5	14.3	38.7	11.4	1.61	ļ	3.27	3.45
Vanadium		mg/kg	10.6	13.9	21.7	16.9	14.2	16.1	13.4	14.6		18.1	20.6	ļ	22.3	22.2	11.9	32	23.7	18	10.5	ļ	17.7	16.4
Zinc	7440666	mg/kg	22.9	30.7	23	28.9	15.9	20.2	27.3	23		22.3	22.6	<u> </u>	25.3	23.7	20.6 J	42.5	38.5	29.1	12.7	<u> </u>	21.5 J	22.3 J
Wall	1 220 (702 (	a !	2 (F.04	105.01	1 (25.05	1 2 45 04	1 2 2 5 2 5 7	1 2 5E 05 X	1 1 05 04		Other	1.05.05.111	1 0 45 05	!	1 00 05 111	1.50.05.111	1 0 500 04	1 405.04	1 2 2 5 2 4	5.00.05.1	1 000 05 1	!	1.75.04	1.25.04
Methylmercury	22967926	mg/kg	2.6E-04 =	4.0E-04 =	6.2E-05 =	2.4E-04 =	3.3E-05 J	2.5E-05 J	1.8E-04 =	4.1E-04 =	 	1.8E-05 UJ	9.4E-05 =		1.8E-05 UJ	1.7E-05 UJ	8.7E-04 =	4.8E-04 =	2.3E-04 =	5.6E-05 J	9.8E-05 J		1.7E-04 =	1.3E-04 =
W (70) 1 (1 1 1 1	1 2101021	л !			!	1	!	!	!	Sequent	tial extraction	<u>n</u> !	!	1 100 04 111	!	!	LIOFOATH	LAGE GATH	LIOFOATI	!	!	1 105 04 1	!	!
Mercury (F0) - volatile elemental mercury	+	mg/kg			ļ			<del> </del>	<u> </u>	<del> </del>	9.3E-04 J 0.0044 J	ļ	<u> </u>	1.8E-04 UJ 0.0015 J	<u> </u>		1.9E-04 UJ 0.0033 J	2.0E-04 UJ 0.0028 J	0.0088 =	<del> </del>		1.8E-04 J 0.0071 =		i 
Mercury (F1) - water soluble mercury	NS1022				<del> </del>	- <del> </del>		<del></del>	<del> </del>	<del> </del>	0.0044 J 0.0016 J		<del> </del>	0.0013 J 0.0012 UJ	<u> </u>	<del> </del>		0.0028 J 0.0021 J		 !	- <del> </del>	0.0071 = 0.0035 J		
Mercury (F2) - pH 2 soluble mercury  Mercury (F3) - IN potassium hydroxide extractable mercury	NS1023				·	<del>- </del>	<del> </del>	<del> </del>	<del> </del>	<del> </del>	0.00163	<del></del>	ļ	0.0012 03	<u> </u>		0.0013 3	0.00213	0.0049 =	<del> </del>	- <del> </del>	0.00333		
Mercury (F4) - 12N nitric acid soluble mercury	NS1024 NS1025				<del> </del>	- <del> </del>	<del> </del>	<del> </del>	<del> </del>	<del> </del>	0.008 – 0.0068 J	<del> </del>	<del> </del>	0.019 – 0.0059 UJ	<del> </del>		0.086 – 0.014 J	0.003 – 0.012 J	0.25 =	 	- <del> </del>	0.12 – 0.015 J		<del></del>
Mercury (F4) - 1218 mitre acta soluble mercury  Mercury (F5) - aqua regia soluble mercury		mg/kg		 	· <del> </del>	- <del> </del>	<del> </del>	<del> </del>		} <del>-</del>	0.0068 3			0.0039 03			0.014 J 0.0022 J	0.012 J 0.003 J	0.023 =	i I		0.0133		 
Mercury (F6) - mineral-bound mercury		mg/kg			<del></del>	- <del> </del>	<del> </del>	<del> </del>		<del> </del>	7.2E-04 J			4.8E-04 J			3.4E-04 UJ	3.6E-04 UJ	8.2E-04 J		- <del></del>	9.3E-04 J		
Mercury (FS) - total mercury by summation	+	mg/kg			<del></del>	<del></del>	<del> </del>	<del> </del>		<del> </del>	0.086 =		<del></del>	0.025 =			0.11 =	0.085 =	0.25 =			0.16 =		
vicious (15) - total incloury by summation	1 1131020	mg/kg			!	<u>.                                      </u>	<u> </u>			<u>:                                    </u>	chemistry	<u> </u>	!	0.023 -	!	!	0.11	1 0.003 -	0.23	!	<u> </u>	1 0.10 -	!	<u></u>
Moisture	N544	%	23.9	22.4	22.6	20	21.6	18.2	27.6	23.4		22.7	19.6		23.2	20.6	30.2	29.7	28.5	24.8	18.8	ļ	23	20.9
Sulfide	18496258			20.7 J	11.3 U	34.9	23.6 J	18 J	31.3 J	17.4 J		43.6	22.1 J		27.6 J	26.3 J	15 J	36	43.3	23.6 J	23.7 J		28.2 J	10.9 U
Total organic carbon average	NS2302			14,600	5550	7350	5320	6640	10,700	11,800		7840	10,400	ļ	5850	5990	16,700	22,100	18,800	9000	4810	ļ	8920	11,200
Tomi organic carbon average	1104304	mg/ng	11,200	17,000	i 2220	i 1330	i 3340	i 0040	i 10,700	i 11,000	<u> </u>	i /070	i 10,400	i	i 2020	i 3770	i 10,700	i 44,100	i 10,000	i 7000	i 7010	i	i 0720	i 11,200

Table C.6. BCV RSE creek bank soil data (cont.)

Chemical	CAS	Units	BCT12B-BS	BCT12B-BSL	BCT12B-BSL-D	BCT12B-BSU	BCT12B-BSU-D	BCT12A-BSL	BCT12A-BSU	BCT13-BSL	BCT13-BSU	BCT14-BS	BCT14-BSL	BCT14-BSL-D	BCT14-BSU	BCT14-BSU-D	BCT15-BSL	BCT15-BSU	HCTREF-BS	HCTREF-BSL	HCTREF-BSU
	1	1 1		1	1	1		1	Anions	1			1 014	1	1		1	1	1		T
Chloride	16887006			6.88	2.84	3.07	2.75	3.98	2.54	5.88	5.13	- <b> </b>	81.3	94.5	6.79	3.44	24.8	11.1	ļ	4.25	2.68
Fluoride	16984488			15.9	20.5	18.5	17.6	12.1	12.7	24.7	15.6	- <b> </b>	5.24	5.6	0.407 U	0.639 J	13.6	9.56	ļ	2.49 J	3.09 J
Nitrate	14797558	mg/kg		1.73	1.56	1.71	1.65	1.2 J	1.33	1.7	1.35	<u> </u>	20	23.1	0.395 U	0.372 U	88.5	29.2		0.49 U	1.06 J
Nitrite	14797650	mg/kg		0.411 U	0.408 U	0.413 U	0.385 U	0.443 U	0.397 U	0.434 U	0.438 U	<u>-</u>	0.407 U	0.403 U	0.395 U	0.372 U	0.525 U	0.473 U		0.49 U	0.392 U
Sulfate	14808798	mg/kg		4.72 J	4.34 J	3.78 J	3.46 J	6.02	3.29 J	26.8 J	25.7 J		72.6	74.9	11.9	11.7	31.9	27.8		8.56	9.5
	1			1	1	1			Metals					1			1			! =	!
Aluminum	7429905			8190	8510	8240	8220	10,500	9870	8310	8030		9450	9010	10,200	9910	9020	8430	ļ	7010	5570
Antimony	7440360	mg/kg		0.392 U	0.377 U	1.34 J	0.842 J	0.45 U	0.633 J	0.846 J	1.46 J	<u> </u>	0.382 U	1.05 J	0.385 U	0.396 U	1.56 J	0.471 U	ļ	0.49 U	0.39 U
Arsenic	7440382	mg/kg		4.35	4.06	5.64	6.72	6.07	5.66	5.49	5.98	<u> </u>	5.12	4.12	4.61	4.83	4.54	3.99		3.3	1.85
Barium	7440393	mg/kg		150	147	102	104	136	151	167	165	<u> </u>	44.2	44.5	47.1	45.6	602	380	<u> </u>	102	77
Beryllium	7440417	mg/kg		0.407 J	0.462 J	0.379 J	0.42 J	0.474 J	0.562 J	0.407 J	0.275 J	<u> </u>	0.417 J	0.357 J	0.449 J	0.414 J	1.29	0.982		0.53 J	0.445 J
Boron	7440428	mg/kg		1.19 U	1.14 U	1.11 U	1.09 U	2.67 J	1.24 U	1.23 U	1.2 U		1.16 U	2.15 J	2.71 J	3.62 J	7.1 J	5.75 J		1.48 U	1.18 U
Cadmium	7440439	mg/kg		4.26	4.04	6.1	7.37	6.84	5.52	16.4	8.44	<u> </u>	0.27	0.15 J	0.0598 J	0.0604 J	15.5	7.95	<u> </u>	0.106 J	0.0364 J
Calcium	7440702	mg/kg		2830	2900	1680	1760	4690	3830	6150	7960		4590	4750	1000	992	6600	7960		3160	3290
Chromium	7440473	mg/kg		18.1	18.2	17.7	18.5	18.8	18.7	18.6	20.1		22.9	20.9	23.2	17.5	12.5 J	15		17.7	13.9
Cobalt	7440484	mg/kg		14.6	17.2	16.1	13.9	16.1	19.4	20.4	15.3		13.4	11.9	16.1	15.3	46.2	24.8		10.6	8.52
Copper	7440508	mg/kg		14.2	21	18.9	18.5	21.7	17.9	17	31.3		5.43	4.48	5.44	5.04	15.1	13.7	<u></u>	9.97	6.93
Iron	7439896	mg/kg		16,600	18,000	17,100	15,400	19,400	17,800	19,100	18,400		38,000	30,200	48,700	46,500	16,700	16,800		15,500	12,900
Lead	7439921	mg/kg		20.6	28.8	27.9	22.2	28.6	23.6	25.4	32.9		15.9	13.1	14.6	14.2	21.7	17.7		14.1 J	11.5 J
Lithium	7439932	mg/kg		24	20.5	24.3	32.6	20.6	22.4	28.8	29.9	<del></del>	9.94	10.9	9.86	10	18.8	16.8		16.4	11.8
Magnesium	7439954	mg/kg		1240	1190	943	932	1920	1310	1230	2200	- <del> </del>	804	833	556	550	2630	3300	<del></del>	1680	1440
Manganese		mg/kg		1610	1990	996	941	1610	1740	2420	2030	†	534	485	448	403	10,100	5620	·	847	694
Mercury	7439976	+ <u></u> -+-		1.3 =	0.62 =	1.9 =	1.5 =	1.6 =	0.86 =	1.6 =	7.1 =		0.026 J	0.041 =	0.059 =	0.054 =	0.32 =	0.42 =	·	0.03 =	0.028 =
Molybdenum		mg/kg		0.683 J	0.849 J	0.729 J	0.691 J	0.801 J	0.377 J	1.18 J	0.644 J		0.519 J	0.351 J	0.242 J	0.437 J	0.307 U	0.315 J	<del> </del>	0.297 U	0.236 U
Nickel	7440020	mg/kg		23.1 J	33.1 J	28.6 J	27.9 J	38 J	39.5 J	60.6 J	38.7 J	<del></del>	11.2 J	8.93 J	7.34 J	6.12 J	120	63.3		11.2	9.24
Phosphorous	7723140	mg/kg		172	171	190	196	215	202	215	250	- <del> </del>	77.9	79	105	110	239	255	·	184	113
Potassium		mg/kg		559	563	582	596	678	586	585	650	<del></del>	626	624	541	533	730	768	†	731	548
Selenium	7782492	mg/kg		1.46	1.29	1.56	1.81	1.7	1.77	1.87	1.81	<del></del>	0.48 J	0.471 J	0.507 J	0.528 J	2.35	1.99	<del> </del>	1.52	1.31
Silicon		mg/kg		2980	3020	2660	2540	3560	3220	2520	2930	- <del> </del>	2830	2770	2750	2870	2680	2700	<del> </del>	687	490
Silver		mg/kg		0.119 U	0.114 U	0.111 U	0.109 U	0.136 U	0.124 U	0.167 J	0.578 J	†	1.52	1.3	1.83	1.76	1.36	0.943		0.742 U	0.591 U
Sodium	7440235	++-		11 J	9.75 J	7.77 U	7.65 U	17.9 J	13.6 J	22.6 J	18.1 J	<del></del>	77.5	112	8.16 U	8.4 U	77.4	64.5		17.8 J	13.4 J
Strontium		+ <del></del> +-		9.32	8.24	7.75	8.32	13.4	19.4	17.4	18.1	<del> </del>	11.2	11.3	5.14	5.21	24.6	20.2		7.78	4.44
Thallium	7440240	mg/kg		0.16 U	0.155 U	0.159 U	0.187 J	0.18 U	0.177 U	0.191 U	0.195 J		0.154 J	0.159 J	0.154 J	0.163 U	0.194 U	0.187 U	ļ	0.18 U	0.174 U
Uranium		mg/kg mg/kg		30.3	33.8	60.7	68.6	36.5	81.4	24.8	17.6	<del> </del>	5.06	4.91	1.33	1.43	7.6	6.86		0.18 0	0.174 0
Vanadium	7440611	mg/kg		22.4	24.5	22.9	21.1	27.6	22.3	21.8	22.2	<del>-</del>	25	24.4	28.9	27.1	16.3	16.3		18.3	15.6
Zinc	7440666			46.5	57.5	49.5	48.7	75	52.1	62.9	79.9	† <u>-</u>	11.6	10.6	11.3	9.39	61.2	58.6		35.6	26.2
Zinc	7440000	mg/kg		1 70.5	31.5	17.5	1 40.7	1 /3	Other	02.5	17.7	-	11.0	10.0	1 11.5	7.57	01.2	30.0	1	35.0	20.2
Methylmercury	22967926	mg/kg		9.8E-05 =	1.4E-04 =	2.8E-04 =	2.0E-04 =	2.6E-04 =	4.2E-04 =	7.1E-04 J	7.7E-04 J		1.7E-05 UJ	1.7E-05 UJ	4.0E-05 J	2.5E-05 J	2.5E-04 =	1.7E-04 =	Ī	4.0E-05 J	1.6E-05 UJ
							!	·	Sequential extractio	:		!		!							
Mercury (F0) - volatile elemental mercury	NS1021	mg/kg	1.7E-04 UJ									1.7E-04 UJ							3.7E-04 J		
Mercury (F1) - water soluble mercury		mg/kg	0.026 J					<del></del>		<del> </del>	<del> </del>	0.0011 UJ	-			<u> </u>	†		0.0028 J		<del> </del>
Mercury (F2) - pH 2 soluble mercury	NS1023	mg/kg	0.02 J								<del> </del>	0.0011 UJ							0.0011 UJ		
Mercury (F3) - 1N potassium hydroxide extractable mercury	NS1024	mg/kg	0.39 J							<u></u>	<del></del>	0.0022 UJ				İ	T		0.013 J		
Mercury (F4) - 12N nitric acid soluble mercury			0.15 J		+	- <del> </del>	 	<del> </del>		÷	} ¦	0.017 J			+	† ¦			0.0053 UJ		
Mercury (F5) - aqua regia soluble mercury	NS1026									<u> </u>		0.0031 J				<del></del>	- <del> </del>		0.0012 J		
Mercury (F6) - mineral-bound mercury	NS1027	mg/kg	0.0068 J						<del>- </del>	<del> </del>	<del> </del>	0.0013 J			<del> </del>	<del> </del>	<del></del>		0.016 J		<del> </del>
Mercury (FS) - total mercury by summation	NS1028			-†	·†	- <del> </del>		†	-†	t	<del></del>	0.021 J			<del> </del>	†	-†		0.033 J	¦	†
		001		1	1	1	1	1	Wet chemistry	L	I	1	I	1	ı	1	-	L	1	1	1
Moisture	N544	%		21.2	21.8	20.3	21.1	26.9	22.3	28.3	27.1		19.3	19.5	18.3	17.9	39.3	34.1		35	21.3
Sulfide	18496258			11.9 J	21.3 J	17.4 J	10.9 U	32.5 J	21.1 J	20 J	21.7 J	- <del> </del>	11.1 U	12 J	22.4 J	24 J	25.9 J	39.7		31.1 J	21.8 J
Total organic carbon average	NS2302	mg/kg		12,700	11,000	10,300	10,500	16,600	13,400	15,900	25,000	- <del> </del>	7950	5370	3790	3360	24,200	26,900		12,300	6850
E = Estimated, matrix interference.	,	00		.,,,,,,	, ,,,,,	.,	,	,	,		,	1					, ,	,	1	, ,	

E = Estimated, matrix interference.

J = The analyte was positively identified; the associated numerical value is the approximate concentration of the analyte in the sample.

U = The analyte was analyzed for, but was not detected above the reported sample quantitation limit.

UJ = Analyte or compound was not detected above the reported detection limit, and the reported detection limit is approximated due to quality deficiency.

<sup>=</sup> Validated result, which is detected and unqualified.

<sup>-- =</sup> Analyte or compound not sampled for.

BCV = Bear Creek Valley CAS = Chemical Abstract Service RSE = remedial site evaluation

Table C.7. BCV RSE channel sediment data

Chemical	CAS	Units	встз-сн	вст4-сн	вст7-сн	вст8-сн	вст9-сн	ВСТ10-СН	вст11-сн	ВСТ12В-СН	BCT12B-CH-D	BCT12A-CH	вст13-сн	ВСТ14-СН	BCT14-CH-D	ВСТ15-СН
	•	•	•			P	Anions	•		•			•		•	•
Chloride	16887006	mg/kg	4.36	3.53	9.88 J	11.5 J	12.9 J	11.5	14.4	21.2	20.5	32.6	31.2	20.9	25	11.5
Fluoride	16984488	mg/kg	0.905 J	0.884 J	1.17 J	1.62	1.83	1.53 J	2.01 J	3.67 J	3.6 J	3.78 J	4.58 J	8.6	9	6.06
Nitrate	14797558	mg/kg	1.22 J	1.1 J	0.464 U	0.459 U	0.894 J	1.08 J	1.43	1.01 J	0.959 J	0.975 J	1.33 J	11.2	14.8	32.4
Nitrite	14797650	mg/kg	0.455 U	0.419 U	0.464 U	0.459 U	0.445 U	0.429 U	0.414 U	0.458 U	0.461 U	0.445 U	0.438 U	0.51 U	0.493 U	0.531 U
Sulfate	14808798	mg/kg	22.1	10.4	20.6	25	35.5	33.7	9.4	13	12.6	14	18.3	30.3	36.3	34.3
	•		•	•	•	I	Metals	•	•	•	•				•	•
Aluminum	7429905	mg/kg	5890	5690	2770	7670 J	5950 J	7040	6130	4830	5990	6510	8230	10,900	10,800	15,400
Antimony	7440360	mg/kg	1.45 J	2.15 J	0.44 U	0.406 U	0.417 U	0.725 J	0.652 J	0.422 U	0.428 U	0.423 U	0.452 U	1.33 J	1.32 J	0.509 U
Arsenic	7440382	mg/kg	6.22	9.8	1.78	15.1	3.46	7.02	11	9.84	8.56	12.2	13.2	5.93	6	6.99
Barium	7440393	mg/kg	71.4	56.4	47.4	80.6	77.8	107	137	176	259	324	615	364	310	1590
Beryllium	7440417	mg/kg	1.03	1.11	0.309 J	1.2	0.901	1.36	1.12	0.497 J	0.545 J	0.61 J	0.775	1.1	1.09	2.56
Boron	7440428	mg/kg	3.74 J	7.31	1.33 U	9.31	3.93 J	9.82	6.2	1.28 U	1.3 U	1.28 U	1.37 U	1.43 U	7.43	10.4
Cadmium	7440439	mg/kg	0.207 J	0.26	0.249 J	1.52	1.36	0.925	2.81	3.18	3.03	2.96	4.7	4.51	5.01	38.5
Calcium	7440702	mg/kg	1220	984	2270	2310 E	3380 E	3670	7510	4890	4830	6500	5600	4730	4630	7680
Chromium	7440473	mg/kg	30.9	51	8.12	55.7	12.7	76	39.6	27.1	31	38.5	48.8	32.8	37.3	25.4 J
Cobalt	7440484	mg/kg	12	16.4	4.99	19.3	11.1	25.8	18.3	19.1	25.2	27.2	36.5	42	39.4	106
Copper		mg/kg	8.23	7.54	3.43	7.51	10.7	11.4	10.2	9.89	11	12.4	17.3	14	13.9	25.4
Iron	7439896	mg/kg	23,800	32,100	7120	29,600 J	12,000 J	44,400	34,100	31,700	35,700	42,900	49,000	52,300	53,200	30,700
Lead	7439921	mg/kg	15.9	16.2	5.17	19.4 J	13.1 J	25.1	22.9	21.3 J	27.3 J	35.2 J	42.9 J	35	28.3	37.7
Lithium	7439932	mg/kg	12.4	9.81	8.4	16.7	17.3	11.1	16.7	10.3	11.3	13.5	14.1	12.3	12.4	23.9
Magnesium	7439954	mg/kg	1380	1290	1230	1420	1350	1990	2780	1280	1050	1320	1430	1320	1300	2390
Manganese	7439965	mg/kg	817	733	478	1140	641	1430	1900	2200	2890	3380	7000	4360	3930	29,600
Mercury	7439976	mg/kg	0.029 J	0.026 J	0.086 =	0.11 =	0.44 =	0.095 J	0.16 J	0.33 =	0.27 J	0.53 =	0.34 =	0.083 =	0.063 =	0.43 =
Molybdenum	7439987	mg/kg	0.546 J	1.14 J	0.267 U	0.246 U	0.253 U	1.15 J	1.22 J	0.255 U	0.259 U	0.256 U	0.274 U	0.41 J	0.435 J	0.703 J
Nickel	7440020	mg/kg	18.3 J	20.1 J	6	22.8 E	17.3 E	24.5	23.1	22.3	30.2	27.2	39.5	38.7 J	34.5 J	289
Phosphorous	7723140	mg/kg	239	340	131	360	138	375	253	196	167	185	259	104	106	245
Potassium	7440097	mg/kg	763	570	395	746	556	610 J	659 J	375	407	478	718	873	853	976
Selenium	7782492	mg/kg	0.778 J	0.903 J	0.581 J	0.965 J	0.666 J	1.21 J	1.64	1.84	1.8	2.19	2.23	1.43 J	1.47 J	2.89
Silicon	7440213	mg/kg	2130	1750	991	1550	1870	3490	2310	458 J	520 J	489 J	572 J	3740	3630	3040
Silver	7440224	mg/kg	0.298 J	0.266 J	0.133 U	0.615 U	0.126 U	0.118 U	0.122 U	0.639 U	0.648 U	0.641 U	1.37 U	2.51	2.51	4.25
Sodium	7440235	mg/kg	9.66 U	8.76 U	22.8 J	35.8 J	42 J	30.3 J	36.8 J	29.4 J	33.1	48	56.2	37.7	34.6	75.9
Strontium	7440246	mg/kg	6.52	4.31	3.89	5.85	4.69	7.23	20.2	20.9	10.5	11.8	25.9	13	13.4	25.5
Thallium	7440280	mg/kg	0.18 U	0.175 U	0.177 U	0.176 U	0.189 U	0.174 U	0.162 U	0.183 U	0.173 U	0.187 U	0.196 U	0.217 J	0.225 U	0.206 U
Uranium	7440611	mg/kg	2.36	4.36	3.29	19.9	12	8.27	8.98	7.53	8.54	15.4	14.6	19.8	19.8	11.1
Vanadium	7440622	mg/kg	31.5	46.1	5.97	43	15.5	45.6	39.6	27.1	36.2	41.3	43.6	35.3	35	26.3
Zinc	7440666	mg/kg	34.7	37.6	15.5 J	46.9	31.7	62.8 J	44.1 J	117	60.8	60.8	63.1	50.5	50.1	116
							Other									
Methylmercury	22967926	mg/kg	6.3E-05 J	5.7E-05 J	2.5E-04 =	1.7E-04 =	8.2E-04 =	2.7E-04 =	1.0E-04 =	1.8E-04 =	2.3E-04 =	4.0E-04 =	1.9E-04 =	2.5E-05 UJ	4.2E-05 J	3.1E-05 J
	1	!		·	·	<del></del>	ial extraction	·					· · · · · · · · · · · · · · · · · · ·		·	т
Mercury (F0) - volatile elemental mercury	NS1021	mg/kg	ļ		2.1E-04 UJ	2.0E-04 UJ	1.8E-04 UJ	<del></del>	5.2E-04 J	1.9E-04 J			<del></del>	0.0074 J	ļ	ļ
Mercury (F1) - water soluble mercury	NS1022	mg/kg	<del> </del>		0.0016 J	0.0034 J	0.0086 =	<del></del>	0.0053 =	0.0065 =			<del> </del>	0.0013 UJ	<del></del>	<del></del>
Mercury (F2) - pH 2 soluble mercury	NS1023	mg/kg	<del> </del>	<del> </del>	0.0016 J	0.0013 J	0.0048 =	<del></del>	0.0043 J	0.0056 J			<del> </del>	0.0013 UJ	<del> </del>	<del></del>
Mercury (F3) - 1N potassium hydroxide extractable mercury	NS1024	mg/kg	ļ	<del> </del>	0.045 =	0.07 =	0.17 =	<del></del>	0.037 =	0.099 =			<del> </del>	0.0027 UJ	<del> </del>	<del></del>
Mercury (F4) - 12N nitric acid soluble mercury	NS1025	mg/kg	<del></del>	<del> </del>	0.0066 UJ	0.0096 J	0.043 =	. <del> </del>	0.0083 J	0.02 =			<del> </del>	0.032 J	}	<del></del>
Mercury (F5) - aqua regia soluble mercury	NS1026	mg/kg		 	0.0024 J	0.0027 J	0.011 =		0.04 =	0.085 =			ļi	0.023 J	ļ	
Mercury (F6) - mineral-bound mercury	NS1027	mg/kg		<del> </del>	3.8E-04 UJ	7.1E-04 J	0.0012 J		0.0019 J	9.1E-04 J			<del> </del>	4.6E-04 J	<del></del>	
Mercury (FS) - total mercury by summation	NS1028	mg/kg	!	<u> </u>	0.051 =	0.088 =	0.24 =	<u> </u>	0.097 =	0.22 =			<u> </u>	0.055 J	!	<u> </u>
Maistone	NEAA	0/	. 27.0	. 22.1	21.5		chemistry	1 220	22.0	20.2	20.4	21.0	20.2	27	1 202	20.6
Moisture	N544	%	27.9	23.1	31.5	29.4	31.6	23.8	22.9	29.2	30.4	31.8	30.3	37	38.2	38.6
Sulfide	18496258	mg/kg	24.4 J	27.6 J	13.1 J	36.7	36.4	12.8 J	11.1 U	12.4 U	12.6 U	12.7 U	17.8 J	14 J	14.4 U	30.7 J
Total organic carbon average	NS2302	mg/kg	2440	2500	7250	5080	10,800	8390	6320	5290	7160	7710	6760	4420	3760	21,200
E = Estimated, matrix interference.																

E = Estimated, matrix interference.

J = The analyte was positively identified; the associated numerical value is the approximate concentration of the analyte in the sample.

U = The analyte was analyzed for, but was not detected above the reported sample quantitation limit.

UJ = Analyte or compound was not detected above the reported detection limit, and the reported detection limit is approximated due to quality deficiency.

<sup>=</sup> Validated result, which is detected and unqualified.

<sup>-- =</sup> Analyte or compound not sampled for.

BCV = Bear Creek Valley CAS = Chemical Abstract Service RSE = remedial site evaluation

# APPENDIX D. PARTICLE SIZE ANALYSIS

FLOODPLAIN SOIL

#### **Particle-Size Distribution**

Form No. TR-D7928-3 Revision No. 0

Log No.:

Revision Date: 03/18/2019


43-4016

BCT1-FP



ASTM D7928 & D6913

	S&ME, Inc.	, 1413 Topsid	e Road, Lou	isville,	TN 37777	
S&ME Project #:	23430142				Report Date:	2/23/2024
Project Name:	UCOR-Geotech				Test Start Date:	2/2/2024
Client Name:	RSI EnTech, LLC					
Address:	203 Victorious Blvd.	, Oak Ridge, TI	N 37830			
Sample ID:	BCR0039-03	Type:	REG		Sample Dat	e: 1/4/2024
	-				Dept	h: NP



Assumed Specific Gravity: Coarse Sand: 0% 2.65 8.4% Clay: Liquid Limit **TNP** Plastic Limit TNP Plastic Index **TNP** 

References / Comments / Deviations:

Victoria Igoe Technical Responsibility

Signature

Senior Engineering Technician Position

2/23/2024 Date

Form No. TR-D7928-3

Revision No. 0

Log No.:

Revision Date: 03/18/2019

43-4016

BCT2-FP



ASTM D7928 & D6913

	S&ME, Inc.	, 1413 Topsid	e Road, Lou	isville,	TN 37777	
S&ME Project #:	23430142				Report Date:	2/19/2024
Project Name:	UCOR-Geotech				Test Start Date:	2/7/2024
Client Name:	RSI EnTech, LLC					
Address:	203 Victorious Blvd.,	Oak Ridge, TN	l 37830			
Sample ID:	BCR0042-03	Type:		Sample Dat	e: 1/4/2024	
					Dept	h: NP



Cobbles	< 300 mm (12") and > 75 mm	Fine Sand	< 0.425 mm and > 0.075 mm
Gravel	< 75 mm and > 4.75 mm (#4)	Silt and Clav	< 0.075 mm
Coarse Sand	< 4.75 mm and >2.00 mm	Clav	<0.002 mm
Medium Sand	< 2.00 mm and > 0.425 mm		

IVICAIAIII SAIIA	<del>Jo IIIIII ana</del>	<u> </u>				_
Maximum Particle Size:	#4	Gravel:	0%	Medium Sand:	4%	
Silt & Clay (% Passing #200):	75%	Total Sand:	25%	Fine Sand:	20%	
Assumed Specific Gravity:	2.65	Coarse Sand:	1%	Clay:	8.4%	
Liquid Limit	TNP	Plastic Limit	TNP	Plastic Index	TNP	

References / Comments / Deviations:

<u>Tyler Copeman</u> Technical Responsibility Signature

<u>Lab Services Manager</u> *Position*  2/19/2024 Date

Form No. TR-D7928-3

Revision No. 0

Log No.:

Revision Date: 03/18/2019


43-4016

BCT3-FP



ASTM D7928 & D6913

S&ME, Inc., 1413 Topside Road, Louisville, TN 37777								
S&ME Project #:	23430142				Report Date:	2/1	9/2024	
Project Name:	UCOR-Geotech				Test Start Date:	2/7	7/2024	
Client Name:	RSI EnTech, LLC							
Address:	203 Victorious Blvd.	, Oak Ridge, TN	N 37830					
Sample ID:	BCR0045-03	Type:	REG		Sample Date: 1/8/		1/8/2024	
					Dept	h:	NP	



Cobbles	< 300 mm (12") and > 75 mm	Fine Sand	< 0.425 mm and > 0.075 mm
Gravel	< 75 mm and > 4.75 mm (#4)	Silt and Clav	< 0.075 mm
Coarse Sand	< 4.75 mm and >2.00 mm	Clav	<0.002 mm
Medium Sand	< 2.00 mm and > 0.425 mm		

Maximum Particle Size:	3/8"	Gravel:	2%	Medium Sand:	6%	
Silt & Clay (% Passing #200):	77%	Total Sand:	21%	Fine Sand:	13%	
Assumed Specific Gravity:	2.65	Coarse Sand:	2%	Clay:	11.8%	
Liquid Limit	TNP	Plastic Limit	TNP	Plastic Index	TNP	

References / Comments / Deviations:

<u>Tyler Copeman</u> Technical Responsibility Signature

<u>Lab Services Manager</u> *Position*  2/19/2024 Date

Form No. TR-D7928-3

Revision No. 0

Log No.:

Revision Date: 03/18/2019


43-4016

BCT4-FP



ASTM D7928 & D6913

	S&ME, Inc., 1413 Topside Road, Louisville, TN 37777						
S&ME Project #:	23430142				Report Date:		1/0/1900
Project Name:	UCOR-Geotech				Test Start Date:	2/1	2/24-2/16/24
Client Name:	RSI EnTech, LLC						
Address:	203 Victorious Blvd.,	Oak Ridge, TN	I 37830				
Sample ID:	BCR0048-03	Type:	REG		Sample D	ate:	1/8/2024
					De <sub>l</sub>	oth:	NP



Cobbles	< 300 mm (12") and > 75 mm	Fine Sand	< 0.425 mm and > 0.075 mm
Gravel	< 75 mm and > 4.75 mm (#4)	Silt and Clav	< 0.075 mm
Coarse Sand	< 4.75 mm and >2.00 mm	Clav	<0.002 mm
Medium Sand	< 2.00 mm and > 0.425 mm		

Wicaiaiii Sana 📗 🔻 🔼	<del>.oo iiiii ana</del>	<u> </u>				_
Maximum Particle Size:	#10	Gravel:	0%	Medium Sand:	3%	
Silt & Clay (% Passing #200):	58%	Total Sand:	42%	Fine Sand:	39%	
Assumed Specific Gravity:	2.65	Coarse Sand:	0%	Clay:	8.1%	
Liquid Limit	TNP	Plastic Limit	TNP	Plastic Index	TNP	

References / Comments / Deviations:

Tyler Copeman Technical Responsibility

Lab Services Manager Position

2/19/2024 Date

### **Particle-Size Distribution**

Form No. TR-D7928-3 Revision No. 0


Revision Date: 03/18/2019

BCT5-FP



Log No.: 43-4016 ASTM D7928 & D6913

3									
	S&ME, Inc., 1413 Topside Road, Louisville, TN 37777								
S&ME Project #:	23430142				Report Date:		3/12/2024		
Project Name:	UCOR-Geotech				Test Start Date:	3/	7/24-3/1/24		
Client Name:	RSI EnTech, LLC								
Address:	203 Victorious Blvd.,	, Oak Ridge, TN	l 37830						
Sample ID:	BCR0051-04	Type:	REG		Sample Da	ite:	2/1/2024		
	·	·	·		Dep	th:	NP		



CODDIES	SOUTHING 2 / 3 / 10 / 13 / 11 / 11 / 12 / 3 / 10 / 12 / 3 / 10 / 12 / 3 / 10 / 10 / 10 / 10 / 10 / 10 / 10		: Saliu	1 \ 0.423 IIIII and /	0.07.3 111111
Gravel	< 75 mm and > 4.75 mm (#4)	Silt a	nd Clav	< 0.075 mr	n
Coarse Sand	< 4.75 mm and >2.00 mm		Clav	<0.002 mr	n
Medium Sand	< 2.00 mm and > 0.425 mm				
Maximum Particle S	Size: 3/4"	Gravel:	0%	Medium Sand:	12%

Silt & Clay (% Passing #200): 41% Total Sand: 59% Fine Sand: 46% Assumed Specific Gravity: 2.65 Coarse Sand: 1% Clay: 10.3% TNP TNP TNP Liquid Limit Plastic Limit Plastic Index

References / Comments / Deviations:

Tyler Copeman Technical Responsibility

Lab Services Manager Position

3/12/2024 Date

#### **Particle-Size Distribution**

Form No. TR-D7928-3 Revision No. 0

Revision Date: 03/18/2019

BCT6-FP



Log No.: 43-4016 ASTM D7928 & D6913

S&ME, Inc., 1413 Topside Road, Louisville, TN 37777							
S&ME Project #:	23430142				Report Date:		3/15/2024
Project Name:	UCOR-Geotech				Test Start Date:	3/1	0/24-3/13/24
Client Name:	RSI EnTech, LLC						
Address:	203 Victorious Blvd.,	Oak Ridge, TN	I 37830				
Sample ID:	BCR0055-04	Туре:	REG		Sample D	ate:	2/1/2024
					De	pth:	NP



Silt & Clay (% Passing #200): 74% Total Sand: 22% Fine Sand: 17% Assumed Specific Gravity: 2.65 Coarse Sand: 2% Clay: 12.8% Liquid Limit TNP Plastic Limit **TNP** Plastic Index TNP

References / Comments / Deviations:

<u>Tyler Copeman</u> Technical Responsibility Signature

<u>Lab Services Manager</u> *Position*  3/15/2024 Date

Revision No. 0

Form No. TR-D7928-3

Revision Date: 03/18/2019

BCT7-FP



Log No.: 43-4016 ASTM D7928 & D6913

S&ME, Inc., 1413 Topside Road, Louisville, TN 37777									
S&ME Project #:	23430142				Report Date:		3/15/2024		
Project Name:	UCOR-Geotech				Test Start Date:	3/1	0/24-3/13/24		
Client Name:	RSI EnTech, LLC								
Address:	203 Victorious Blvd.,	Oak Ridge, TN	37830		•				
Sample ID:	BCR0059-04	Туре:	REG		Sample D	ate:	2/7/2024		
					De	oth:	NP		

Sample Description: **Brown Sandy Silt** 1" 3/4" 3/8" #4 #20 #40 #60 #100 #200 100% 90% 80% 70% Percent Passing 60% 50% 40% 30% 20% 10% 0% 10 0.1 0.01 0.001 100 Particle Size (mm) < 300 mm (12") and > 75 mm < 0.425 mm and > 0.075 mm Cohbles Fine Sand

CODDIC3	<u> </u>		- Jana	<u> </u>	<del></del>
Gravel	< 75 mm and > 4.75 mm (#4)	Silt a	nd Clav	< 0.075 mm	n
Coarse Sand	< 4.75 mm and >2.00 mm		Clav	<0.002 mm	1
Medium Sand	< 2.00 mm and > 0.425 mm				
Maximum Particle S	Size: 3/8"	Gravel:	0%	Medium Sand:	3%

Silt & Clay (% Passing #200): 60% Total Sand: 40% Fine Sand: 37% Assumed Specific Gravity: 2.65 Coarse Sand: 0% Clay: 10.0% TNP TNP **TNP** Liquid Limit Plastic Limit Plastic Index

References / Comments / Deviations:

Tyler Copeman Technical Responsibility

Lab Services Manager Position

3/15/2024 Date

Form No. TR-D7928-3 Revision No. 0

Revision Date: 03/18/2019

BCT8-FP



Log No.: 43-4016 ASTM D7928 & D6913

S&ME, Inc., 1413 Topside Road, Louisville, TN 37777									
S&ME Project #:	23430142				Report Date:		3/18/2024		
Project Name:	UCOR-Geotech				Test Start Date:	3/1	2/24-3/14/24		
Client Name:	RSI EnTech, LLC								
Address:	203 Victorious Blvd.	Oak Ridge, TN	I 37830						
Sample ID:	BCR0081-04	Туре:	REG		Sample D	ate:	2/5/2024		
					De	pth:	NP		

Sample Description: Brown Silty Sand with Gravel 1" 3/4" 3/8" #20 #40 #60 #100 #200 100% 90% 80% 70% Percent Passing 60% 50% 40% 30% 20% 10% 0% 10 1 0.1 0.01 0.001 100 Particle Size (mm) < 300 mm (12") and > 75 mm < 0.425 mm and > 0.075 mm Cobbles Fine Sand Gravel < 75 mm and > 4.75 mm (#4) Silt and Clav < 0.075 mm < 4.75 mm and >2.00 mm <0.002 mm Coarse Sand Clav

Medium Sand	< 2.00	mm and >	0.425 mm				
Maximum Particle S	ize:	3/4''		Gravel:	17%	Medium Sand:	14%
Silt & Clay (% Passing #2	00):	42%	To	otal Sand:	41%	Fine Sand:	20%
Assumed Specific Gra	vity:	2.65	Coa	rse Sand:	7%	Clay:	7.7%
Liquid L	imit	TNP	Pla	stic Limit	TNP	Plastic Index	TNP

References / Comments / Deviations:

<u>Tyler Copeman</u> Technical Responsibility Signature

<u>Lab Services Manager</u> <u>Position</u> 3/18/2024 Date

Form No. TR-D7928-3 Revision No. 0


Revision Date: 03/18/2019

BCT9-FP



Log No.: 43-4016 ASTM D7928 & D6913

S&ME, Inc., 1413 Topside Road, Louisville, TN 37777										
S&ME Project #:	23430142				Report Date:	:	3/14/2024			
Project Name:	UCOR-Geotech				Test Start Date:	3/1	0/24-3/13/24			
Client Name:	RSI EnTech, LLC									
Address:	203 Victorious Blvd.	, Oak Ridge, TN	I 37830							
Sample ID:	BCR0070-04	Туре:	REG		Sample Date: 2/		2/5/2024			
					De	oth:	NP			



Silt & Clay (% Passing #200):78%Total Sand:22%Fine Sand:15%Assumed Specific Gravity:2.65Coarse Sand:2%Clay:18.1%Liquid LimitTNPPlastic LimitTNPPlastic IndexTNP

References / Comments / Deviations:

<u>Tyler Copeman</u> Technical Responsibility Signature

<u>Lab Services Manager</u> *Position*  3/14/2024 Date

Form No. TR-D7928-3

Revision No. 0

Log No.:

Revision Date: 03/18/2019

43-4016

BCT10-FP



ASTM D7928 & D6913

	S&ME, Inc., 1413 Topside Road, Louisville, TN 37777									
S&ME Project #:	23430142				Report Date:		2/20/2024			
Project Name:	UCOR-Geotech				Test Start Date:	2/1	7/24-2/20/24			
Client Name:	RSI EnTech, LLC									
Address:	203 Victorious Blvd.	, Oak Ridge, TN	l 37830							
Sample ID:	BRC0074-03	Type:	REG		Sample D	ate:	1/11/2024			
					De <sub>l</sub>	oth:	NP			

Sample Description: **Brown Sandy Clay** 1" 3/4" #60 #100 #200 100% 90% 80% 70% Percent Passing 60% 50% 40% 30% 20% 10% 0% 0.1 0.01 0.001 Particle Size (mm)

Cobbles	< 300 mm (12") and > 75 mm	Fine Sand	< 0.425 mm and > 0.075 mm
Gravel	< 75 mm and > 4.75 mm (#4)	Silt and Clav	< 0.075 mm
Coarse Sand	< 4.75 mm and >2.00 mm	Clav	<0.002 mm
Medium Sand	< 2.00 mm and > 0.425 mm		

Tricatatii Saria		0. 123 111111				
Maximum Particle Size:	3/8"	Gravel:	1%	Medium Sand:	6%	
Silt & Clay (% Passing #200):	60%	Total Sand:	39%	Fine Sand:	33%	
Assumed Specific Gravity:	2.65	Coarse Sand:	0%	Clay:	7.2%	
Liquid Limit	TNP	Plastic Limit	TNP	Plastic Index	TNP	

References / Comments / Deviations:

<u>Tyler Copeman</u> Technical Responsibility Signature

<u>Lab Services Manager</u> *Position*  2/20/2024 Date

Revision No. 0

Form No. TR-D7928-3

Revision Date: 03/18/2019

BCT11-FP




Log No.: 43-4016 ASTM D7928 & D6913

S&ME, Inc., 1413 Topside Road, Louisville, TN 37777									
S&ME Project #:	23430142				Report Date:		3/18/2024		
Project Name:	UCOR-Geotech				Test Start Date:	3/1	2/24-3/14/24		
Client Name:	RSI EnTech, LLC								
Address:	203 Victorious Blvd.,	Oak Ridge, TN	37830						
Sample ID:	BCR0077-04	Туре:	REG		Sample D	ate:	1/31/2024		
					De	pth:	NP		

Sample Description:

Brown Clay with Sand

1" 3/4" 3/8" #4 #10 #20 #40 #60 #100 #200



Cobbles	< 300 mm (12") and > 75 mm	Fine Sand	< 0.425 mm and > 0.075 mm
Gravel	< 75 mm and > 4.75 mm (#4)	Silt and Clav	< 0.075 mm
Coarse Sand	< 4.75 mm and >2.00 mm	Clav	<0.002 mm
Medium Sand	< 2.00 mm and > 0.425 mm		

Maximum Particle Size:	3/8''	Gravel:	0%	Medium Sand:	1%
Silt & Clay (% Passing #200):	75%	Total Sand:	25%	Fine Sand:	23%
Assumed Specific Gravity:	2.65	Coarse Sand:	1%	Clay:	11.9%
Liquid Limit	TNP	Plastic Limit	TNP	Plastic Index	TNP

References / Comments / Deviations:

Tyler Copeman
Technical Responsibility

Signature

<u>Lab Services Manager</u> *Position*  3/18/2024 Date

Revision No. 0

Log No.:

Revision Date: 03/18/2019


43-4016

### BCT12B-FP



ASTM D7928 & D6913

S&ME, Inc., 1413 Topside Road, Louisville, TN 37777											
S&ME Project #:	23430142				Report Date:		1/0/1900				
Project Name:	UCOR-Geotech				Test Start Date:	2/1	2/24-2/16/24				
Client Name:	RSI EnTech, LLC										
Address:	203 Victorious Blvd	., Oak Ridge, TN	37830								
Sample ID:	BRC0065-04	Type:	REG		Sample D	ate:	1/10/2024				
					De	pth:	NP				



Gravel	< 75	<u>mm and &gt; 4.</u>	75 mm (#4)	Silt	: and Clay	< 0.075 m	m
Coarse Sand	< 4.75	5 mm and >2.	00 mm (#10)		Clav	<0.002 m	m
Medium Sand	< 2.	.00 mm and >	0.425 mm				
Maximum Particle S	ize:	3/4''		Gravel:	4%	Medium Sand:	14%
Silt & Clay (% Passing #2	200):	65%	To	tal Sand:	31%	Fine Sand:	13%
Assumed Specific Gra	vity:	2.65	Coa	rse Sand:	4%	Clay:	12.5%
Liquid L	imit	TNP	Pla	stic Limit	TNP	Plastic Index	TNP

References / Comments / Deviations:

Tyler Copeman
Technical Responsibility

Signature

<u>Lab Services Manager</u> *Position*  2/19/2024 Date

Form No. TR-D7928-3


Revision No. 0

Revision Date: 03/18/2019

BCT12B-FP



Log No.:	43-4016	ASTM D7928 & D6913						
		S&ME, Inc	, 1413 Topsid	e Road, Loui	isville, T	N 37777		
S&ME Proje	ect #:	23430142				Report Date:		2/20/2024
Project Nam	ne:	UCOR-Geotech				Test Start Date:	2/1	2/24-2/16/24
Client Name	e:	RSI EnTech, LLC						
Address:		203 Victorious Blvd	., Oak Ridge, TN	I 37830				
Sample ID:	·	BRC0066-03	Type:	REG		Sample D	ate:	1/11/2024
						De	pth:	NP



Cobbles	< 300 mm (12") and > 75 mm	Fine Sand	< 0.425 mm and > 0.075 mm
Gravel	< 75 mm and > 4.75 mm (#4)	Silt and Clav	< 0.075 mm
Coarse Sand	< 4.75 mm and >2.00 mm	Clav	<0.002 mm
Medium Sand	< 2.00 mm and > 0.425 mm		

Maximum Particle Size:	3/4''	Gravel:	2%	Medium Sand:	16%
Silt & Clay (% Passing #200):	62%	Total Sand:	36%	Fine Sand:	18%
Assumed Specific Gravity:	2.65	Coarse Sand:	2%	Clay:	11.5%
Liquid Limit	TNP	Plastic Limit	TNP	Plastic Index	TNP

References / Comments / Deviations:

Tyler Copeman Technical Responsibility

Lab Services Manager Position

2/20/2024 Date

Form No. TR-D7928-3

Revision No. 0

Log No.:

Revision Date: 03/18/2019


43-4016

BCT12A-FP



ASTM D7928 & D6913

S&ME, Inc., 1413 Topside Road, Louisville, TN 37777							
S&ME Project #:	23430142				Report Date:		2/22/2024
Project Name:	UCOR-Geotech				Test Start Date:	2/2	0/24-2/22/24
Client Name:	RSI EnTech, LLC						
Address:	203 Victorious Blvd.	, Oak Ridge, TN	I 37830				
Sample ID:	Sample ID: BRC0106-03 Type: REG			Sample D	ate:	1/10/2024	
		-			De	oth:	NP



Cobbles	< 300 mm (12") and > 75 mm	Fine Sand	< 0.425 mm and > 0.075 mm
Gravel	< 75 mm and > 4.75 mm (#4)	Silt and Clav	< 0.075 mm
Coarse Sand	< 4.75 mm and >2.00 mm	Clav	<0.002 mm
Medium Sand	< 2.00  mm and  > 0.425  mm		

Maximum Particle Size:	3/8''	Gravel:	0%	Medium Sand:	7%	
Silt & Clay (% Passing #200):	79%	Total Sand:	21%	Fine Sand:	13%	
Assumed Specific Gravity:	2.65	Coarse Sand:	1%	Clay:	14.0%	
Liquid Limit	TNP	Plastic Limit	TNP	Plastic Index	TNP	

References / Comments / Deviations:

<u>Tyler Copeman</u> Technical Responsibility Signature

<u>Lab Services Manager</u> *Position*  2/22/2024 Date

Form No. TR-D7928-3

Revision No. 0

Log No.:

Revision Date: 03/18/2019


43-4016

BCT13-FP



ASTM D7928 & D6913

S&ME, Inc., 1413 Topside Road, Louisville, TN 37777							
S&ME Project #:	23430142				Report Date:		2/20/2024
Project Name:	UCOR-Geotech				Test Start Date:	2/1	7/24-2/20/24
Client Name:	RSI EnTech, LLC						
Address:	203 Victorious Blvd	., Oak Ridge, TN	I 37830				
Sample ID:	BRC0085-03	Type:	REG		Sample D	ate:	1/11/2024
					De	oth:	NP



Cobbles	< 300 mm (12") and > 75 mm	Fine Sand	< 0.425 mm and > 0.075 mm
Gravel	< 75 mm and > 4.75 mm (#4)	Silt and Clav	< 0.075 mm
Coarse Sand	< 4.75 mm and >2.00 mm	Clav	<0.002 mm
Medium Sand	< 2.00 mm and > 0.425 mm		

Wicaidili Salid \ Z.C	<u> </u>	/ U.TEJ IIIII				4
Maximum Particle Size:	#4	Gravel:	0%	Medium Sand:	10%	
Silt & Clay (% Passing #200):	55%	Total Sand:	45%	Fine Sand:	34%	
Assumed Specific Gravity:	2.65	Coarse Sand:	1%	Clay:	15.0%	
Liquid Limit	TNP	Plastic Limit	TNP	Plastic Index	TNP	

References / Comments / Deviations:

<u>Tyler Copeman</u> Technical Responsibility Signature

<u>Lab Services Manager</u> *Position*  2/20/2024 Date

Revision No. 0

Log No.:

Revision Date: 03/18/2019

43-4016

BCT14-FP



ASTM D7928 & D6913

	S&ME, Inc.	uisville,	TN 37777				
S&ME Project #:	23430142				Report Date:		2/22/2024
Project Name:	Name: UCOR-Geotech		Test Start Date:	2/2	20/24-2/22/24		
Client Name:	RSI EnTech, LLC						
Address:	203 Victorious Blvd.,	Oak Ridge, TN	I 37830				
Sample ID:	BRC0090-04	Type:	REG		Sample D	ate:	12/28/2023
					De <sub>l</sub>	oth:	NP

Sample Description: Brown Clay with Sand 1" 3/4" #40 #60 #100 #200 100% 90% 80% 70% Percent Passing 60% 50% 40% 30% 20% 10% 0% 0.1 0.01 0.001 Particle Size (mm)

Cobbles	< 300 mm (12") and > 75 mm	Fine Sand	< 0.425 mm and > 0.075 mm
Gravel	< 75 mm and > 4.75 mm (#4)	Silt and Clav	< 0.075 mm
Coarse Sand	< 4.75 mm and >2.00 mm	Clav	<0.002 mm
Medium Sand	< 2.00 mm and > 0.425 mm		

Wicaiaiii Sana	<del>oo miin ana</del>	<u> </u>				_
Maximum Particle Size:	#10	Gravel:	0%	Medium Sand:	6%	
Silt & Clay (% Passing #200):	81%	Total Sand:	19%	Fine Sand:	13%	
Assumed Specific Gravity:	2.65	Coarse Sand:	0%	Clay:	16.0%	
Liquid Limit	TNP	Plastic Limit	TNP	Plastic Index	TNP	

References / Comments / Deviations:

<u>Tyler Copeman</u> Technical Responsibility Signature

<u>Lab Services Manager</u> *Position*  2/22/2024 Date

This report shall not be reproduced, except in full, without the written approval of S&ME, Inc.

3201 Spring Forest Road

Raleigh, NC. 27616

Revision No. 0


Revision Date: 03/18/2019

BCT14-FP



Log No.: 43-4016 ASTM D7928 & D6913

	S&ME, Inc., 1413 Topside Road, Louisville, TN 37777								
S&ME Project #:	5&ME Project #: 23430142						2/22/2024		
Project Name:	UCOR-Geotech	COR-Geotech				2/2	0/24-2/22/24		
Client Name:	RSI EnTech, LLC								
Address:	203 Victorious Blvd.	., Oak Ridge, TN	I 37830						
Sample ID:	BRC0091-03	Type:	REG		Sample D	ate:	12/28/2023		
					De	oth:	NP		



Cobbles	< 300 mm (12") and > 75 mm	Fine Sand	< 0.425 mm and > 0.075 mm
Gravel	< 75 mm and > 4.75 mm (#4)	Silt and Clav	< 0.075 mm
Coarse Sand	< 4.75 mm and >2.00 mm	Clav	<0.002 mm
Medium Sand	< 2.00 mm and > 0.425 mm		

Wicaiaiii Sana C.	<del>JO IIIIII GIIG</del>	<u> </u>				_
Maximum Particle Size:	#4	Gravel:	0%	Medium Sand:	6%	
Silt & Clay (% Passing #200):	79%	Total Sand:	21%	Fine Sand:	15%	
Assumed Specific Gravity:	2.65	Coarse Sand:	0%	Clay:	15.1%	
Liquid Limit	TNP	Plastic Limit	TNP	Plastic Index	TNP	

References / Comments / Deviations:

<u>Tyler Copeman</u> Technical Responsibility Signature

<u>Lab Services Manager</u> *Position*  2/22/2024 Date


Revision No. 0


Revision Date: 03/18/2019

ASTM D7928 & D6913









Cobbles	< 300 mm (12") and > 75 mm	Fine Sand	< 0.425 mm and > 0.075 mm
Gravel	< 75 mm and > 4.75 mm (#4)	Silt and Clav	< 0.075 mm
Coarse Sand	< 4.75 mm and >2.00 mm	Clav	<0.002 mm
Medium Sand	< 2.00 mm and > 0.425 mm		

Maximum Particle Size:	3/4"	Gravel:	2%	Medium Sand:	4%
Silt & Clay (% Passing #200):	74%	Total Sand:	24%	Fine Sand:	18%
Assumed Specific Gravity:	2.65	Coarse Sand:	2%	Clay:	13.4%
Liquid Limit	TNP	Plastic Limit	TNP	Plastic Index	TNP

References / Comments / Deviations:

Tyler Copeman Technical Responsibility

Lab Services Manager Position

2/22/2024 Date

Form No. TR-D7928-3 Revision No. 0


Revision Date: 03/18/2019

**HCTREF-FP** 



Log No.: 43-4016 ASTM D7928 & D6913

	S&ME, Inc., 1413 Topside Road, Louisville, TN 37777							
S&ME Project #:	ያME Project #: 23430142						3/18/2024	
Project Name:	UCOR-Geotech				Test Start Date:	3/1	2/24-3/14/24	
Client Name:	RSI EnTech, LLC							
Address:	203 Victorious Blvd.,	Oak Ridge, TN	37830		•			
Sample ID:	BCR0098-04	Type:	REG		Sample D	ate:	2/2/2024	
					De	pth:	NP	



Medium Sand: Maximum Particle Size: 3/4" Gravel: 6% 13% Silt & Clay (% Passing #200): 56% Total Sand: 38% Fine Sand: 24% Assumed Specific Gravity: 2.65 Coarse Sand: 1% Clay: 7.9% Liquid Limit **TNP** Plastic Limit **TNP** Plastic Index TNP

References / Comments / Deviations:

<u>Tyler Copeman</u> Technical Responsibility Signature

<u>Lab Services Manager</u> *Position*  3/18/2024 Date

This page intentionally left blank.

**UPPER CREEK BANK SOIL** 

This page intentionally left blank.

Form No. TR-D7928-3 Revision No. 0

Log No.:

Revision Date: 03/18/2019

43-4016

**BCT1-BSU** 



ASTM D7928 & D6913

	S&ME, Inc., 1413 Topside Road, Louisville, TN 37777								
S&ME Project #:	23430142			Report	Date:	2/23/2024			
Project Name:	UCOR-Geotech			Test Star	t Date:	2/1/2024			
Client Name:	RSI EnTech, LLC								
Address:	203 Victorious Blvd	., Oak Ridge, TN	37830						
Sample ID:	BCR0038-03	Type:	REG	S	ample Date:	1/4/2024			
					Denth:	NP			

Sample Description: Dark Brown Clay with Sand #60 #100 #200 3/8" 100% 90% 80% 70% Percent Passing 60% 50% 40% 30% 20% 10% 0% 10 0.01 0.001 100 Particle Size (mm) 300 mm (12") and > 75 mm Cobbles Fine Sand < 0.425 mm and > 0.075 mm Gravel < 75 mm and > 4.75 mm (#4) Silt and Clav < 0.075 mm Coarse Sand < 4.75 mm and > 2.00 mm Clav <0.002 mm Medium Sand < 2.00 mm and > 0.425 mm Maximum Particle Size: #20 Gravel: 0% Medium Sand: 1% Silt & Clay (% Passing #200): 76% Total Sand: 24% Fine Sand: 23%

References / Comments / Deviations:

Assumed Specific Gravity:

Liquid Limit

<u>Victoria Igoe</u> Technical Responsibility

Signature

2.65

**TNP** 

Senior Engineering Technician

Position

0%

TNP

2/23/2024 Date

9.4%

**TNP** 

Clay:

Plastic Index

This report shall not be reproduced, except in full, without the written approval of S&ME, Inc.

Coarse Sand:

Plastic Limit

Revision No. 0


Revision Date: 03/18/2019

BCT2-BSU



Log No.: 43-4016 ASTM D7928 & D6913

	S&ME, Inc., 1413 Topside Road, Louisville, TN 37777								
S&ME Project #:	23430142			Report Date:	2/16/20	024			
Project Name:	UCOR-Geotech	UCOR-Geotech			Test Start Date: 2/1/2024				
Client Name:	RSI EnTech, LLC								
Address:	203 Victorious Blvd.	, Oak Ridge, TN	37830						
Sample ID:	BCR0041-03	Туре:	REG		Sample Dat	:e: 1/4/	2024		
				·	Dept	h: N	IP		



Cobbles	< 300 mm (12") and > 75 mm	Fine Sand	< 0.425 mm and > 0.075 mm
Gravel	< 75 mm and > 4.75 mm (#4)	Silt and Clav	< 0.075 mm
Coarse Sand	< 4.75 mm and >2.00 mm	Clav	<0.002 mm
Medium Sand	< 2.00 mm and > 0.425 mm		

Wicalaili Salia V	<u> </u>	9 0. <del>4</del> 23 111111				_
Maximum Particle Size:	#4	Gravel:	0%	Medium Sand:	2%	
Silt & Clay (% Passing #200):	84%	Total Sand:	16%	Fine Sand:	14%	
Assumed Specific Gravity:	2.65	Coarse Sand:	0%	Clay:	13.7%	
Liquid Limit	TNP	Plastic Limit	TNP	Plastic Index	TNP	

References / Comments / Deviations:

<u>Tyler Copeman</u> Technical Responsibility Signature

<u>Lab Services Manager</u> *Position*  2/19/2024 Date

Revision No. 0


Revision Date: 03/18/2019

**BCT3-BSU** 



Log No.: 43-4016 ASTM D7928 & D6913

	S&ME, Inc	., 1413 Topsid	e Road, Loui	sville, 1	ΓN 37777		
S&ME Project #:	23430142		Report Date:	2/19/20	024		
Project Name:	UCOR-Geotech	UCOR-Geotech				2/7/20	24
Client Name:	RSI EnTech, LLC						
Address:	203 Victorious Blvd.	, Oak Ridge, TN	I 37830				
Sample ID:	BCR0044-03	Type:	REG		Sample Dat	:e: 1/8/	2024
					Dept	h: N	1P



Cobbles	< 300 mm (12") and > 75 mm	Fine Sand	< 0.425 mm and > 0.075 mm
Gravel	< 75 mm and > 4.75 mm (#4)	Silt and Clav	< 0.075 mm
Coarse Sand	< 4.75 mm and >2.00 mm	Clav	<0.002 mm
Medium Sand	< 2.00 mm and > 0.425 mm		

Maximum Particle Size:	3/8"	Gravel:	6%	Medium Sand:	6%	
Silt & Clay (% Passing #200):	82%	Total Sand:	12%	Fine Sand:	3%	
Assumed Specific Gravity:	2.65	Coarse Sand:	3%	Clay:	13.9%	
Liquid Limit	TNP	Plastic Limit	TNP	Plastic Index	TNP	

References / Comments / Deviations:

<u>Tyler Copeman</u> Technical Responsibility Signature

<u>Lab Services Manager</u> *Position*  2/19/2024 Date

Form No. TR-D7928-3

Revision No. 0

Log No.:

Revision Date: 03/18/2019


43-4016

BCT4-BSU



ASTM D7928 & D6913

	S&ME, Inc.	, 1413 Topsid	e Road, Lou	iisville, <sup>-</sup>	TN 37777	
S&ME Project #:	23430142		Report Date:	1/0/1900		
Project Name:	UCOR-Geotech				Test Start Date:	2/7/2024
Client Name:	Client Name: RSI EnTech, LLC					
Address:	203 Victorious Blvd.,	Oak Ridge, TN	l 37830			
Sample ID:	le ID: BCR0047-03 Type: REG				Sample Date	e: 1/8/2024
		Depth	n: NP			



Cobbles	< 300 mm (12") and > 75 mm	Fine Sand	< 0.425 mm and > 0.075 mm
Gravel	< 75 mm and > 4.75 mm (#4)	Silt and Clav	< 0.075 mm
Coarse Sand	< 4.75 mm and >2.00 mm	Clav	<0.002 mm
Medium Sand	< 2.00 mm and > 0.425 mm		

Tricalatii Salla	<u>00                                   </u>	O. ILS IIIII				
Maximum Particle Size:	#10	Gravel:	0%	Medium Sand:	4%	
Silt & Clay (% Passing #200):	62%	Total Sand:	38%	Fine Sand:	34%	
Assumed Specific Gravity:	2.65	Coarse Sand:	0%	Clay:	8.4%	
Liquid Limit	TNP	Plastic Limit	TNP	Plastic Index	TNP	

References / Comments / Deviations:

<u>Tyler Copeman</u> Technical Responsibility Signature

<u>Lab Services Manager</u> *Position*  2/19/2024 Date

Revision No. 0

Revision Date: 03/18/2019


Form No. TR-D7928-3

**BCT5-BSU** 



Log No.: 43-4016 ASTM D7928 & D6913

	S&ME, Inc., 1413 Topside Road, Louisville, TN 37777									
S&ME Project #:	23430142	23430142			Report Date:		3/12/2024			
Project Name:	UCOR-Geotech				Test Start Date:	3/7	7/24-3/11/24			
Client Name:	RSI EnTech, LLC									
Address:	203 Victorious Blvd.	, Oak Ridge, TN	I 37830							
Sample ID:	BCR0050-03	Туре:	REG		Sample Date:		2/1/2024			
					Dep	oth:	NP			



References / Comments / Deviations:

Silt & Clay (% Passing #200):

Assumed Specific Gravity:

Liquid Limit

<u>Tyler Copeman</u> Technical Responsibility Signature

52%

2.65

TNP

Lab Services Manager
Position

46%

3%

**TNP** 

3/12/2024 Date

36%

11.7%

TNP

This report shall not be reproduced, except in full, without the written approval of S&ME, Inc.

Total Sand:

Coarse Sand:

Plastic Limit

Fine Sand:

Plastic Index

Clay:

Form No. TR-D7928-3 Revision No. 0


Revision Date: 03/18/2019

**BCT6-BSU** 



Log No.: 43-4016 ASTM D7928 & D6913

S&ME, Inc., 1413 Topside Road, Louisville, TN 37777									
S&ME Project #:	ME Project #: 23430142						3/14/2024		
Project Name:	UCOR-Geotech				Test Start Date:	3/1	0/24-3/13/24		
Client Name:	RSI EnTech, LLC								
Address:	203 Victorious Blvd.	, Oak Ridge, TN	I 37830						
Sample ID:	BCR0054-03	Туре:	REG		Sample Date:		2/1/2024		
					De	pth:	NP		



References / Comments / Deviations:

Assumed Specific Gravity:

Liquid Limit

<u>Tyler Copeman</u> Technical Responsibility Signature

2.65

TNP

<u>Lab Services Manager</u> <u>Position</u>

2%

**TNP** 

3/14/2024 Date

10.9%

TNP

This report shall not be reproduced, except in full, without the written approval of S&ME, Inc.

Coarse Sand:

Plastic Limit

Clay:

Plastic Index

Form No. TR-D7928-3 Revision No. 0

Revision Date: 03/18/2019


BCT7-BS



Log No.: 43-4016 ASTM D7928 & D6913

S&ME, Inc., 1413 Topside Road, Louisville, TN 37777									
S&ME Project #:	Report Date:	3/14/2024							
Project Name:	UCOR-Geotech				Test Start Date:	3/1	0/24-3/13/24		
Client Name:	RSI EnTech, LLC								
Address:	203 Victorious Blvd.,	Oak Ridge, TN	I 37830						
Sample ID:	BCR0057-03	Туре:	REG		Sample D	ate:	2/7/2024		
					De	pth:	NP		

Sample Description: **Brown Sandy Silt** 1" 3/4" 3/8" #4 #20 #40 #60 #100 #200 100%



Cobbles	< 300 mm (12") and > 75 mm	Fine Sand	< 0.425 mm and > 0.075 mm
Gravel	< 75 mm and > 4.75 mm (#4)	Silt and Clav	< 0.075 mm
Coarse Sand	< 4.75 mm and >2.00 mm	Clav	<0.002 mm
Medium Sand	< 2.00 mm and > 0.425 mm		

Maximum Particle Size:	3/4''	Gravel:	1%	Medium Sand:	10%
Silt & Clay (% Passing #200):	57%	Total Sand:	42%	Fine Sand:	32%
Assumed Specific Gravity:	2.65	Coarse Sand:	0%	Clay:	10.9%
Liquid Limit	TNP	Plastic Limit	TNP	Plastic Index	TNP

References / Comments / Deviations:

Tyler Copeman Technical Responsibility

Lab Services Manager Position

3/14/2024 Date

Form No. TR-D7928-3 Revision No. 0


Revision Date: 03/18/2019

BCT8-BS



Log No.: 43-4016 ASTM D7928 & D6913

	S&ME, Inc., 1413 Topside Road, Louisville, TN 37777									
S&ME Project #:	S&ME Project #: 23430142						3/18/2024			
Project Name:	UCOR-Geotech	OR-Geotech			Test Start Date:	3/1	2/24-3/14/24			
Client Name:	RSI EnTech, LLC									
Address:	203 Victorious Blvd.,	Oak Ridge, TN	37830							
Sample ID:	BCR0079-03	Type:	REG		Sample D	ate:	2/5/2024			
					De	pth:	NP			



References / Comments / Deviations:

Silt & Clay (% Passing #200):

Assumed Specific Gravity:

Liquid Limit

<u>Tyler Copeman</u> Technical Responsibility Signature

38%

2.65

**TNP** 

<u>Lab Services Manager</u> *Position* 

49%

10%

TNP

3/18/2024 Date

25%

6.5%

TNP

This report shall not be reproduced, except in full, without the written approval of S&ME, Inc.

Total Sand:

Coarse Sand:

Plastic Limit

Fine Sand:

Plastic Index

Clay:

Revision No. 0

Form No. TR-D7928-3


Revision Date: 03/18/2019

BCT9-BS



Log No.: 43-4016 ASTM D7928 & D6913

3									
S&ME, Inc., 1413 Topside Road, Louisville, TN 37777									
S&ME Project #:	23430142		Report Date:			3	3/15/2024		
Project Name:	UCOR-Geotech				Test Start Date:	3/10	0/24-3/13/24		
Client Name:	RSI EnTech, LLC								
Address:	203 Victorious Blvd.	, Oak Ridge, TN	l 37830						
Sample ID:	BCR0068-03	Type:	REG		Sample Date:		2/5/2024		
					De	pth:	NP		



*References / Comments / Deviations:* 

Assumed Specific Gravity:

Liquid Limit

<u>Tyler Copeman</u> Technical Responsibility Signature

2.65

TNP

<u>Lab Services Manager</u> <u>Position</u>

4%

**TNP** 

3/15/2024 Date

12.2%

TNP

This report shall not be reproduced, except in full, without the written approval of S&ME, Inc.

Coarse Sand:

Plastic Limit

Clay:

Plastic Index

Form No. TR-D7928-3

Revision No. 0

Log No.:

Revision Date: 03/18/2019

43-4016

BCT10-BSU



ASTM D7928 & D6913

S&ME, Inc., 1413 Topside Road, Louisville, TN 37777							
S&ME Project #:	23430142	23430142			Report Date: 2		2/20/2024
Project Name:	UCOR-Geotech				Test Start Date:	2/1	2/24-2/16/24
Client Name:	RSI EnTech, LLC						
Address:	203 Victorious Blvd.	, Oak Ridge, TN	I 37830				
Sample ID:	BRC0073-03	Type:	REG		Sample D	ate:	1/11/2024
					De	oth:	NP

Sample Description: Brown Clay with Sand 1" 3/4" #40 #60 #100 #200 100% 90% 80% 70% Percent Passing 60% 50% 40% 30% 20% 10% 0% 0.1 0.01 0.001 Particle Size (mm)

Cobbles	< 300 mm (12") and > 75 mm	Fine Sand	< 0.425 mm and > 0.075 mm
Gravel	< 75 mm and > 4.75 mm (#4)	Silt and Clav	< 0.075 mm
Coarse Sand	< 4.75 mm and >2.00 mm	Clav	<0.002 mm
Medium Sand	< 2.00 mm and > 0.425 mm		

Wicalain Sana C.C	<del>//                                     </del>	2 0.423 HIIII				_
Maximum Particle Size:	3/8''	Gravel:	1%	Medium Sand:	1%	
Silt & Clay (% Passing #200):	77%	Total Sand:	22%	Fine Sand:	21%	
Assumed Specific Gravity:	2.65	Coarse Sand:	0%	Clay:	5.6%	
Liquid Limit	TNP	Plastic Limit	TNP	Plastic Index	TNP	

References / Comments / Deviations:

<u>Tyler Copeman</u> Technical Responsibility Signature

<u>Lab Services Manager</u> *Position*  2/20/2024 Date

Form No. TR-D7928-3

Revision No. 0

Revision Date: 03/18/2019 Log No.: 43-4016


BCT11-BSU



ASTM D7928 & D6913

S&ME, Inc., 1413 Topside Road, Louisville, TN 37777							
S&ME Project #:	&ME Project #: 23430142						3/18/2024
Project Name:	UCOR-Geotech				Test Start Date:	3/1	2/24-3/14/24
Client Name:	RSI EnTech, LLC						
Address:	203 Victorious Blvd.,	Oak Ridge, TN	37830				
Sample ID:	BCR0076-03	Type:	REG		Sample Date: 1/3		1/31/2024
					De	nth:	NP

NΡ



	Copples	South the state of the state		e sanu	\( \text{0.423 } \text{11111 } \text{and } \( \text{2.073 } \text{11111} \)		
Gravel < 75 mm and > 4.75 mm (#4)			Silt and Clav		< 0.075 mm		
	Coarse Sand	Clav		<0.002 mm			
	Medium Sand	< 2.00 mm and > 0.425 mm					
	Maximum Particle S	Size: 3/8"	Gravel:	0%	Medium Sand:	2%	

THAMITTATTT AT CICIC DIEC.	3,0	Grave	0,0	mealani bana.	-70
Silt & Clay (% Passing #200):	77%	Total Sand:	23%	Fine Sand:	20%
Assumed Specific Gravity:	2.65	Coarse Sand:	1%	Clay:	12.1%
Liquid Limit	TNP	Plastic Limit	TNP	Plastic Index	TNP

References / Comments / Deviations:

Tyler Copeman Technical Responsibility

Lab Services Manager Position

3/18/2024 Date

Revision No. 0

Log No.:

Revision Date: 03/18/2019


43-4016

BCT12B-BSU



#### ASTM D7928 & D6913

	S&ME, Inc., 1413 Topside Road, Louisville, TN 37777							
S&ME Project #:	23430142	30142 Report Date:				2/19/2024		
Project Name:	UCOR-Geotech			Test Start Date:	2/1	2/24-2/16/24		
Client Name:	RSI EnTech, LLC							
Address:	203 Victorious Blvd.	, Oak Ridge, TN 37830						
Sample ID:	BCR0063-03	Type: REG		Sample D	ate:	1/10/2024		
				Do	nth:	NID		



Coarse Sand	< 4.75 mm and >	mm and >2.00 mm (#10) Clav		<0.002 m	m	
Medium Sand	< 2.00 mm and	> 0.425 mm				
Maximum Particle Si	ize: 3/8"		Gravel:	4%	Medium Sand:	4%
Silt & Clay (% Passing #2	00): 83%	To	otal Sand:	13%	Fine Sand:	7%
Assumed Specific Grav	vity: 2.65	Coa	rse Sand:	2%	Clay:	14.1%
Liquid Li	imit TNP	Pla	stic Limit	TNP	Plastic Index	TNP

**References / Comments / Deviations:** 

Tyler Copeman
Technical Responsibility

Signature

<u>Lab Services Manager</u> *Position*  2/19/2024 Date


Revision No. 0

Revision Date: 03/18/2019

# BCT12B-BSU



Log No.:	43-4016						
		S&ME, Inc	lle, TN 37777				
S&ME Proje	ect #:	23430142			Report Date:	;	2/19/2024
Project Nam	ne:	UCOR-Geotech			Test Start Date:	2/1	2/24-2/16/24
Client Name	e:	RSI EnTech, LLC					
Address: 203 Victorious Blvd., Oak Ridge, TN 37830		37830					
Sample ID:		BCR0064-03	Type:	REG	Sample D	ate:	1/10/2024
					De	pth:	NP



Gravel	< /5	<u>mm and &gt; 4.7</u>	/5 mm (#4)	#4) Silt and Clay		< 0.075 m	< 0.075 mm		
Coarse Sand	< 4.7	5 mm and >2.0	00 mm (#10)		Clav	<0.002 mr	n		
Medium Sand	< 2	.00 mm and >	0.425 mm						
Maximum Particle S	ize:	3/4''		Gravel:	5%	Medium Sand:	5%		
Silt & Clay (% Passing #2	200):	72%	To	tal Sand:	23%	Fine Sand:	16%		
Assumed Specific Gra	vity:	2.65	Coa	rse Sand:	2%	Clay:	12.8%		
Liquid L	imit	TNP	Pla	stic Limit	TNP	Plastic Index	TNP		

**References / Comments / Deviations:** 

Tyler Copeman Technical Responsibility

Lab Services Manager Position

2/19/2024 Date

Form No. TR-D7928-3

Revision No. 0

Revision Date: 03/18/2019

BCT12A-BSU



Log No.: 43-4016 ASTM D7928 & D6913

	S&ME, Inc., 1413 Topside Road, Louisville, TN 37777								
S&ME Project #:	23430142				Report Date:		2/22/2024		
Project Name:	UCOR-Geotech	UCOR-Geotech			Test Start Date:	2/2	0/24-2/22/24		
Client Name:	RSI EnTech, LLC								
Address:	203 Victorious Blvd.	, Oak Ridge, TN	N 37830						
Sample ID:	BRC0105-03	Type:	REG		Sample D	ate:	1/10/2024		
					De <sub>l</sub>	oth:	NP		

Sample Description: Brown Clay with Sand 1" 3/4" #40 #60 #100 #200 100% 90% 80% 70% Percent Passing 60% 50% 40% 30% 20% 10% 0% 0.1 0.01 0.001 Particle Size (mm)

Cobbles	< 300 mm (12") and > 75 mm	Fine Sand	< 0.425 mm and > 0.075 mm		
Gravel	< 75 mm and > 4.75 mm (#4)	Silt and Clav	< 0.075 mm		
Coarse Sand	< 4.75 mm and >2.00 mm	Clav	<0.002 mm		
Medium Sand	< 2.00 mm and > 0.425 mm				

IVICAIAITI SAITA \ L.	<del>oo iiiiii ana</del>	> 0. <del>1</del> 23 11111				
Maximum Particle Size:	3/4"	Gravel:	5%	Medium Sand:	3%	
Silt & Clay (% Passing #200):	80%	Total Sand:	15%	Fine Sand:	10%	
Assumed Specific Gravity:	2.65	Coarse Sand:	2%	Clay:	14.5%	
Liquid Limit	TNP	Plastic Limit	TNP	Plastic Index	TNP	

References / Comments / Deviations:

<u>Tyler Copeman</u> Technical Responsibility Signature

<u>Lab Services Manager</u> *Position*  2/22/2024 Date

Revision No. 0

Log No.:

Revision Date: 03/18/2019


43-4016

BCT13-BSU



ASTM D7928 & D6913

	S&ME, Inc., 1413 Topside Road, Louisville, TN 37777						
S&ME Project #:	23430142				Report Date:		2/20/2024
Project Name:	UCOR-Geotech				Test Start Date:	2/1	2/24-2/16/24
Client Name:	RSI EnTech, LLC						
Address:	203 Victorious Blvd.	, Oak Ridge, TN	N 37830				
Sample ID:	BRC0084-03	Type:	REG		Sample D	ate:	1/11/2024
					De	oth:	NP



Cobbles	< 300 mm (12") and > 75 mm	Fine Sand	< 0.425 mm and > 0.075 mm
Gravel	< 75 mm and > 4.75 mm (#4)	Silt and Clav	< 0.075 mm
Coarse Sand	< 4.75 mm and >2.00 mm	Clav	<0.002 mm
Medium Sand	< 2.00 mm and > 0.425 mm		

Maximum Particle Size:	3/8"	Gravel:	2%	Medium Sand:	6%
Silt & Clay (% Passing #200):	68%	Total Sand:	30%	Fine Sand:	23%
Assumed Specific Gravity:	2.65	Coarse Sand:	1%	Clay:	15.2%
Liquid Limit	TNP	Plastic Limit	TNP	Plastic Index	TNP

References / Comments / Deviations:

<u>Tyler Copeman</u> Technical Responsibility Signature

<u>Lab Services Manager</u> *Position*  2/20/2024 Date

Form No. TR-D7928-3

Revision No. 0

Log No.:

Revision Date: 03/18/2019


43-4016

BCT14-BSU



ASTM D7928 & D6913

	S&ME, Inc., 1413 Topside Road, Louisville, TN 37777							
S&ME Project #:	23430142				Report Date:	á	2/20/2024	
Project Name:	UCOR-Geotech				Test Start Date:	2/1	7/24-2/20/24	
Client Name:	RSI EnTech, LLC							
Address:	203 Victorious Blvd	., Oak Ridge, TN	I 37830					
Sample ID:	BRC0088-03	Type:	REG		Sample D	ate:	1/2/2024	
					De	oth:	NP	



Cobbles	< 300 mm (12") and > 75 mm	Fine Sand	< 0.425 mm and > 0.075 mm
Gravel	< 75 mm and > 4.75 mm (#4)	Silt and Clav	< 0.075 mm
Coarse Sand	< 4.75 mm and >2.00 mm	Clav	<0.002 mm
Medium Sand	< 2.00 mm and > 0.425 mm		

Maximum Particle Size:	#10	Gravel:	0%	Medium Sand:	5%	
Silt & Clay (% Passing #200):	86%	Total Sand:	14%	Fine Sand:	9%	
Assumed Specific Gravity:	2.65	Coarse Sand:	0%	Clay:	23.1%	
Liquid Limit	TNP	Plastic Limit	TNP	Plastic Index	TNP	

References / Comments / Deviations:

<u>Tyler Copeman</u> Technical Responsibility Signature

<u>Lab Services Manager</u> *Position*  2/20/2024 Date

Revision No. 0


Revision Date: 03/18/2019

BCT14-BSU



Log No.: 43-4016 ASTM D7928 & D6913

	S&ME, Inc., 1413 Topside Road, Louisville, TN 37777							
S&ME Project #:	23430142				Report Date:	ï	2/21/2024	
Project Name:	UCOR-Geotech				Test Start Date:	2/1	2/24-2/16/24	
Client Name:	RSI EnTech, LLC							
Address:	203 Victorious Blvd.	, Oak Ridge, TN	I 37830					
Sample ID:	BRC0089-03	Type:	REG		Sample D	ate:	1/2/2024	
					De	pth:	NP	



Cobbles	< 300 mm (12") and > 75 mm	Fine Sand	< 0.425 mm and > 0.075 mm
Gravel	< 75 mm and > 4.75 mm (#4)	Silt and Clav	< 0.075 mm
Coarse Sand	< 4.75 mm and >2.00 mm	Clav	<0.002 mm
Medium Sand	< 2.00 mm and > 0.425 mm		

Maximum Particle Size:	#10	Gravel:	0%	Medium Sand:	5%
Silt & Clay (% Passing #200):	86%	Total Sand:	14%	Fine Sand:	9%
Assumed Specific Gravity:	2.65	Coarse Sand:	0%	Clay:	22.2%
Liquid Limit	TNP	Plastic Limit	TNP	Plastic Index	TNP

**References / Comments / Deviations:** 

<u>Tyler Copeman</u> Technical Responsibility Signature

<u>Lab Services Manager</u> *Position*  2/21/2024 Date

Form No. TR-D7928-3

Revision No. 0

Log No.:

Revision Date: 03/18/2019


43-4016

BCT15-BSU



ASTM D7928 & D6913

S&ME, Inc., 1413 Topside Road, Louisville, TN 37777							
S&ME Project #:	23430142				Report Date:	2	/21/2024
Project Name:	UCOR-Geotech				Test Start Date:	2/17	//24-2/20/24
Client Name:	RSI EnTech, LLC						
Address:	203 Victorious Blvd.	, Oak Ridge, TN	I 37830				
Sample ID:	BRC0094-03	Type:	REG		Sample D	ate:	NP
					De	pth:	NP



Cobbles	< 300 mm (12") and > 75 mm	Fine Sand	< 0.425 mm and > 0.075 mm
Gravel	< 75 mm and > 4.75 mm (#4)	Silt and Clav	< 0.075 mm
Coarse Sand	< 4.75 mm and >2.00 mm	Clav	<0.002 mm
Medium Sand	< 2.00 mm and > 0.425 mm		

Wicaiaiii Sana 🔍 Z	<u> /</u>	0.423 IIIII				
Maximum Particle Size:	#4	Gravel:	0%	Medium Sand:	3%	
Silt & Clay (% Passing #200):	77%	Total Sand:	23%	Fine Sand:	19%	
Assumed Specific Gravity:	2.65	Coarse Sand:	1%	Clay:	12.5%	
Liquid Limit	TNP	Plastic Limit	TNP	Plastic Index	TNP	

References / Comments / Deviations:

<u>Tyler Copeman</u> Technical Responsibility Signature

<u>Lab Services Manager</u> *Position*  2/21/2024 Date

Form No. TR-D7928-3

Revision No. 0

Revision Date: 03/18/2019 Log No.: 43-4016 **HCTREF-BSU** 



ASTM D7928 & D6913

S&ME, Inc., 1413 Topside Road, Louisville, TN 37777							
S&ME Project #:	23430142		Report Date:		3/18/2024		
Project Name:	UCOR-Geotech	UCOR-Geotech		Test Start Date:	3/12/24-3/14/24		
Client Name:	RSI EnTech, LLC						
Address:	203 Victorious Blvd.,	, Oak Ridge, TN	I 37830				
Sample ID:	BCR0097-03	Туре:	REG		Sample Date: 2/2		2/2/2024
					De	NP	

Depth: NΡ Sample Description: **Brown Sandy Silt** 1" 3/4" 3/8" #4 #20 #40 #60 #100 #200 100% 90% 80% 70% Percent Passing 60% 50% 40% 30% 20% 10% 0% 10 0.1 1 0.01 0.001 100 Particle Size (mm) < 0.425 mm and > 0.075 mm Cobbles < 300 mm (12") and > 75 mm Fine Sand Gravel < 75 mm and > 4.75 mm (#4) Silt and Clav < 0.075 mm < 4.75 mm and >2.00 mm <0.002 mm Coarse Sand Clav Medium Sand < 2.00 mm and > 0.425 mm

Maximum Particle Size:	3/8''	Gravel:	1%	Medium Sand:	6%
Silt & Clay (% Passing #200):	55%	Total Sand:	44%	Fine Sand:	36%
Assumed Specific Gravity:	2.65	Coarse Sand:	2%	Clay:	9.7%
Liquid Limit	TNP	Plastic Limit	TNP	Plastic Index	TNP

References / Comments / Deviations:

<u>Tyler Copeman</u> Technical Responsibility Signature

<u>Lab Services Manager</u> *Position*  3/18/2024 Date

This page intentionally left blank.

LOWER CREEK BANK SOIL

This page intentionally left blank.

Form No. TR-D7928-3 Revision No. 0

Log No.:

Revision Date: 03/18/2019


43-4016

**BCT1-BSL** 



ASTM D7928 & D6913

S&ME, Inc., 1413 Topside Road, Louisville, TN 37777							
S&ME Project #:	23430142				Report Date:		2/23/2024
Project Name:	UCOR-Geotech				Test Start Date:		2/1/2024
Client Name:	RSI EnTech, LLC						
Address:	203 Victorious Blvd.	, Oak Ridge, TN	N 37830				
Sample ID:	BRC0037-03	Type:	REG		Sample Date: 1/4/2024		1/4/2024
·			·		Dept	h:	NP



Silt & Clay (% Passing #200): 75% Total Sand: 24% Fine Sand: 17% Assumed Specific Gravity: Coarse Sand: 10.0% 2.65 1% Clay: Liquid Limit **TNP** Plastic Limit TNP Plastic Index **TNP** 

References / Comments / Deviations:

<u>Victoria Igoe</u> Technical Responsibility

Signature

Senior Engineering Technician

Position

2/23/2024 Date

Form No. TR-D7928-3

Revision No. 0

Log No.:

Revision Date: 03/18/2019


43-4016

**BCT2-BSL** 



ASTM D7928 & D6913

	S&ME, Inc., 1413 Topside Road, Louisville, TN 37777								
S&ME Project #:	23430142				Report Date:	2/16/2024			
Project Name:	UCOR-Geotech				Test Start Date:	2/1/2024			
Client Name:	RSI EnTech, LLC								
Address:	203 Victorious Blvd.	, Oak Ridge, TN	l 37830						
Sample ID:	BCR0040-03	Type:	REG		Sample Date: 1/4/2024				
					Dept	h: NP			



Cobbles	< 300 mm (12") and > 75 mm	Fine Sand	< 0.425 mm and > 0.075 mm
Gravel	< 75 mm and > 4.75 mm (#4)	Silt and Clav	< 0.075 mm
Coarse Sand	< 4.75 mm and >2.00 mm	Clav	<0.002 mm
Medium Sand	< 2.00  mm and  > 0.425  mm		

TVICAIAITI SAITA	<del>70 111111 ana</del>	/ U.TES IIIII				_
Maximum Particle Size:	#4	Gravel:	4%	Medium Sand:	11%	
Silt & Clay (% Passing #200):	63%	Total Sand:	33%	Fine Sand:	19%	
Assumed Specific Gravity:	2.65	Coarse Sand:	3%	Clay:	12.5%	
Liquid Limit	TNP	Plastic Limit	TNP	Plastic Index	TNP	

**References / Comments / Deviations:** 

<u>Tyler Copeman</u> Technical Responsibility Signature

<u>Lab Services Manager</u> *Position*  2/19/2024 Date

Form No. TR-D7928-3 Revision No. 0


Revision Date: 03/18/2019

Log No.: 43-4016 ASTM D7928 & D6913

**BCT3-BSL** 



9		,		_				
	S&ME, Inc., 1413 Topside Road, Louisville, TN 37777							
S&ME Project #:	23430142				Report Date:	2/19/2024		
Project Name:	UCOR-Geotech				Test Start Date:	2/7/2024		
Client Name:	RSI EnTech, LLC							
Address:	203 Victorious Blvd.	, Oak Ridge, TN	I 37830					
Sample ID:	BCR0043-03	Type:	REG		Sample Dat	e: 1/4/202	24	
					Dept	h: NP		



Cobbles	< 300 mm (12") and > 75 mm	Fine Sand	< 0.425 mm and > 0.075 mm
Gravel	< 75 mm and > 4.75 mm (#4)	Silt and Clav	< 0.075 mm
Coarse Sand	< 4.75 mm and >2.00 mm	Clav	<0.002 mm
Medium Sand	< 2.00 mm and > 0.425 mm		

Wicalam Sana 2.0	<del>Jo IIIIII and</del>	/ U.TLJ IIIII			
Maximum Particle Size:	#10	Gravel:	0%	Medium Sand:	2%
Silt & Clay (% Passing #200):	91%	Total Sand:	9%	Fine Sand:	7%
Assumed Specific Gravity:	2.65	Coarse Sand:	0%	Clay:	12.4%
Liquid Limit	TNP	Plastic Limit	TNP	Plastic Index	TNP

References / Comments / Deviations:

<u>Tyler Copeman</u> Technical Responsibility Signature

<u>Lab Services Manager</u> *Position*  2/19/2024 Date

Form No. TR-D7928-3


Revision No. 0

Revision Date: 03/18/2019





Log No.:	43-4016		ASTM D7	928 & D691	3		
S&ME, Inc., 1413 Topside Road, Louisville, TN 37777							
S&ME Proje	ect #:	23430142				Report Date:	2/16/2024
Project Nam	ne:	UCOR-Geotech	-			Test Start Date:	2/7/2024
Client Name	e:	RSI EnTech, LLC					
Address:		203 Victorious Blvd	., Oak Ridge, TN	N 37830			
Sample ID:		BCR0046-03	Type:	REG		Sample Date:	1/8/2024
	·					Depth	. NP



Cobbles	< 300 mm (12") and > 75 mm	Fine Sand	< 0.425 mm and > 0.075 mm
Gravel	< 75 mm and > 4.75 mm (#4)	Silt and Clav	< 0.075 mm
Coarse Sand	< 4.75 mm and >2.00 mm	Clav	<0.002 mm
Medium Sand	< 2.00 mm and > 0.425 mm		

Maximum Particle Size:	#4	Gravel:	0%	Medium Sand:	5%
Silt & Clay (% Passing #200):	71%	Total Sand:	29%	Fine Sand:	24%
Assumed Specific Gravity:	2.65	Coarse Sand:	0%	Clay:	11.1%
Liquid Limit	TNP	Plastic Limit	TNP	Plastic Index	TNP

References / Comments / Deviations:

Tyler Copeman Technical Responsibility

Lab Services Manager Position

2/16/2024 Date

Form No. TR-D7928-3 Revision No. 0

Revision Date: 03/18/2019

**BCT5-BSL** 



Log No.: 43-4016 ASTM D7928 & D6913

	S&ME, Inc., 1413 Topside Road, Louisville, TN 37777								
S&ME Project #:	23430142				Report Date:		3/12/2024		
Project Name:	UCOR-Geotech				Test Start Date:	3/7	7/24-3/11/24		
Client Name:	RSI EnTech, LLC								
Address:	203 Victorious Blvd.	, Oak Ridge, TN	I 37830						
Sample ID:	BCR0049-03	Туре:	REG		Sample Date: 2/1/2024		2/1/2024		
					Dep	oth:	NP		

Brown Clay with Sand Sample Description: 1" 3/4" 3/8" #4 #20 #40 #60 #100 #200 100% 90% 80% 70% Percent Passing 60% 50% 40% 30% 20% 10% 0% 10 1 0.1 0.01 0.001 100 Particle Size (mm) < 300 mm (12") and > 75 mm < 0.425 mm and > 0.075 mm Cohbles Fine Sand

CODDICS	<u> </u>		- Janu	<u> </u>	0.07.3 111111
Gravel	< 75 mm and > 4.75 mm (#4)	Silt a	nd Clav	< 0.075 mr	n
Coarse Sand	< 4.75 mm and >2.00 mm		Clav	<0.002 mm	า
Medium Sand	< 2.00 mm and > 0.425 mm				
Maximum Particle	Size: #4	Gravel:	0%	Medium Sand:	2%

Maximani i di dicic Size.		Gravei.	070	Micalain Sana.	270
Silt & Clay (% Passing #200):	74%	Total Sand:	26%	Fine Sand:	24%
Assumed Specific Gravity:	2.65	Coarse Sand:	0%	Clay:	13.4%
Liquid Limit	TNP	Plastic Limit	TNP	Plastic Index	TNP

References / Comments / Deviations:

Tyler Copeman Technical Responsibility

Lab Services Manager Position

3/12/2024 Date

Form No. TR-D7928-3 Revision No. 0


Revision Date: 03/18/2019

**BCT6-BSL** 



Log No.: 43-4016 ASTM D7928 & D6913

	S&ME, Inc., 1413 Topside Road, Louisville, TN 37777								
S&ME Project #:	23430142				Report Date:		3/14/2024		
Project Name:	UCOR-Geotech				Test Start Date:	3/1	0/24-3/13/24		
Client Name:	RSI EnTech, LLC								
Address:	203 Victorious Blvd.,	Oak Ridge, TN	37830						
Sample ID:	BCR0053-03	Туре:	REG		Sample D	ate:	2/1/2024		
					De	pth:	NP		



Silt & Clay (% Passing #200): 46% Total Sand: 53% Fine Sand: 39% Assumed Specific Gravity: 2.65 Coarse Sand: 1% Clay: 12.4% Liquid Limit TNP Plastic Limit **TNP** Plastic Index TNP

References / Comments / Deviations:

<u>Tyler Copeman</u> Technical Responsibility Signature

<u>Lab Services Manager</u> *Position*  3/14/2024 Date

Form No. TR-D7928-3

Revision No. 0


Revision Date: 03/18/2019

BCT10-BSL



Log No.: 43-4016 ASTM D7928 & D6913

	S&ME, Inc., 1413 Topside Road, Louisville, TN 37777							
S&ME Project #:	23430142				Report Date:		2/20/2024	
Project Name:	UCOR-Geotech				Test Start Date:	2/1	2/24-2/16/24	
Client Name:	RSI EnTech, LLC							
Address:	203 Victorious Blvd.	, Oak Ridge, TN	l 37830					
Sample ID:	BRC0072-03	Type:	REG		Sample D	ate:	1/11/2024	
					De	pth:	NP	



Cobbles	< 300 mm (12") and > 75 mm	Fine Sand	< 0.425 mm and > 0.075 mm
Gravel	< 75 mm and > 4.75 mm (#4)	Silt and Clav	< 0.075 mm
Coarse Sand	< 4.75 mm and >2.00 mm	Clav	<0.002 mm
Medium Sand	< 2.00 mm and > 0.425 mm		

TVICAIAITI SAITA	<del>//                                     </del>	2 0.423 HIIII				_
Maximum Particle Size:	3/8''	Gravel:	1%	Medium Sand:	5%	
Silt & Clay (% Passing #200):	73%	Total Sand:	26%	Fine Sand:	20%	
Assumed Specific Gravity:	2.65	Coarse Sand:	1%	Clay:	10.2%	
Liquid Limit	TNP	Plastic Limit	TNP	Plastic Index	TNP	

References / Comments / Deviations:

<u>Tyler Copeman</u> Technical Responsibility Signature

<u>Lab Services Manager</u> *Position*  2/20/2024 Date

Form No. TR-D7928-3 Revision No. 0

20%

10%

0%

100

Revision Date: 03/18/2019

BCT11-BSL



Log No.: 43-4016 ASTM D7928 & D6913

10

S&ME, Inc., 1413 Topside Road, Louisville, TN 37777								
S&ME Project #: 23430142					Report Date:		3/14/2024	
Project Name:	UCOR-Geotech				Test Start Date:	3/1	0/24-3/13/24	
Client Name:	RSI EnTech, LLC							
Address:	203 Victorious Blvd.,	Oak Ridge, TN	37830		•			
Sample ID:	BCR0075-03	Туре:	REG		Sample D	ate:	1/31/2024	
					De	oth:	NP	

Brown Clay with Sand Sample Description: 1" 3/4" 3/8" #4 #20 #40 #60 #100 #200 100% 90% 80% 70% Percent Passing 60% 50% 40% 30%

Cobbles	< 300 mm (12") and > 75 mm	Fine Sand	< 0.425 mm and > 0.075 mm
Gravel	< 75 mm and > 4.75 mm (#4)	Silt and Clav	< 0.075 mm
Coarse Sand	< 4.75 mm and >2.00 mm	Clav	<0.002 mm
Medium Sand	< 2.00 mm and > 0.425 mm		

Particle Size (mm)

0.1

0.01

Maximum Particle Size:	3/8''	Gravel:	2%	Medium Sand:	4%
Silt & Clay (% Passing #200):	71%	Total Sand:	27%	Fine Sand:	22%
Assumed Specific Gravity:	2.65	Coarse Sand:	1%	Clay:	10.3%
Liquid Limit	TNP	Plastic Limit	TNP	Plastic Index	TNP

References / Comments / Deviations:

Tyler Copeman
Technical Responsibility

Signature

<u>Lab Services Manager</u> *Position*  3/14/2024 Date

0.001

Form No. TR-D7928-3

Revision No. 0

Log No.:

Revision Date: 03/18/2019


43-4016

BCT12B-BSL



ASTM D7928 & D6913

S&ME, Inc., 1413 Topside Road, Louisville, TN 37777							
S&ME Project #:	23430142	23430142					2/19/2024
Project Name:	UCOR-Geotech				Test Start Date:		2/7/2024
Client Name:	RSI EnTech, LLC						
Address:	203 Victorious Blvd.	, Oak Ridge, TN	I 37830				
Sample ID:	BCR0061-03	Type:	REG		Sample Dat	:e:	1/10/2024
					Dept	:h:	NP



Cobbles	< 300 mm (12") and > 75 mm	Fine Sand	< 0.425 mm and > 0.075 mm
Gravel	< 75 mm and > 4.75 mm (#4)	Silt and Clav	< 0.075 mm
Coarse Sand	< 4.75 mm and >2.00 mm	Clav	<0.002 mm
Medium Sand	< 2.00 mm and > 0.425 mm		

Maximum Particle Size:	3/4"	Gravel:	1%	Medium Sand:	5%	
Silt & Clay (% Passing #200):	69%	Total Sand:	30%	Fine Sand:	24%	
Assumed Specific Gravity:	2.65	Coarse Sand:	1%	Clay:	15.7%	
Liquid Limit	TNP	Plastic Limit	TNP	Plastic Index	TNP	

References / Comments / Deviations:

<u>Tyler Copeman</u> Technical Responsibility Signature

<u>Lab Services Manager</u> *Position*  2/19/2024 Date

Form No. TR-D7928-3

Revision No. 0

Log No.:

Revision Date: 03/18/2019


43-4016

## BCT12B-BSL



#### ASTM D7928 & D6913

S&ME, Inc., 1413 Topside Road, Louisville, TN 37777								
S&ME Project #: 23430142					Report Date:		1/0/1900	
Project Name:	UCOR-Geotech				Test Start Date:	2/1	2/24-2/16/24	
Client Name:	RSI EnTech, LLC							
Address:	203 Victorious Blvd	., Oak Ridge, TN	37830					
Sample ID:	BCR0062-03	Type:	REG		Sample D	ate:	1/10/2024	
					De	nth:	NID	



Coarse Sand < 4.7		< 4.75 mm and	4.75 mm and >2.00 mm (#10)		Clav	<0.002 m	m
	Medium Sand	< 2.00 mm a	nd > 0.425 mm				
	Maximum Particle Si	ize: #10		Gravel:	0%	Medium Sand:	2%
	Silt & Clay (% Passing #2	00): 94%	To	otal Sand:	6%	Fine Sand:	4%
	Assumed Specific Grav	vity: 2.65	Coa	rse Sand:	0%	Clay:	15.1%
	Liquid Li	imit TNP	Pla	stic Limit	TNP	Plastic Index	TNP

Silt and Clay

References / Comments / Deviations:

Gravel

Tyler Copeman
Technical Responsibility

Signature

< 75 mm and > 4.75 mm (#4)

<u>Lab Services Manager</u> *Position*  2/19/2024 Date

This report shall not be reproduced, except in full, without the written approval of S&ME, Inc.

< 0.075 mm

Form No. TR-D7928-3

Revision No. 0

Log No.:

Revision Date: 03/18/2019


43-4016

BCT12A-BSL



ASTM D7928 & D6913

	S&ME, Inc., 1413 Topside Road, Louisville, TN 37						
S&ME Project #:	23430142				Report Date:		2/22/2024
Project Name:	Name: UCOR-Geotech			Test Start Date:	2/2	0/24-2/22/24	
Client Name:	RSI EnTech, LLC						
Address:	203 Victorious Blvd.	, Oak Ridge, TN	I 37830				
Sample ID:	BRC0104-03	Type:	REG		Sample D	ate:	1/10/2024
					Dej	oth:	NP



Cobbles	< 300 mm (12") and > 75 mm	Fine Sand	< 0.425 mm and > 0.075 mm
Gravel	< 75 mm and > 4.75 mm (#4)	Silt and Clav	< 0.075 mm
Coarse Sand	< 4.75 mm and >2.00 mm	Clav	<0.002 mm
Medium Sand	< 2.00 mm and > 0.425 mm		

Maximum Particle Size:	3/8''	Gravel:	1%	Medium Sand:	5%
Silt & Clay (% Passing #200):	69%	Total Sand:	30%	Fine Sand:	24%
Assumed Specific Gravity:	2.65	Coarse Sand:	1%	Clay:	16.6%
Liquid Limit	TNP	Plastic Limit	TNP	Plastic Index	TNP

References / Comments / Deviations:

<u>Tyler Copeman</u> Technical Responsibility Signature

<u>Lab Services Manager</u> *Position*  2/22/2024 Date

Form No. TR-D7928-3

Revision No. 0

Log No.:

Revision Date: 03/18/2019


43-4016

BCT13-BSL



ASTM D7928 & D6913

S&ME, Inc., 1413 Topside Road, Louisville, TN 37777							
S&ME Project #:	Report Date:		2/20/2024				
Project Name:	UCOR-Geotech				Test Start Date:	2/1	2/24-2/16/24
Client Name:	RSI EnTech, LLC						
Address:	203 Victorious Blvd	., Oak Ridge, TN	I 37830				
Sample ID:	BRC0083-03	Type:	REG		Sample D	ate:	1/11/2024
					De	oth:	NP



Cobbles	< 300 mm (12") and > 75 mm	Fine Sand	< 0.425 mm and > 0.075 mm
Gravel	< 75 mm and > 4.75 mm (#4)	Silt and Clav	< 0.075 mm
Coarse Sand	< 4.75 mm and >2.00 mm	Clav	<0.002 mm
Medium Sand	< 2.00 mm and > 0.425 mm		

Maximum Particle Size:	3/4"	Gravel:	3%	Medium Sand:	5%	
Silt & Clay (% Passing #200):	65%	Total Sand:	32%	Fine Sand:	24%	
Assumed Specific Gravity:	2.65	Coarse Sand:	3%	Clay:	6.9%	
Liquid Limit	TNP	Plastic Limit	TNP	Plastic Index	TNP	

References / Comments / Deviations:

<u>Tyler Copeman</u> Technical Responsibility Signature

<u>Lab Services Manager</u> *Position*  2/20/2024 Date

Form No. TR-D7928-3

Revision No. 0

Log No.:


Revision Date: 03/18/2019

BCT14-BSL



43-4016 ASTM D7928 & D6913

S&ME, Inc., 1413 Topside Road, Louisville, TN 37777							
S&ME Project #:	ያME Project #: 23430142					Report Date: 2/20/20	
Project Name:	ne: UCOR-Geotech				Test Start Date:	2/1	2/24-2/16/24
Client Name:	RSI EnTech, LLC						
Address:	203 Victorious Blvd.	, Oak Ridge, TN	l 37830				
Sample ID:	BRC0086-03	Type:	REG		Sample D	ate:	1/2/2024
					De <sub>l</sub>	pth:	NP



Cobbles	< 300 mm (12") and > 75 mm	Fine Sand	< 0.425 mm and > 0.075 mm
Gravel	< 75 mm and > 4.75 mm (#4)	Silt and Clav	< 0.075 mm
Coarse Sand	< 4.75 mm and >2.00 mm	Clav	<0.002 mm
Medium Sand	< 2.00 mm and > 0.425 mm		

Maximum Particle Size:	#10	Gravel:	0%	Medium Sand:	3%
Silt & Clay (% Passing #200):	85%	Total Sand:	15%	Fine Sand:	12%
Assumed Specific Gravity:	2.65	Coarse Sand:	0%	Clay:	25.4%
Liquid Limit	TNP	Plastic Limit	TNP	Plastic Index	TNP

References / Comments / Deviations:

<u>Tyler Copeman</u> Technical Responsibility Signature

<u>Lab Services Manager</u> *Position*  2/20/2024 Date

Form No. TR-D7928-3

Revision No. 0

Log No.:

Revision Date: 03/18/2019


43-4016

BCT14-BSL



ASTM D7928 & D6913

	S&ME, Inc., 1413 Topside Road, Louisville, TN 3777						
S&ME Project #:	23430142				Report Date:		2/20/2024
Project Name:	UCOR-Geotech				Test Start Date:	2/1	2/24-2/16/24
Client Name:	RSI EnTech, LLC						
Address:	203 Victorious Blvd.,	Oak Ridge, TN	I 37830				
Sample ID:	BRC0087-03	Type:	REG		Sample D	ate:	1/2/2024
					Dej	oth:	NP



Cobbles	< 300 mm (12") and > 75 mm	Fine Sand	< 0.425 mm and > 0.075 mm
Gravel	< 75 mm and > 4.75 mm (#4)	Silt and Clav	< 0.075 mm
Coarse Sand	< 4.75 mm and >2.00 mm	Clav	<0.002 mm
Medium Sand	< 2.00 mm and > 0.425 mm		

Maximum Particle Size:	#10	Gravel:	0%	Medium Sand:	2%
Silt & Clay (% Passing #200):	88%	Total Sand:	12%	Fine Sand:	10%
Assumed Specific Gravity:	2.65	Coarse Sand:	0%	Clay:	28.0%
Liquid Limit	TNP	Plastic Limit	TNP	Plastic Index	TNP

References / Comments / Deviations:

<u>Tyler Copeman</u> Technical Responsibility Signature

<u>Lab Services Manager</u> *Position*  2/20/2024 Date

Form No. TR-D7928-3

Revision No. 0


Revision Date: 03/18/2019

BCT15-BSL



Log No.: 43-4016 ASTM D7928 & D6913

S&ME, Inc., 1413 Topside Road, Louisville, TN 37777											
S&ME Project #:	23430142	23430142			Report Date:	e: 2/22/2024					
Project Name:	UCOR-Geotech				Test Start Date:	2/2	0/24-2/22/24				
Client Name:	RSI EnTech, LLC										
Address:	203 Victorious Blvd.	, Oak Ridge, TN	l 37830								
Sample ID:	BRC0093-03	Type:	REG		Sample Date: 1/3/202		1/3/2024				
					De <sub>l</sub>	oth:	NP				



Cobbles	< 300 mm (12") and > 75 mm	Fine Sand	< 0.425 mm and > 0.075 mm
Gravel	< 75 mm and > 4.75 mm (#4)	Silt and Clav	< 0.075 mm
Coarse Sand	< 4.75 mm and >2.00 mm	Clav	<0.002 mm
Medium Sand	< 2.00 mm and > 0.425 mm		

Maximum Particle Size:	#4	Gravel:	0%	Medium Sand:	5%
Silt & Clay (% Passing #200):	69%	Total Sand:	31%	Fine Sand:	25%
Assumed Specific Gravity:	2.65	Coarse Sand:	1%	Clay:	11.2%
Liquid Limit	TNP	Plastic Limit	TNP	Plastic Index	TNP

References / Comments / Deviations:

<u>Tyler Copeman</u> Technical Responsibility Signature

<u>Lab Services Manager</u> *Position*  2/22/2024 Date

Form No. TR-D7928-3 Revision No. 0


Revision Date: 03/18/2019

**HCTREF-BSL** 



Log No.: 43-4016 ASTM D7928 & D6913

3										
S&ME, Inc., 1413 Topside Road, Louisville, TN 37777										
S&ME Project #:	23430142				Report Date:	3	3/18/2024			
Project Name:	UCOR-Geotech			-	Test Start Date:	3/12	2/24 - 3/14/24			
Client Name:	RSI EnTech, LLC									
Address:	203 Victorious Blvd.,	, Oak Ridge, TN	l 37830							
Sample ID:	BCR0096-03	Type:	REG		Sample Date: 2,		2/2/2024			
		·	·		De	pth:	NP			



CODDICS	<u> </u>		e Janu	\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	$\mu_{\rm c}$
Gravel	< 75 mm and > 4.75 mm (#4)	Silt a	and Clav	< 0.075 mn	า
Coarse Sand	< 4.75 mm and >2.00 mm		Clav	<0.002 mm	1
Medium Sand	< 2.00 mm and > 0.425 mm				
Maximum Particle S	Size: 3/8"	Gravel:	1%	Medium Sand:	7%

7%	Medium Sand:	1%	Gravel:	3/8"	Maximum Particle Size:
35%	Fine Sand:	45%	Total Sand:	54%	Silt & Clay (% Passing #200):
13.2%	Clay:	3%	Coarse Sand:	2.65	Assumed Specific Gravity:
TNP	Plastic Index	TNP	Plastic Limit	TNP	Liquid Limit

References / Comments / Deviations:

Tyler Copeman Technical Responsibility

Lab Services Manager Position

3/18/2024 Date

**CHANNEL SEDIMENT** 

This page intentionally left blank.

Revision No. 0

Revision Date: 08/30/17

# **Sieve Analysis of Soils**

BCT1-CH



ASTM D6913 Method A **Quality Assurance** 

	S&N	ME, Inc Knox	ville: 1413 T	opside Roa	d, Louisv	/ille, TN 3	37777		
Project #:	2343014	42			Re	eport Da	te:	2/22/2024	
	UCOR-Ge				T	est Date(	(s):	1/30/2024	
	RSI EnTed					_			
	Oak Ridg	e, TN							
Sample ID: BCR			Туре:	REG		Sample		1/4/2024	
Sample Log No.:			Sample:				Depth:	NP	
Sample Description	on: F	Poorly Graded	Gravel with Sa	nd					
_	" 1.5"	1" 3/4" 3/8"	#4 #10	#20	#40 #6	50 #100	#200		
100%		• • • • • • • • • • • • • • • • • • • •				$\overline{}$			7 I
90%									
H		-+							- 1
80%									
70%		$\longrightarrow$							- 1
Sin									$\blacksquare$
Percent Passing (%)  900  900  900  900  900  900  900  9									11
50%			$\longleftarrow$						$\blacksquare$
40%									11
<b> </b>			+N $+$ +						-
30%									
20%									_
4004			+++						-
10%									
0%		10.00				<u> </u>			
100.00		10.00	Millimeters	1.00			.10		0.01
Cobbles	< :	300 mm (12") and	> 75 mm (3")	Fi	ne Sand		< 0.425 m	m and > 0.075 i	mm
Gravel		< 75 mm and > 4.	. ,	Silt	and Clay		<	0.075 mm	
Coarse Sand		4.75 mm and >2.0							
Medium Sand Maximum Part		2.00 mm and > 0.4	, ,	rca Cand	28%		Fina	Sand 09	)/
Maximum Part		3/4		rse Sand um Sand					
Lieu	Gravel	61%		tic Limit:	10%		Plastic I	k Clay 09	
Liqu	ıid Limit:	TNP	Plas	uc Limit.	TNP		Plastic i	ndex: TN	IP .
Descripti	ion of San	nd & Gravel Par	ticles:	Round	led		Angula	ar 🗵	
•	& Durable						d & Friabl		
Notes / Deviations /	References:								
,	<u> </u>								
			1. Cal	/					
•	<u>opeman</u>		per con			<u>ices Mar</u>	<u>nager</u>	2/22/20	<u> 24</u>
Technical Re		/	Signature			Position		Date	
	This r	eport shall not be rep	produced, except in	full, without the	e written app	proval of S&	ME, Inc.		

Form No. TR-D7928-3 Revision No. 0

Log No.:

Revision Date: 03/18/2019


43-4016

BCT2-CH



ASTM D7928 & D6913

S&ME, Inc., 1413 Topside Road, Louisville, TN 37777										
S&ME Project #:	23430142			Report Date:		2/23/2024				
Project Name:	UCOR-Geotech			Test Start Date:		2/1/2024				
Client Name:	RSI EnTech, LLC									
Address:	203 Victorious Blvd	., Oak Ridge, TN	37830							
Sample ID:	BCR0020-03	Type:	REG	Sample	Sample Date: 1/4/2					
				Γ	enth.	NP				



References / Comments / Deviations:

Liquid Limit

Victoria Igoe Technical Responsibility Signature

2.65

**TNP** 

Senior Engineering Technician Position

TNP

2/23/2024 Date

3.2%

**TNP** 

Clay:

Plastic Index

This report shall not be reproduced, except in full, without the written approval of S&ME, Inc.

Plastic Limit


Form No. TR-D7928-3 Revision No. 0

Revision Date: 03/18/2019 Log No.: 43-4016 BCT3-CH



ASTM D7928 & D6913

_og		ASTITUT	20 a D0513	,					
S&ME, Inc., 1413 Topside Road, Louisville, TN 37777									
S&ME Project #:	23430142				Report Date:	2	/23/2024		
Project Name:	UCOR-Geotech				Test Start Date:	2	2/1/2024		
Client Name:	RSI EnTech, LLC								
Address:	203 Victorious Blvd.	, Oak Ridge, TN	l 37830						
Sample ID:	BCR0021-03	Type:	REG		Sample Dat	e:	1/8/2024		
		_			Dept	h:	NP		



Maximum Particle Size:	#10	Gravel:	0%	Medium Sand:	57%
Silt & Clay (% Passing #200):	7%	Total Sand:	93%	Fine Sand:	36%
Assumed Specific Gravity:	2.65	Coarse Sand:	0%	Clay:	3.7%
Liquid Limit	TNP	Plastic Limit	TNP	Plastic Index	TNP

**References / Comments / Deviations:** 

<u>Victoria Igoe</u> Technical Responsibility Signature

Senior Engineering Technician

Position

2/23/2024 Date

Form No. TR-D7928-3 Revision No. 0


Revision Date: 03/18/2019

BCT4-CH



Log No.: 43-4016 ASTM D7928 & D6913

	S&ME, Inc.	isville,	TN 37777				
S&ME Project #:	23430142				Report Date:	2/23/2024	
Project Name:	UCOR-Geotech				Test Start Date:	2/1/2024	
Client Name:	RSI EnTech, LLC						
Address:	203 Victorious Blvd.	, Oak Ridge, TN	N 37830				
Sample ID:	BCR0022-03	Type:	REG		Sample Date: 1/8/20		
·					Dept	h: NP	



Silt & Clay (% Passing #200): 15% Total Sand: 85% Fine Sand: 19% Assumed Specific Gravity: Coarse Sand: 0% 2.65 3.3% Clay: Liquid Limit **TNP** Plastic Limit TNP Plastic Index **TNP** 

References / Comments / Deviations:

<u>Victoria Igoe</u> Technical Responsibility Signature

Senior Engineering Technician

Position

2/23/2024 Date

Revision No. 0

Form No. TR-D7928-3

Revision Date: 03/18/2019


BCT5-CH



Log No.: 43-4016 ASTM D7928 & D6913

S&ME, Inc., 1413 Topside Road, Louisville, TN 37777									
S&ME Project #:	23430142				Report Date:		3/8/2024		
Project Name:	UCOR-Geotech				Test Start Date:	3/	/3/24-3/6/24		
Client Name:	RSI EnTech, LLC								
Address:	203 Victorious Blvd.	, Oak Ridge, TN	I 37830						
Sample ID:	BCR0023-04	Туре:	REG		Sample Da	ite:	2/1/2024		
					Dep	th:	NP		

Sample Description: Brown Poorly Graded Sand with Silt



Cobbles	< 300 mm (12") and > 75 mm	Fine Sand	< 0.425 mm and > 0.075 mm
Gravel	< 75 mm and > 4.75 mm (#4)	Silt and Clav	< 0.075 mm
Coarse Sand	< 4.75 mm and >2.00 mm	Clav	<0.002 mm
Medium Sand	< 2.00 mm and > 0.425 mm		

75%	Medium Sand:	0%	Gravel:	#4	Maximum Particle Size:
15%	Fine Sand:	91%	Total Sand:	9%	Silt & Clay (% Passing #200):
1.8%	Clay:	1%	Coarse Sand:	2.65	Assumed Specific Gravity:
TNP	Plastic Index	TNP	Plastic Limit	TNP	Liquid Limit

References / Comments / Deviations:

Tyler Copeman
Technical Responsibility

Signature

<u>Lab Services Manager</u> *Position*  3/8/2024 Date

Revision No. 0

Revision Date: 03/18/2019


Form No. TR-D7928-3

BCT6-CH



Log No.: 43-4016 ASTM D7928 & D6913

	S&ME, Inc	., 1413 Topsid	e Road, Loui	sville, 1	TN 37777		
S&ME Project #:	23430142				Report Date:		3/8/2024
Project Name:	UCOR-Geotech				Test Start Date:	3/	3/24-3/6/24
Client Name:	RSI EnTech, LLC						
Address:	203 Victorious Blvd.,	Oak Ridge, TN	37830				
Sample ID:	BCR0024-04	Type:	REG		Sample Da	ite:	2/1/2024
					Dep	th:	NP



Cobbles	< 300 mm (12") and > 75 mm	Fine Sand	< 0.425 mm and > 0.075 mm
Gravel	< 75 mm and > 4.75 mm (#4)	Silt and Clav	< 0.075 mm
Coarse Sand	< 4.75 mm and >2.00 mm	Clav	<0.002 mm
Medium Sand	< 2.00 mm and > 0.425 mm		

Med	%	(	Gravel:		#4	icle Size:	Maximum Par
	%	7	Total Sand:		29%	ng #200):	Silt & Clay (% Pass
	%	(	Coarse Sand:	(	2.65	c Gravity:	Assumed Specif
Pla	NΡ	Т	Plastic Limit		TNP	quid Limit	Li

References / Comments / Deviations:

Tyler Copeman Technical Responsibility

Lab Services Manager Position

3/8/2024 Date

Form No. TR-D7928-3 Revision No. 0

Revision Date: 03/18/2019

BCT7-CH



Log No.: 43-4016 ASTM D7928 & D6913

	S&ME, Inc	., 1413 Topsid	e Road, Loui	sville, 1	TN 37777		
S&ME Project #:	23430142				Report Date:		3/8/2024
Project Name:	UCOR-Geotech				Test Start Date:	3/	/3/24-3/6/24
Client Name:	RSI EnTech, LLC						
Address:	203 Victorious Blvd.,	Oak Ridge, TN	37830				
Sample ID:	BCR0025-04	Туре:	REG		Sample Da	ite:	2/7/2024
					Dep	th:	NP

Sample Description: Grayish Brown Silt 1" 3/4" 3/8" #20 #40 #60 #100 #200 100% 90% 80% 70% Percent Passing 60% 50% 40% 30% 20% 10% 0% 10 1 0.1 0.01 0.001 100 Particle Size (mm) < 0.425 mm and > 0.075 mm Cobbles < 300 mm (12") and > 75 mm Fine Sand Gravel < 75 mm and > 4.75 mm (#4) Silt and Clav < 0.075 mm < 4.75 mm and >2.00 mm <0.002 mm Coarse Sand Clav < 2.00 mm and > 0.425 mm Medium Sand Medium Sand: Maximum Particle Size: #20 Gravel: 0% 2%

References / Comments / Deviations:

Silt & Clay (% Passing #200):

Assumed Specific Gravity:

Liquid Limit

<u>Tyler Copeman</u> Technical Responsibility Signature

86%

2.65

**TNP** 

<u>Lab Services Manager</u> <u>Position</u>

14%

0%

**TNP** 

3/8/2024 Date

12%

4.1%

TNP

This report shall not be reproduced, except in full, without the written approval of S&ME, Inc.

Total Sand:

Coarse Sand:

Plastic Limit

Fine Sand:

Plastic Index

Clay:

Revision No. 0

Revision Date: 03/18/2019

Form No. TR-D7928-3


BCT8-CH



Log No.: 43-4016 ASTM D7928 & D6913

	S&ME, Inc	., 1413 Topsid	e Road, Louis	sville, T	N 37777		
S&ME Project #:	23430142				Report Date:		3/19/2024
Project Name:	UCOR-Geotech				Test Start Date:	3/7	7/24-3/11/24
Client Name:	RSI EnTech, LLC						
Address:	203 Victorious Blvd.,	Oak Ridge, TN	I 37830				
Sample ID:	BCR0031-04	Type:	REG		Sample Da	ate:	2/5/2024
					Dep	oth:	NP

Sample Description: Brown Well-graded Sand with Silt 1" 3/4" 3/8" #20 #60 #100 #200 100%



Cobbles	< 300 mm (12") and > 75 mm	Fine Sand	< 0.425 mm and > 0.075 mm
Gravel	< 75 mm and > 4.75 mm (#4)	Silt and Clav	< 0.075 mm
Coarse Sand	< 4.75 mm and >2.00 mm	Clav	<0.002 mm
Medium Sand	< 2.00 mm and > 0.425 mm		

Maximum Particle Size:	#10	Gravel:	0%	Medium Sand:	67%
Silt & Clay (% Passing #200):	8%	Total Sand:	92%	Fine Sand:	25%
Assumed Specific Gravity:	2.65	Coarse Sand:	0%	Clay:	3.8%
Liquid Limit	TNP	Plastic Limit	TNP	Plastic Index	TNP

References / Comments / Deviations:

Tyler Copeman Technical Responsibility

Lab Services Manager Position

3/19/2024 Date

Form No. TR-D7928-3 Revision No. 0


Revision Date: 03/18/2019

Log No.: 43-4016 ASTM D7928 & D6913 BCT9-CH



9	• • •	, 13111 27	20000					
S&ME, Inc., 1413 Topside Road, Louisville, TN 37777								
S&ME Project #:	23430142				Report Date:		3/8/2024	
Project Name:	UCOR-Geotech				Test Start Date:	3/3	/24 - 3/6/24	
Client Name:	RSI EnTech, LLC							
Address:	203 Victorious Blvd.,	Oak Ridge, TN	37830					
Sample ID:	BCR0028-04	Type:	REG		Sample Da	ate:	2/5/2024	
					Dor	+b.	ND	

Depth: NΡ



References / Comments / Deviations:

Silt & Clay (% Passing #200):

Assumed Specific Gravity:

Liquid Limit

Tyler Copeman Technical Responsibility

69%

2.65

TNP

Lab Services Manager Position

31%

0%

**TNP** 

3/8/2024 Date

21%

5.9%

TNP

This report shall not be reproduced, except in full, without the written approval of S&ME, Inc.

Total Sand:

Coarse Sand:

Plastic Limit

Fine Sand:

Plastic Index

Clay:

Revision No. 0

Revision Date: 03/18/2019

Form No. TR-D7928-3


ASTM D7928 & D6913





Log No.:	43-4016		ASTM D7	928 & D6913					
S&ME, Inc., 1413 Topside Road, Louisville, TN 37777									
S&ME Proje	ect #:	23430142			Report Dat	te:	3/12/2024		
Project Nan	ne:	UCOR-Geotech			Test Start D	ate: 3,	/3/24-3/6/24		
Client Name	e:	RSI EnTech, LLC							
Address:		203 Victorious Blvd.,	Oak Ridge, TN	I 37830					
Sample ID:		BCR0029-03	Type:	REG	Sam	ple Date:	1/31/2024		
						Donth:	ND		

Depth: NΡ



Gravei	< /	<u> </u>	[[][[](# <del>4</del> )	SIII	. and Clav	< 0.075 11	1111	
Coarse Sand	<	4.75 mm and >2	.00 mm		Clav	<0.002 m	m	
Medium Sand	< 7	2.00 mm and > 0.	425 mm					
Maximum Particle S	ize:	#10		Gravel:	0%	Medium Sand:	44%	
Silt & Clay (% Passing #2	200):	21%	To	tal Sand:	79%	Fine Sand:	35%	

Assumed Specific Gravity: 2.65 Coarse Sand: 0% Clay: 2.5% TNP Liquid Limit TNP Plastic Limit TNP Plastic Index

References / Comments / Deviations:

Tyler Copeman Technical Responsibility

<u>Lab Services Manager</u> Position

3/12/2024 Date

Form No. TR-D7928-3

Revision No. 0

Revision Date: 03/18/2019 Log No.: 43-4016


BCT11-CH



ASTM D7928 & D6913

		4440 = 11									
	S&ME, Inc., 1413 Topside Road, Louisville, TN 37777										
S&ME Project #:	23430142				Report Date:		3/12/2024				
Project Name:	UCOR-Geotech				Test Start Date:	3/	7/24-3/11/24				
Client Name:	RSI EnTech, LLC										
Address:	203 Victorious Blvd.	, Oak Ridge, TN	I 37830								
Sample ID:	BCR0030-04	Type:	REG		Sample Date: 1/31		1/31/2024				
					Der	oth:	NP				





Gravel	< 75 mm and > 4.75 mm (#4)	Silt and Clav		< 0.075 mm	
Coarse Sand	< 4.75 mm and >2.00 mm		Clav	<0.002 mr	n
Medium Sand	< 2.00 mm and > 0.425 mm				
Maximum Particle S	Size: 3/8"	Gravel:	1%	Medium Sand:	53%

	- , -				
Silt & Clay (% Passing #200):	11%	Total Sand:	88%	Fine Sand:	34%
Assumed Specific Gravity:	2.65	Coarse Sand:	1%	Clay:	5.0%
Liquid Limit	TNP	Plastic Limit	TNP	Plastic Index	TNP

References / Comments / Deviations:

Tyler Copeman Technical Responsibility

Lab Services Manager Position

3/12/2024 Date

Form No. TR-D7928-3

Revision No. 0

Log No.:

Revision Date: 03/18/2019

43-4016

BCT12B-CH



ASTM D7928 & D6913

	S&ME, Inc., 1413 Topside Road, Louisville, TN 37777										
S&ME Project #:	23430142				Report Date:		3/8/2024				
Project Name:	UCOR-Geotech				Test Start Date:	3/	3/24-3/6/24				
Client Name:	RSI EnTech, LLC										
Address:	203 Victorious Blvd.,	Oak Ridge, TN	l 37830								
Sample ID:	BCR0026-04	Type:	REG		Sample Date: 1/		1/29/2024				
					Dep	th:	NP				

Sample Description: Brown Poorly Graded Sand with Silt 1" 3/4" #20 #60 #100 #200 100% 90% 80% 70% Percent Passing 60% 50% 40% 30% 20% 10% 0% 10 0.1 0.01 0.001 100 Particle Size (mm)

Cobbles	< 300 mm (12") and > 75 mm	Fine Sand	< 0.425 mm and > 0.075 mm
Gravel	< 75 mm and > 4.75 mm (#4)	Silt and Clav	< 0.075 mm
Coarse Sand	< 4.75 mm and >2.00 mm	Clav	<0.002 mm
Madium Cand	2.00 mm and > 0.425 mm		

Maximum Particle Size:	#10	Gravel:	0%	Medium Sand:	44%	
Silt & Clay (% Passing #200):	11%	Total Sand:	89%	Fine Sand:	45%	
Assumed Specific Gravity:	2.65	Coarse Sand:	0%	Clay:	4.7%	
Liquid Limit	TNP	Plastic Limit	TNP	Plastic Index	TNP	

References / Comments / Deviations:

Tyler Copeman
Technical Responsibility

Signature

<u>Lab Services Manager</u> *Position*  3/8/2024 Date

Revision No. 0

Form No. TR-D7928-3

Revision Date: 03/18/2019

BCT12B-CH



Log No.: 43-4016 ASTM D7928 & D6913

S&ME, Inc., 1413 Topside Road, Louisville, TN 37777									
S&ME Project #:	23430142				Report Date:		3/8/2024		
Project Name:	UCOR-Geotech				Test Start Date:	3/	/3/24-3/6/24		
Client Name:	RSI EnTech, LLC								
Address:	203 Victorious Blvd.	, Oak Ridge, TN	37830						
Sample ID:	BCR0027-04	Туре:	REG		Sample Date:		1/29/2024		
					Dep	th:	NP		

Depth: NΡ



Copples	< 500 mm (12 ) and > 75 mm	Fille Salid	1 < 0.425 mm and > 0.075 mm
Gravel	< 75 mm and > 4.75 mm (#4)	Silt and Clav	< 0.075 mm
Coarse Sand	< 4.75 mm and >2.00 mm	Clav	<0.002 mm
Medium Sand	< 2.00 mm and > 0.425 mm		
Maximum Particle 9	Sizo: #10	Gravel: 0%	Madium Cand: 410/

%	edium Sand:	0%	Gravei:	#10	Maximum Particle Size:
5%	Fine Sand:	56%	Total Sand:	44%	Silt & Clay (% Passing #200):
3%	Clay:	0%	Coarse Sand:	2.65	Assumed Specific Gravity:
NΡ	Plastic Index	TNP	Plastic Limit	TNP	Liquid Limit

References / Comments / Deviations:

Tyler Copeman Technical Responsibility

Lab Services Manager Position

3/8/2024 Date

Revision No. 0

Form No. TR-D7928-3


Revision Date: 03/18/2019

BCT12A-CH



Log No.: 43-4016 ASTM D7928 & D6913

	S&ME, Inc., 1413 Topside Road, Louisville, TN 37777									
S&ME Project #:	23430142				Report Date:		3/18/2024			
Project Name:	UCOR-Geotech		Test Start Date:	rt Date: 3/12/24-3/14/						
Client Name:	RSI EnTech, LLC									
Address:	203 Victorious Blvd.	Oak Ridge, TN	37830							
Sample ID:	BCR0103-03	Туре:	REG		Sample Date: 1/29/		1/29/2024			
					De	pth:	NP			



CODDIC3	<u> </u>		- Juliu	V 0.423 Hilli and > 0.013 Hilli		
Gravel	< 75 mm and > 4.75 mm (#4)	Silt and Clav		< 0.075 mm		
Coarse Sand	< 4.75 mm and >2.00 mm	(	Clav	<0.002 mm	n	
Medium Sand	< 2.00 mm and > 0.425 mm					
Maximum Particle S	ize: #4	Gravel:	0%	Medium Sand:	40%	

Silt & Clay (% Passing #200): 22% Total Sand: 78% Fine Sand: 38% Assumed Specific Gravity: 2.65 Coarse Sand: 0% Clay: 8.5% TNP TNP **TNP** Liquid Limit Plastic Limit Plastic Index

References / Comments / Deviations:

Tyler Copeman Technical Responsibility

Lab Services Manager Position

3/18/2024 Date

Form No. TR-D7928-3

Revision No. 0

Log No.:

Revision Date: 03/18/2019


43-4016

BCT13-CH



ASTM D7928 & D6913

3							
S&ME, Inc., 1413 Topside Road, Louisville, TN 37777							
S&ME Project #:	23430142				Report Date: 3/12/20		3/12/2024
Project Name:	UCOR-Geotech				Test Start Date:	3/7	7/24-3/11/24
Client Name:	RSI EnTech, LLC						
Address:	203 Victorious Blvd.,	Oak Ridge, TN	l 37830				
Sample ID:	BCR0032-03	Type:	REG		Sample Da	ate:	1/29/2024
					Dep	oth:	NP



Cobbles	< 300 mm (12") and > 75 mm	Fine Sand	< 0.425 mm and > 0.075 mm
Gravel	< 75 mm and > 4.75 mm (#4)	Silt and Clav	< 0.075 mm
Coarse Sand	< 4.75 mm and > 2.00 mm	Clav	<0.002 mm
Medium Sand	< 2.00 mm and > 0.425 mm		

Maximum Particle Size:	3/4''	Gravel:	2%	Medium Sand:	55%
Silt & Clay (% Passing #200):	8%	Total Sand:	90%	Fine Sand:	35%
Assumed Specific Gravity:	2.65	Coarse Sand:	0%	Clay:	5.3%
Liquid Limit	TNP	Plastic Limit	TNP	Plastic Index	TNP

References / Comments / Deviations:

Tyler Copeman
Technical Responsibility

Signature

<u>Lab Services Manager</u> *Position*  3/12/2024 Date


Form No. TR-D7928-3 Revision No. 0

Revision Date: 03/18/2019 Log No.: 43-4016 BCT14-CH



ASTM D7928 & D6913

S&ME, Inc., 1413 Topside Road, Louisville, TN 37777							
S&ME Project #:	23430142			Report D	ate:	2/23/2024	
Project Name:	UCOR-Geotech			Test Start	Date:	2/1/2024	
Client Name:	RSI EnTech, LLC						
Address:	203 Victorious Blvd	., Oak Ridge, TN	37830				
Sample ID:	BCR0033-04	Type:	REG	Sa	mple Date:	1/2/2024	
					Denth:	NP	



References / Comments / Deviations:

Assumed Specific Gravity:

Liquid Limit

<u>Victoria Igoe</u> Technical Responsibility Signature

2.65

**TNP** 

Senior Engineering Technician

Position

0%

TNP

2/23/2024 Date

29.0%

**TNP** 

Clay:

Plastic Index

This report shall not be reproduced, except in full, without the written approval of S&ME, Inc.

Coarse Sand:

Plastic Limit

## **Particle-Size Distribution**

Form No. TR-D7928-3 Revision No. 0

Log No.:

Revision Date: 03/18/2019


43-4016

BCT14-CH



ASTM D7928 & D6913

	S&ME, Inc.	, 1413 Topsid	e Road, Loui	isville, <sup>-</sup>	TN 37777		
S&ME Project #:	23430142				Report Date:		2/23/2024
Project Name:	UCOR-Geotech				Test Start Date:		2/1/2024
Client Name:	RSI EnTech, LLC						
Address:	203 Victorious Blvd.	, Oak Ridge, TN	l 37830				
Sample ID:	BCR0034-04	Туре:	REG		Sample Dat	te:	1/2/2024
					Dept	:h:	NP



Maximum Particle Size:	#10	Gravel:	0%	Medium Sand:	6%
Silt & Clay (% Passing #200):	90%	Total Sand:	10%	Fine Sand:	4%
Assumed Specific Gravity:	2.65	Coarse Sand:	0%	Clay:	29.5%
Liquid Limit	TNP	Plastic Limit	TNP	Plastic Index	TNP

**References / Comments / Deviations:** 

<u>Victoria Igoe</u> Technical Responsibility

Signature

Senior Engineering Technician
Position

2/23/2024 Date

This report shall not be reproduced, except in full, without the written approval of S&ME, Inc.

## **Particle-Size Distribution**

Form No. TR-D7928-3 Revision No. 0


Revision Date: 03/18/2019

BCT15-CH



Log No.: 43-4016 ASTM D7928 & D6913

	S&ME, Inc	, 1413 Topside	Road, Louis	ville, TN 37777		
S&ME Project #:	23430142			Report Date:		2/23/2024
Project Name:	UCOR-Geotech			Test Start Dat	e:	2/1/2024
Client Name:	RSI EnTech, LLC					
Address:	203 Victorious Blvd	., Oak Ridge, TN	37830			
Sample ID:	BCR0035-03	Туре:	REG	Sampl	e Date:	1/3/2024
					Depth:	NP



Maximum Particle Size:	#10	Gravel:	0%	Medium Sand:	29%
Silt & Clay (% Passing #200):	52%	Total Sand:	48%	Fine Sand:	19%
Assumed Specific Gravity:	2.65	Coarse Sand:	0%	Clay:	7.5%
Liquid Limit	TNP	Plastic Limit	TNP	Plastic Index	TNP

**References / Comments / Deviations:** 

Victoria Igoe Technical Responsibility

Signature

Senior Engineering Technician Position

2/23/2024 Date

This report shall not be reproduced, except in full, without the written approval of S&ME, Inc.

## **Particle-Size Distribution**

Revision No. 0

Form No. TR-D7928-3


Revision Date: 03/18/2019

**HCTREF-CH** 



Log No.: 43-4016 ASTM D7928 & D6913

3					
	S&ME, Inc	, 1413 Topsid	e Road, Louis	ville, TN 37777	
S&ME Project #:	23430142			Report Date:	3/12/2024
Project Name:	UCOR-Geotech			Test Start Date:	3/7/24-3/11/24
Client Name:	RSI EnTech, LLC				
Address:	203 Victorious Blvd.	, Oak Ridge, TN	I 37830		
Sample ID:	BCR0036-04	Type:	REG	Sample Da	ate: 2/2/2024
				Dep	oth: NP



References / Comments / Deviations:

Silt & Clay (% Passing #200):

Assumed Specific Gravity:

Liquid Limit

<u>Tyler Copeman</u> Technical Responsibility Signature

3%

2.65

TNP

<u>Lab Services Manager</u> <u>Position</u>

95%

53%

TNP

3/12/2024 Date

1%

0.5%

TNP

This report shall not be reproduced, except in full, without the written approval of S&ME, Inc.

Total Sand:

Coarse Sand:

Plastic Limit

Fine Sand:

Plastic Index

Clay:

# APPENDIX E. BEAR CREEK VALLEY REMEDIAL SITE EVALUATION ANALYTE CORRELATIONS

# **E.1.** MULTIVARIATE ANALYSIS

Results of analyses of Bear Creek floodplain and bank soil samples (combined) and Bear Creek channel sediment samples were evaluated for analyte pairwise and multivariate correlation. Correlations range from -1 (perfect negative correlation) to +1 (perfect positive correlation). Negative correlations indicate that, as concentrations of one variable increase, the concentrations of the second variable decrease. Positive correlations indicate that, as concentrations of one variable increase, the concentrations of the second variable also increase. Correlations close to zero indicate no correlation between the variables. When all concentrations for both pairs of variables were detected, the Pearson correlation was calculated because it uses all the information available in ratio concentration data. When at least one concentration in either variable was not detected. Kendall's tau correlation was calculated because it allows for non-detections (Helsel 2005). With each correlation, a two-sided p-value was also calculated to test the null hypothesis of zero correlation versus the alternative hypothesis that the correlation is not zero. A p-value is the probability of observing a test statistic at least as large as the one observed given the null hypothesis is true. A significance level of 0.05 was used to determine whether a correlation was significantly different from zero. At least one-half of the concentrations for both variables must be detected to calculate the correlation. Individual pairwise significance levels were not adjusted for multiple comparisons because the purpose of the correlations was to identify analytes as potential candidate variables for multivariate regressions.

The pairwise correlation tables were used to identify analytes that are significantly correlated with mercury in soil and sediment. The tables contain the correlation coefficient for each analyte pair and use a color scheme to reflect the degree of significance of each correlation pair. Correlations that test the null hypothesis of zero correlation and are significant at the 0.05 significance level are highlighted by significance. Table E.1 contains all pairwise correlations for floodplain and creek bank soil combined. Table E.2 contains all pairwise correlations for floodplain and creek bank soil combined for particle size #10. Table E.3 contains all pairwise correlations for all sediment combined. To examine correlation with particle size, Table E.4 contains all pairwise correlations for sediment combined for particle size #10. There were insufficient sediment data to calculate correlations for the other particle sizes.

Combinations resulting in correlations that are significant at the 0.05 significance level are highlighted by significance:

- Two-sided *p*-values ranging between 0.01 and 0.05 are highlighted in blue.
- Two-sided p-values ranging between 0.001 and < 0.01 are highlighted in green.
- Two-sided p-values ranging between 0.0001 and < 0.001 are highlighted in yellow.
- Two-sided p-values that are < 0.0001 are highlighted in red.

Non-highlighted correlations are not significant at the 0.05 significance level. Two-sided p-values for correlations are highly influenced by the number of samples used in the correlations. For example, a correlation of 0.8 will likely be highly significant when the number of samples (n) = 48 samples are used, but may not be significant when n = 3.

Table E.1. Analyte pairwise correlation matrix for Bear Creek floodplain and creek bank soils

	E		Anions	<b>S</b>				Seq	uential extraction	on .			Other inorganics	Wet che	mistry
Chemical	Freq. of detection	Chloride	Fluoride	Nitrate	Sulfate	Mercury (F1)	Mercury (F2)	Mercury (F3)	Mercury (F4)	Mercury (F5)	Mercury (F6)	Mercury (FS)	Methylmercury	Sulfide	TOC
Chloride	48 / 48	NA	0.067	0.219	0.853	0.061	0.076	0.061	0.107	-0.305	-0.443	-0.311	-0.003	0.091	-0.061
Fluoride	48 / 48	0.067	NA	0.155	0.277	0.667	0.473	0.667	0.687	0.866	0.168	0.933	0.089	-0.050	0.145
Nitrate	44 / 48	0.219	0.155	NA	0.167	0.424	0.412	0.485	0.382	0.455	0.290	0.473	0.332	-0.089	0.468
Sulfate	48 / 48	0.853	0.277	0.167	NA	-0.121	-0.290	-0.182	-0.076	-0.543	-0.382	-0.492	0.100	-0.010	0.052
Aluminum	48 / 48	0.255	0.458	0.111	0.343	0.394	0.382	0.333	0.412	0.039	0.229	0.186	0.068	0.093	0.337
Arsenic	48 / 48	0.070	0.316	0.166	0.165	0.321	0.492	0.260	0.339	0.109	0.339	0.185	0.196	0.034	0.430
Barium	48 / 48	0.092	0.452	0.264	0.322	0.364	0.168	0.364	0.321	0.072	-0.137	0.104	0.319	0.051	0.387
Beryllium	46 / 48	0.257	-0.093	0.245	0.217	0.061	0.107	0.061	0.076	0.091	-0.351	0.076	0.081	0.140	0.092
Boron	26 / 48	0.347	0.003	0.282	0.213	0.254	0.288	0.191	0.288	0.350	-0.144	0.208	0.100	0.189	0.176
Cadmium	48 / 48	0.029	0.792	0.433	0.303	0.485	0.534	0.485	0.504	0.724	0.015	0.872	0.529	0.039	0.466
Calcium	48 / 48	0.071	0.395	0.372	0.355	-0.061	-0.168	-0.121	-0.015	-0.318	-0.351	-0.247	0.324	-0.056	0.385
Chromium	48 / 48	0.216	0.142	0.068	0.254	-0.031	0.140	-0.031	-0.016	-0.326	-0.016	-0.259	0.092	0.121	0.292
Cobalt	48 / 48	0.205	0.538	0.220	0.376	0.182	0.351	0.182	0.199	0.354	0.015	0.237	0.160	0.092	0.308
Copper	48 / 48	-0.137	0.629	0.257	0.023	0.576	0.595	0.515	0.565	0.750	0.473	0.889	0.383	0.152	0.598
Iron	48 / 48	0.473	0.113	0.089	0.502	0.062	0.171	0.000	0.078	-0.226	-0.016	-0.196	0.095	0.119	-0.004
Lead	48 / 48	-0.018	0.617	0.212	0.047	0.485	0.718	0.485	0.504	0.862	0.382	0.941	0.234	0.098	0.494
Lithium	48 / 48	-0.101	0.781	0.226	0.088	0.424	0.443	0.364	0.473	0.416	0.137	0.566	0.394	0.057	0.477
Magnesium	48 / 48	-0.002	0.145	0.127	0.232	-0.242	-0.290	-0.242	-0.199	-0.389	-0.473	-0.359	0.234	0.105	0.442
Manganese	48 / 48	0.132	0.435	0.226	0.339	0.273	0.076	0.273	0.229	0.215	-0.168	0.198	0.250	0.032	0.347
Mercury	48 / 48	-0.097	0.551	0.373	0.119	0.939	0.809	0.939	0.962	0.843	0.260	0.977	0.546	-0.031	0.464
Molybdenum	32 / 48	0.015	0.350	0.102	-0.045	0.599	0.523	0.599	0.604	0.399	0.181	0.604	0.172	-0.178	0.051
Nickel	48 / 48	0.097	0.624	0.394	0.334	0.424	0.565	0.424	0.473	0.657	0.199	0.805	0.438	0.077	0.469
Phosphorous	48 / 48	-0.247	0.233	0.416	-0.092	0.394	0.473	0.394	0.443	0.164	0.168	0.308	0.553	0.090	0.907
Potassium	48 / 48	0.048	-0.016	0.210	0.141	-0.121	0.076	-0.121	-0.076	-0.121	0.076	-0.070	0.130	0.167	0.348
Selenium	48 / 48	-0.157	0.595	0.132	0.038	0.154	0.202	0.215	0.109	0.610	0.326	0.557	0.119	0.050	0.417
Silicon	48 / 48	0.147	0.435	0.192	0.235	0.788	0.840	0.788	0.809	0.680	0.137	0.669	0.207	-0.069	0.210
Sodium	32 / 48	0.635	0.321	0.197	0.394	-0.061	-0.076	-0.061	-0.046	-0.091	-0.504	-0.046	0.033	0.157	0.182
Strontium	48 / 48	0.204	0.662	0.313	0.459	0.394	0.290	0.333	0.443	0.051	-0.137	0.231	0.339	-0.057	0.461
Uranium	48 / 48	-0.113	0.373	0.342	-0.178	0.606	0.534	0.667	0.565	0.728	0.199	0.843	0.479	0.025	0.416
Vanadium	48 / 48	0.131	0.210	0.021	0.106	0.273	0.382	0.212	0.290	0.017	0.229	0.131	0.023	0.174	0.357
Zinc	48 / 48	-0.146	0.638	0.316	0.086	0.394	0.473	0.333	0.443	0.395	0.412	0.563	0.465	0.112	0.731
Mercury (F1)	16 / 18	0.061	0.667	0.424	-0.121	NA	0.779	0.862	0.768	0.673	0.211	0.911	0.290	-0.061	0.273
Mercury (F2)	13 / 18	0.076	0.473	0.412	-0.290	0.779	NA	0.801	0.743	0.653	0.170	0.824	0.092	0.046	0.229
Mercury (F3)	17 / 18	0.061	0.667	0.485	-0.182	0.862	0.801	NA	0.749	0.603	0.131	0.931	0.290	-0.061	0.273
Mercury (F4)	14 / 18	0.107	0.687	0.382	-0.076	0.768	0.743	0.749	NA	0.631	0.208	0.772	0.308	-0.046	0.260
Mercury (F5)	18 / 18	-0.305	0.866	0.455	-0.543	0.673	0.653	0.603	0.631	NA	0.257	0.557	0.076	0.152	0.155
Mercury (F6)	16 / 18	-0.443	0.168	0.290	-0.382	0.211	0.170	0.131	0.208	0.257	NA	0.191	-0.092	-0.046	0.229
Mercury (FS)	18 / 18	-0.311	0.933	0.473	-0.492	0.911	0.824	0.931	0.772	0.557	0.191	NA	0.277	-0.046	0.386
Methylmercury	43 / 48	-0.003	0.089	0.332	0.100	0.290	0.092	0.290	0.308	0.076	-0.092	0.277	NA	-0.078	0.569
Sulfide	45 / 48	0.091	-0.050	-0.089	-0.010	-0.061	0.046	-0.061	-0.046	0.152	-0.046	-0.046	-0.078	NA	0.027
TOC	48 / 48	-0.061	0.145	0.468	0.052	0.273	0.229	0.273	0.260	0.155	0.229	0.386	0.569	0.027	NA

Table E.1. Analyte pairwise correlation matrix for Bear Creek floodplain and creek bank soils (cont.)

	Fred of							Metals							
Chemical	detection	Aluminum	Arsenic	Barium	Beryllium	Boron	Cadmium	Calcium	Chromium	Cobalt	Copper	Iron	Lead	Lithium	Magnesium
Chloride	48 / 48	0.255	0.070	0.092	0.257	0.347	0.029	0.071	0.216	0.205	-0.137	0.473	-0.018	-0.101	-0.002
Fluoride	48 / 48	0.458	0.316	0.452	-0.093	0.003	0.792	0.395	0.142	0.538	0.629	0.113	0.617	0.781	0.145
Nitrate	44 / 48	0.111	0.166	0.264	0.245	0.282	0.433	0.372	0.068	0.220	0.257	0.089	0.212	0.226	0.127
Sulfate	48 / 48	0.343	0.165	0.322	0.217	0.213	0.303	0.355	0.254	0.376	0.023	0.502	0.047	0.088	0.232
Aluminum	48 / 48	NA	0.650	0.421	0.089	0.124	0.551	0.194	0.612	0.590	0.595	99.0	0.681	0.529	0.315
Arsenic	48 / 48	0.650	NA	0.107	-0.026	0.117	0.356	0.153	0.627	0.293	0.591	0.439	0.619	0.530	0.014
Barium	48 / 48	0.421	0.107	NA	0.268	-0.068	0.740	0.335	0.045	0.918	0.346	0.047	0.310	0.331	0.577
Beryllium	46 / 48	0.089	-0.026	0.268	NA	0.270	0.086	0.094	0.094	0.219	0.039	0.150	-0.010	-0.049	0.334
Boron	26 / 48	0.124	0.117	-0.068	0.270	NA	0.134	0.158	0.013	0.056	0.025	0.039	-0.082	0.070	0.158
Cadmium	48 / 48	0.551	0.356	0.740	0.086	0.134	NA	0.417	0.173	0.782	0.675	0.123	0.633	0.754	0.437
Calcium	48 / 48	0.194	0.153	0.335	0.094	0.158	0.417	NA	0.193	0.313	0.338	0.140	0.186	0.419	0.432
Chromium	48 / 48	0.612	0.627	0.045	0.094	0.013	0.173	0.193	NA	0.249	0.272	0.706	0.342	0.324	0.265
Cobalt	48 / 48	0.590	0.293	0.918	0.219	0.056	0.782	0.313	0.249	NA	0.393	0.303	0.432	0.407	0.478
Copper	48 / 48	0.595	0.591	0.346	0.039	0.025	0.675	0.338	0.272	0.393	NA	0.084	0.910	0.828	0.205
Iron	48 / 48	0.665	0.439	0.047	0.150	0.039	0.123	0.140	0.706	0.303	0.084	NA	0.219	0.106	0.062
Lead	48 / 48	0.681	0.619	0.310	-0.010	-0.082	0.633	0.186	0.342	0.432	0.910	0.219	NA	0.769	0.020
Lithium	48 / 48	0.529	0.530	0.331	-0.049	0.070	0.754	0.419	0.324	0.407	0.828	0.106	692.0	NA	0.229
Magnesium	48 / 48	0.315	0.014	0.577	0.334	0.158	0.437	0.432	0.265	0.478	0.205	0.062	0.020	0.229	NA
Manganese	48 / 48	0.364	0.061	0.982	0.164	-0.111	0.724	0.341	0.036	0.918	0.269	0.062	0.235	0.268	0.579
Mercury	48 / 48	0.387	0.437	0.173	-0.021	0.071	0.538	0.428	0.203	0.203	0.862	0.092	0.779	0.746	0.100
Molybdenum	32 / 48	0.217	0.203	0.170	-0.304	-0.102	0.282	990.0	0.039	0.237	0.211	0.102	0.346	0.198	-0.153
Nickel	48 / 48	0.558	0.260	0.929	0.170	0.140	0.914	0.367	0.141	0.919	0.527	0.122	0.484	0.547	0.573
Phosphorous	48 / 48	0.366	0.487	0.424	0.116	0.232	0.529	0.371	0.315	0.375	0.640	-0.078	0.518	0.568	0.509
Potassium	48 / 48	0.505	0.242	0.360	0.353	0.299	0.291	0.113	0.405	0.375	0.225	0.218	0.143	0.173	0.690
Selenium	48 / 48	0.471	0.291	0.717	0.008	-0.131	0.764	0.302	0.197	0.697	0.617	-0.044	0.615	0.589	0.434
Silicon	48 / 48	0.469	0.374	0.215	0.019	0.068	0.372	0.182	0.040	0.307	0.472	0.254	0.501	0.384	-0.027
Sodium	32 / 48	0.306	0.129	0.236	0.324	0.173	0.189	0.355	0.334	0.324	0.130	0.332	0.211	0.141	0.282
Strontium	48 / 48	0.596	0.293	0.773	0.081	0.161	0.798	0.539	0.220	0.756	0.534	0.201	0.448	0.559	0.563
Uranium	48 / 48	0.483	0.510	0.071	-0.052	0.098	0.430	0.013	0.174	0.185	0.707	0.023	0.760	669.0	-0.080
Vanadium	48 / 48	0.706	0.825	-0.025	-0.039	0.020	0.173	0.100	0.794	0.189	0.485	0.635	0.594	0.379	-0.084
Zinc	48 / 48	0.624	0.579	0.540	0.095	0.026	0.784	0.492	0.371	0.549	0.900	0.094	0.783	0.827	0.476
Mercury (F1)	16/18	0.394	0.321	0.364	0.061	0.254	0.485	-0.061	-0.031	0.182	0.576	0.062	0.485	0.424	-0.242
Mercury (F2)	13 / 18	0.382	0.492	0.168	0.107	0.288	0.534	-0.168	0.140	0.351	0.595	0.171	0.718	0.443	-0.290
Mercury (F3)	17 / 18	0.333	0.260	0.364	0.061	0.191	0.485	-0.121	-0.031	0.182	0.515	0.000	0.485	0.364	-0.242
Mercury (F4)	14 / 18	0.412	0.339	0.321	0.076	0.288	0.504	-0.015	-0.016	0.199	0.565	0.078	0.504	0.473	-0.199
Mercury (F5)	18 / 18	0.039	0.109	0.072	0.091	0.350	0.724	-0.318	-0.326	0.354	0.750	-0.226	0.862	0.416	-0.389
Mercury (F6)	16 / 18	0.229	0.339	-0.137	-0.351	-0.144	0.015	-0.351	-0.016	0.015	0.473	-0.016	0.382	0.137	-0.473
Mercury (FS)	18 / 18	0.186	0.185	0.104	0.076	0.208	0.872	-0.247	-0.259	0.237	0.889	-0.196	0.941	0.566	-0.359
Methylmercury	43 / 48	0.068	0.196	0.319	0.081	0.100	0.529	0.324	0.092	0.160	0.383	0.095	0.234	0.394	0.234
Sulfide	45 / 48	0.093	0.034	0.051	0.140	0.189	0.039	-0.056	0.121	0.092	0.152	0.119	0.098	0.057	0.105
TOC	48 / 48	0.337	0.430	0.387	0.092	0.176	0.466	0.385	0.292	0.308	0.598	-0.004	0.494	0.477	0.442
							1								

Table E.1. Analyte pairwise correlation matrix for Bear Creek floodplain and creek bank soils (cont.)

	Freq. of						Me	etals						
Chemical	detection	Manganese	Mercury	Molybdenum	Nickel	Phosphorous	Potassium	Selenium	Silicon	Sodium	Strontium	Uranium	Vanadium	Zinc
Chloride	48 / 48	0.132	-0.097	0.015	0.097	-0.247	0.048	-0.157	0.147	0.635	0.204	-0.113	0.131	-0.146
Fluoride	48 / 48	0.435	0.551	0.350	0.624	0.233	-0.016	0.595	0.435	0.321	0.662	0.373	0.210	0.638
Nitrate	44 / 48	0.226	0.373	0.102	0.394	0.416	0.210	0.132	0.192	0.197	0.313	0.342	0.021	0.316
Sulfate	48 / 48	0.339	0.119	-0.045	0.334	-0.092	0.141	0.038	0.235	0.394	0.459	-0.178	0.106	0.086
Aluminum	48 / 48	0.364	0.387	0.217	0.558	0.366	0.505	0.471	0.469	0.306	0.596	0.483	0.706	0.624
Arsenic	48 / 48	0.061	0.437	0.203	0.260	0.487	0.242	0.291	0.374	0.129	0.293	0.510	0.825	0.579
Barium	48 / 48	0.982	0.173	0.170	0.929	0.424	0.360	0.717	0.215	0.236	0.773	0.071	-0.025	0.540
Beryllium	46 / 48	0.164	-0.021	-0.304	0.170	0.116	0.353	0.008	0.019	0.324	0.081	-0.052	-0.039	0.095
Boron	26 / 48	-0.111	0.071	-0.102	0.140	0.232	0.299	-0.131	0.068	0.173	0.161	0.098	0.020	0.026
Cadmium	48 / 48	0.724	0.538	0.282	0.914	0.529	0.291	0.764	0.372	0.189	0.798	0.430	0.173	0.784
Calcium	48 / 48	0.341	0.428	0.066	0.367	0.371	0.113	0.302	0.182	0.355	0.539	0.013	0.100	0.492
Chromium	48 / 48	0.036	0.203	0.039	0.141	0.315	0.405	0.197	0.040	0.334	0.220	0.174	0.794	0.371
Cobalt	48 / 48	0.918	0.203	0.237	0.919	0.375	0.375	0.697	0.307	0.324	0.756	0.185	0.189	0.549
Copper	48 / 48	0.269	0.862	0.211	0.527	0.640	0.225	0.617	0.472	0.130	0.534	0.707	0.485	0.900
Iron	48 / 48	0.062	0.092	0.102	0.122	-0.078	0.218	-0.044	0.254	0.332	0.201	0.023	0.635	0.094
Lead	48 / 48	0.235	0.779	0.346	0.484	0.518	0.143	0.615	0.501	0.211	0.448	0.760	0.594	0.783
Lithium	48 / 48	0.268	0.746	0.198	0.547	0.568	0.173	0.589	0.384	0.141	0.559	0.699	0.379	0.827
Magnesium	48 / 48	0.579	0.100	-0.153	0.573	0.509	0.690	0.434	-0.027	0.282	0.563	-0.080	-0.084	0.476
Manganese	48 / 48	NA	0.116	0.219	0.917	0.375	0.335	0.692	0.160	0.310	0.757	-0.020	-0.059	0.478
Mercury	48 / 48	0.116	NA	0.378	0.333	0.470	0.055	0.454	0.412	0.104	0.419	0.507	0.335	0.718
Molybdenum	32 / 48	0.219	0.378	NA	0.224	0.112	-0.126	0.146	0.557	-0.120	0.183	0.270	0.164	0.145
Nickel	48 / 48	0.917	0.333	0.224	NA	0.516	0.415	0.765	0.322	0.148	0.843	0.274	0.100	0.704
Phosphorous	48 / 48	0.375	0.470	0.112	0.516	NA	0.447	0.523	0.254	0.227	0.493	0.455	0.342	0.789
Potassium	48 / 48	0.335	0.055	-0.126	0.415	0.447	NA	0.296	0.069	0.115	0.348	0.070	0.205	0.391
Selenium	48 / 48	0.692	0.454	0.146	0.765	0.523	0.296	NA	0.063	0.283	0.658	0.377	0.165	0.713
Silicon	48 / 48	0.160	0.412	0.557	0.322	0.254	0.069	0.063	NA	-0.074	0.379	0.416	0.260	0.395
Sodium	32 / 48	0.310	0.104	-0.120	0.148	0.227	0.115	0.283	-0.074	NA	0.333	0.024	0.187	0.175
Strontium	48 / 48	0.757	0.419	0.183	0.843	0.493	0.348	0.658	0.379	0.333	NA	0.205	0.181	0.682
Uranium	48 / 48	-0.020	0.507	0.270	0.274	0.455	0.070	0.377	0.416	0.024	0.205	NA	0.384	0.566
Vanadium	48 / 48	-0.059	0.335	0.164	0.100	0.342	0.205	0.165	0.260	0.187	0.181	0.384	NA	0.463
Zinc	48 / 48	0.478	0.718	0.145	0.704	0.789	0.391	0.713	0.395	0.175	0.682	0.566	0.463	NA
Mercury (F1)	16 / 18	0.273	0.939	0.599	0.424	0.394	-0.121	0.154	0.788	-0.061	0.394	0.606	0.273	0.394
Mercury (F2)	13 / 18	0.076	0.809	0.523	0.565	0.473	0.076	0.202	0.840	-0.076	0.290	0.534	0.382	0.473
Mercury (F3)	17 / 18	0.273	0.939	0.599	0.424	0.394	-0.121	0.215	0.788	-0.061	0.333	0.667	0.212	0.333
Mercury (F4)	14 / 18	0.229	0.962	0.604	0.473	0.443	-0.076	0.109	0.809	-0.046	0.443	0.565	0.290	0.443
Mercury (F5)	18 / 18	0.215	0.843	0.399	0.657	0.164	-0.121	0.610	0.680	-0.091	0.051	0.728	0.017	0.395
Mercury (F6)	16 / 18	-0.168	0.260	0.181	0.199	0.168	0.076	0.326	0.137	-0.504	-0.137	0.199	0.229	0.412
Mercury (FS)	18 / 18	0.198	0.977	0.604	0.805	0.308	-0.070	0.557	0.669	-0.046	0.231	0.843	0.131	0.563
Methylmercury	43 / 48	0.250	0.546	0.172	0.438	0.553	0.130	0.119	0.207	0.033	0.339	0.479	0.023	0.465
Sulfide	45 / 48	0.032	-0.031	-0.178	0.077	0.090	0.167	0.050	-0.069	0.157	-0.057	0.025	0.174	0.112
TOC	48 / 48	0.347	0.464	0.051	0.469	0.907	0.348	0.417	0.210	0.182	0.461	0.416	0.357	0.731

Table E.1. Analyte pairwise correlation matrix for Bear Creek floodplain and creek bank soils (cont.)

# Correlation is significant at the 0.05 significance level where $0.01 \le \text{two-sided } p\text{-value} \le 0.05$ .

Correlation is significant at the 0.01 significance level where  $0.001 \le \text{two-sided } p\text{-value} \le 0.01$ .

Correlation is significant at the 0.001 significance level where  $0.0001 \le \text{two-sided } p\text{-value} \le 0.001$ .

Pearson correlations are calculated when both variables have all detections.

Kendall tau correlations are calculated when there is at least one non-detection in one or both variables, but at least half detections in each variable.

Freq. = frequency

NA = not applicable

TOC = Total organic carbon average

Table E.2. Correlation matrix for soil analytes for BCV RSE for locations with particle size #10

Character 1	Freq. of		Anions			Other inorganics	Wet cher	nistry			Me	etals		
Chemical	detection	Chloride	Methylmercury	Nitrate	Sulfate	Methylmercury	Sulfide	TOC	Aluminum	Arsenic	Barium	Beryllium	Boron	Cadmium
Chloride	6 / 6	NA	0.647	0.138	0.995	-0.733	0.067	-0.258	0.459	0.138	-0.645	-0.333	-0.507	0.121
Fluoride	6 / 6	0.647	NA	0.414	0.694	-0.200	-0.200	-0.243	0.044	-0.093	-0.309	-0.333	-0.388	-0.257
Nitrate	4 / 6	0.138	0.414	NA	-0.138	0.138	-0.690	0.552	0.000	0.414	0.276	-0.138	-0.276	0.552
Sulfate	6 / 6	0.995	0.694	-0.138	NA	-0.733	0.333	-0.337	0.446	0.072	-0.689	-0.333	-0.479	0.050
Aluminum	6 / 6	0.459	0.044	0.000	0.446	-0.200	0.333	-0.023	NA	0.651	-0.641	-0.600	-0.744	-0.429
Arsenic	6 / 6	0.138	-0.093	0.414	0.072	0.200	-0.067	0.735	0.651	NA	0.138	-0.733	-0.825	-0.064
Barium	6 / 6	-0.645	-0.309	0.276	-0.689	0.733	-0.333	0.755	-0.641	0.138	NA	0.333	0.235	0.405
Beryllium	5 / 6	-0.333	-0.333	-0.138	-0.333	0.067	-0.200	0.067	-0.600	-0.733	0.333	NA	0.867	0.200
Boron	6 / 6	-0.507	-0.388	-0.276	-0.479	-0.067	-0.067	-0.358	-0.744	-0.825	0.235	0.867	NA	0.333
Cadmium	6 / 6	0.121	-0.257	0.552	0.050	0.067	-0.733	0.354	-0.429	-0.064	0.405	0.200	0.333	NA
Calcium	6 / 6	0.818	0.474	0.966	0.764	0.067	-0.733	0.335	0.351	0.503	-0.145	-0.067	-0.621	0.420
Chromium	6 / 6	0.433	-0.045	0.276	0.391	0.333	0.067	0.245	0.953	0.824	-0.429	-0.600	-0.820	-0.231
Cobalt	6 / 6	0.391	-0.229	0.000	0.391	-0.200	0.067	-0.451	0.782	0.155	-0.823	-0.067	-0.240	-0.234
Copper	6 / 6	-0.292	-0.204	0.276	-0.360	0.733	-0.333	0.968	0.108	0.813	0.682	0.333	-0.490	0.116
Iron	6 / 6	0.461	-0.098	0.357	0.460	0.000	0.000	-0.343	0.906	0.327	-0.824	-0.276	-0.435	-0.334
Lead	6 / 6	0.202	-0.008	0.357	0.143	0.138	0.000	0.675	0.688	0.992	0.067	-0.828	-0.887	-0.137
Lithium	6 / 6	-0.055	-0.151	0.414	-0.134	0.067	-0.467	0.919	0.036	0.681	0.610	-0.333	-0.300	0.496
Magnesium	6 / 6	-0.222	0.057	0.000	-0.237	0.200	-0.333	0.226	-0.927	-0.452	0.682	0.600	0.570	0.670
Manganese	6 / 6	-0.332	-0.255	0.414	-0.408	0.600	-0.467	0.988	-0.179	0.622	0.844	0.200	-0.234	0.414
Mercury	6 / 6	-0.270	-0.233	0.414	-0.340	0.733	-0.333	0.968	0.163	0.842	0.645	0.067	-0.499	0.128
Molybdenum	5 / 6	0.733	0.467	0.414	0.467	-0.467	-0.200	0.067	0.467	0.333	-0.200	-0.600	-0.467	-0.067
Nickel	6 / 6	-0.055	-0.180	0.552	-0.142	0.467	-0.600	0.841	-0.313	0.426	0.742	0.067	-0.119	0.751
Phosphorous	6 / 6	-0.551	-0.340	0.276	-0.610	0.733	-0.333	0.939	-0.214	0.554	0.882	0.333	-0.128	0.236
Potassium	6 / 6	0.464	0.329	0.690	0.411	-0.200	-0.733	0.351	-0.366	0.042	0.319	0.200	-0.073	0.780
Selenium	6 / 6	-0.646	-0.114	0.000	-0.667	0.467	-0.067	0.708	-0.587	0.149	0.951	0.067	0.139	0.157
Silicon	6 / 6	0.539	0.078	0.138	0.526	-0.067	0.200	-0.094	0.984	0.599	-0.698	-0.467	-0.751	-0.397
Sodium	3 / 6	0.745	0.894	0.386	0.596	-0.447	-0.149	0.000	0.298	0.149	-0.298	-0.447	-0.298	-0.149
Strontium	6/6	0.577	0.349	0.828	0.511	0.200	-0.600	0.634	0.405	0.769	0.093	-0.200	-0.764	0.298
Thallium	3 / 6	0.183	0.183	0.378	0.000	0.183	-0.183	0.365	0.183	0.730	0.183	-0.730	-0.730	0.000
Uranium	6/6	0.021	-0.325	0.552	-0.059	0.333	-0.733	0.512	-0.401	0.070	0.522	-0.067	0.275	0.982
Vanadium	6 / 6	0.142	-0.120	0.138	0.094	0.200	0.200	0.537	0.817	0.957	-0.086	-0.733	-0.830	-0.275
Zinc	6 / 6	-0.394	-0.275	0.414	-0.464	0.600	-0.467	0.986	-0.110	0.671	0.826	0.200	-0.286	0.281
Methylmercury	5 / 6	-0.733	-0.200	0.138	-0.733	NA	-0.067	0.467	-0.200	0.200	0.733	0.067	-0.067	0.067
Sulfide	5 / 6	0.067	-0.200	-0.690	0.333	-0.067	NA	-0.600	0.333	-0.067	-0.333	-0.200	-0.067	-0.733
TOC	6 / 6	-0.258	-0.243	0.552	-0.337	0.467	-0.600	NA	-0.023	0.735	0.755	0.067	-0.358	0.354

Table E.2. Correlation matrix for soil analytes for BCV RSE for locations with particle size #10 (cont.)

148.0	<i>L</i> 90.0	896.0	886.0	922.0	616.0	£75.0	£4£.0-	896.0	124.0-	6,245	<b>255.0</b>	9/9	TOC
009.0-	002.0-	£££.0-	<i>₹</i> 94.0-	£££.0-	794.0-	000.0	000.0	£££.0-	L90 <sup>.</sup> 0	L90.0	££7.0-	9/5	əpiliu2
<u></u>	L94.0-	6.733	009.0	0.200	L90 <sup>.</sup> 0	8£1.0	000.0	££7.0	002.0-	£££.0	L90 <sup>.</sup> 0	9/5	Methylmercury
108.0	L90 <sup>.</sup> 0-	796.0	686.0	197.0	£98.0	119.0	164.0-	176.0	152.0-	941.0	261.0	9/9	əniS
221.0	£££.0	869.0	204.0	<i>₹</i> 89.0-	764.0	096.0	645.0	659.0	475.0	716.0	£7£.0	9/9	muibene V
188.0	L90 <sup>.</sup> 0-	00£.0	ç9ç.0	059.0	<b>₽</b> £9.0	210.0-	₹9£.0-	987.0	067.0-	<i>₽</i> ∠1.0-	604.0	9/9	muins1U
295.0	842.0	681.0	£9£.0	£81.0-	0.730	9\$ <i>L</i> .0	681.0	681.0	£81.0-	842.0	£9£.0	9/8	muilledT
609.0	£££.0	129.0	<b>SSS.0</b>	LS0.0-	127.0	STT.0	211.0	£09.0	9\$0.0-	£6£.0	816.0	9/9	Strontium
0.000	St7.0	862.0-	641.0-	862.0-	867.0	0.232	000.0	867.0-	867.0-	0.000	<i>L</i> ħħ.0	9/8	muibo2
£1£.0-	£££.0	180.0	942.0-	606.0-	<i>5</i> 20.0-	649.0	076.0	150.0	918.0	046.0	88£.0	9/9	Silicon
482.0	L90 <sup>°</sup> 0	449.0	887.0	£6 <b>2</b> .0	888.0	960.0	9£8.0 <b>-</b>	889.0	888.0-	<i>\</i> 24.0-	261.0-	9/9	Selenium
227.0	0.200	421.0	7e£.0	999.0	0.440	2£0.0	164.0-	S71.0	724.0-	171.0-	22 <i>T</i> .0	9/9	muissatoA
427.0	002.0-	<b>4</b> 16.0	686.0	015.0	818.0	784.0	802.0-	<b>₽</b> 76.0	£6 <b>2</b> .0-	610.0	710.0	9/9	Phosphorous
٧N	L90 <sup>.</sup> 0	769.0	978.0	LLS.0	708.0	89£.0	764.0-	<i>₽</i> 07.0	912.0-	910.0-	L84.0	9/9	Nickel
790.0	٧N	-0.200	L90 <sup>.</sup> 0-	L9t <sup>.</sup> 0-	L9t <sup>.</sup> 0	414.0	972.0	002.0-	۷90 <sup>.</sup> 0-	0.200	L94.0	9/5	Molybdenum
769.0	002.0-	٧N	826.0	100.0	<i>L</i> \$8.0	96 <i>L</i> .0	161.0-	966.0	188.0-	86£.0	<i>LL</i> 7.0	9/9	Mercury
978.0	L90 <sup>.</sup> 0-	826.0	٧N	£9£.0	968.0	<b>7</b> \$\$.0	874.0-	7£6.0	<i></i> ₽9 <b>5</b> .0-	£60.0	L97.0	9/9	Manganese
rre.0	L94.0-	100.0	295.0	VΝ	0.240	∠6t <sup>.</sup> 0-	968.0-	940.0	808.0-	<b>≿</b> 08.0-	110.0	9/9	Magnesium
708.0	L9t <sup>.</sup> 0	7 <b>č</b> 8.0	968.0	040.0	ΨN	£09.0	8£2.0-	££8.0	985.0-	S72.0	902.0	9/9	muidii.1
89£.0	t1t.0	96Ľ.0	<b>722.0</b>	<i>L</i> 6⊅ <sup>.</sup> 0−	£09.0	ΨN	825.0	277.0	6LI.0	948.0	0.522	9/9	Lead
∠6t <sup>·</sup> 0-	972.0	191.0-	874.0-	968.0-	8£2.0-	82£.0	VΝ	-0.255	696.0	86 <i>L</i> .0	781.0	9/9	Iron
407.0	002.0-	966'0	7£6.0	940.0	££8.0	2 <i>LL</i> .0	522.0-	ΨN	765.0-	945.0	952.0	9/9	Copper
912.0-	L90 <sup>.</sup> 0-	188.0-	<i></i> ₽9 <i>S</i> ·0-	808.0-	9 <b>2</b> £.0-	671.0	696.0	76£.0-	ΑN	999.0	170.0	9/9	Cobalt
910.0-	0.200	86£.0	£60.0	₹08.0-	SLZ.0	948.0	867.0	946.0	999.0	ΨN	064.0	9/9	Chromium
784.0	L9t <sup>.</sup> 0	<i>LL</i> 7.0	L97.0	110.0	902.0	0.522	781.0	922.0	170.0	064.0	٧N	9/9	Calcium
127.0	L90 <sup>.</sup> 0-	821.0	414.0	076.0	964.0	781.0-	466.0-	911.0	<b>₽</b> £2.0-	162.0-	0.420	9/9	Cadmium
911.0-	L9t <sup>.</sup> 0-	664.0-	-0.234	07 <b>2.</b> 0	00£.0-	∠88 <sup>.</sup> 0-	2£4.0-	064.0-	042.0-	028.0-	129.0-	9/9	Boron
790.0	009.0-	<i>L</i> 90'0	0.200	009.0	£££.0-	828.0-	972.0-	£££.0	۲90 <sup>.</sup> 0-	009.0-	۷90 <sup>.</sup> 0-	9/5	Beryllium
<u>7</u> ₽₽	002.0-	S49.0	<i>4</i> 48.0	789.0	019.0	<i>L</i> 90'0	<b>₽</b> 28.0-	789.0	628.0-	624.0-	541.0-	9/9	Barium
924.0	£££.0	7 <del>+</del> 8.0	229.0	754.0-	189.0	766.0	72£.0	£18.0	\$\$1.0	<del>7</del> 28.0	£0 <b>2</b> .0	9/9	əinəs1A
£1£.0-	794.0	691.0	671.0-	726.0-	9£0.0	889.0	906.0	801.0	287.0	£26.0	125.0	9/9	munimulA
-0.142	L9t <sup>.</sup> 0	045.0-	804.0-	7£2.0-	4£1.0-	6,143	094.0	09£.0-	16£.0	16£.0	<del>7</del> 9∠'0	9/9	Sulfate
252.0	0.414	414.0	414.0	000.0	414.0	72£.0	72£.0	972.0	000.0	972.0	996:0	9/1	Nitrate
081.0-	794.0	££2.0-	-0.255	720.0	121.0-	800.0-	860.0-	<b>₽</b> 02.0-	622.0-	<i>≿</i> ₽0.0-	<i>ኮ</i> /ኒታ <sup>.</sup> 0	9/9	Fluoride
\$\$0.0-	££7.0	072.0-	266.0-	222.0-	\$\$0.0-	202.0	194.0	262.0-	16£.0	654.0	818.0	9/9	Chloride
Nickel	Molybdenum	Mercury	Manganese	muisəngaM	Lithium	Геяд	Iron	Copper	Cobalt	Сһтотіит	Calcium	noitootob	[kəimədə
					sls	њМ						Freq. of	[ooimod)

Table E.2. Correlation matrix for soil analytes for BCV RSE for locations with particle size #10 (cont.)

G1 1 1	Freq. of					Metals					
Chemical	detection	Phosphorous	Potassium	Selenium	Silicon	Sodium	Strontium	Thallium	Uranium	Vanadium	Zinc
Chloride	6 / 6	-0.551	0.464	-0.646	0.539	0.745	0.577	0.183	0.021	0.142	-0.394
Fluoride	6 / 6	-0.340	0.329	-0.114	0.078	0.894	0.349	0.183	-0.325	-0.120	-0.275
Nitrate	4 / 6	0.276	0.690	0.000	0.138	0.386	0.828	0.378	0.552	0.138	0.414
Sulfate	6 / 6	-0.610	0.411	-0.667	0.526	0.596	0.511	0.000	-0.059	0.094	-0.464
Aluminum	6 / 6	-0.214	-0.366	-0.587	0.984	0.298	0.405	0.183	-0.401	0.817	-0.110
Arsenic	6 / 6	0.554	0.042	0.149	0.599	0.149	0.769	0.730	0.070	0.957	0.671
Barium	6 / 6	0.882	0.319	0.951	-0.698	-0.298	0.093	0.183	0.522	-0.086	0.826
Beryllium	5 / 6	0.333	0.200	0.067	-0.467	-0.447	-0.200	-0.730	-0.067	-0.733	0.200
Boron	6 / 6	-0.128	-0.073	0.139	-0.751	-0.298	-0.764	-0.730	0.275	-0.830	-0.286
Cadmium	6 / 6	0.236	0.780	0.157	-0.397	-0.149	0.298	0.000	0.982	-0.275	0.281
Calcium	6 / 6	0.017	0.725	-0.192	0.388	0.447	0.918	0.365	0.409	0.373	0.192
Chromium	6 / 6	0.013	-0.171	-0.427	0.940	0.000	0.595	0.548	-0.174	0.917	0.146
Cobalt	6 / 6	-0.593	-0.427	-0.888	0.816	-0.298	-0.056	-0.183	-0.290	0.374	-0.531
Copper	6 / 6	0.924	0.175	0.688	0.031	-0.298	0.603	0.183	0.286	0.659	0.971
Iron	6 / 6	-0.508	-0.431	-0.836	0.920	0.000	0.112	0.189	-0.365	0.543	-0.431
Lead	6 / 6	0.482	0.035	0.096	0.649	0.232	0.775	0.756	-0.015	0.960	0.611
Lithium	6 / 6	0.818	0.440	0.558	-0.055	0.298	0.721	0.730	0.634	0.492	0.863
Magnesium	6 / 6	0.310	0.666	0.593	-0.909	-0.298	-0.057	-0.183	0.650	-0.687	0.261
Manganese	6 / 6	0.959	0.397	0.788	-0.246	-0.149	0.555	0.365	0.565	0.402	0.989
Mercury	6 / 6	0.914	0.154	0.644	0.081	-0.298	0.621	0.183	0.300	0.698	0.962
Molybdenum	5 / 6	-0.200	0.200	0.067	0.333	0.745	0.333	0.548	-0.067	0.333	-0.067
Nickel	6 / 6	0.724	0.752	0.584	-0.313	0.000	0.609	0.365	0.831	0.155	0.801
Phosphorous	6 / 6	NA	0.157	0.872	-0.311	-0.298	0.356	0.183	0.407	0.377	0.978
Potassium	6 / 6	0.157	NA	0.186	-0.296	0.149	0.597	0.000	0.741	-0.211	0.271
Selenium	6 / 6	0.872	0.186	NA	-0.668	0.000	0.083	0.183	0.285	-0.043	0.797
Silicon	6 / 6	-0.311	-0.296	-0.668	NA	0.149	0.400	0.183	-0.394	0.761	-0.186
Sodium	3 / 6	-0.298	0.149	0.000	0.149	NA	0.298	0.306	-0.149	0.149	-0.149
Strontium	6 / 6	0.356	0.597	0.083	0.400	0.298	NA	0.548	0.359	0.622	0.518
Thallium	3 / 6	0.183	0.000	0.183	0.183	0.306	0.548	NA	0.365	0.548	0.365
Uranium	6 / 6	0.407	0.741	0.285	-0.394	-0.149	0.359	0.365	NA	-0.153	0.445
Vanadium	6 / 6	0.377	-0.211	-0.043	0.761	0.149	0.622	0.548	-0.153	NA	0.478
Zinc	6 / 6	0.978	0.271	0.797	-0.186	-0.149	0.518	0.365	0.445	0.478	NA
Methylmercury	5 / 6	0.733	-0.200	0.467	-0.067	-0.447	0.200	0.183	0.333	0.200	0.600
Sulfide	5 / 6	-0.333	-0.733	-0.067	0.200	-0.149	-0.600	-0.183	-0.733	0.200	-0.467
TOC	6 / 6	0.939	0.351	0.708	-0.094	0.000	0.634	0.365	0.512	0.537	0.986

Correlation is significant at the 0.05 significance level where  $0.01 \le \text{two-sided } p\text{-value} \le 0.05$ .

Correlation is significant at the 0.01 significance level where  $0.001 \le \text{two-sided } p\text{-value} \le 0.01$ .

Correlation is significant at the 0.001 significance level where  $0.0001 \le \text{two-sided } p\text{-value} \le 0.001$ .

Correlation is significant at the 0.0001 significance level where two-sided *p*-Pearson correlations are calculated when both variables have all detections.

Kendall tau correlations are calculated when there is at least one non-detection in one or both variables, but at least half detections in each variable.

BCV = Bear Creek Valley

Freq. = frequency
NA = not applicable
RSE = remedial site evaluation

TOC = Total organic carbon aver

Table E.3. Analyte pairwise correlation matrix for Bear Creek channel sediment

	F 6		Anior	ns					Sequentia	al extraction				Other inorganics	Wet che	mistry
Chemical	Freq. of detection	Chloride	Fluoride	Nitrate	Sulfate	Mercury (F0)	Mercury (F1)	Mercury (F2)	Mercury (F3)	Mercury (F4)	Mercury (F5)	Mercury (F6)	Mercury (FS)	Methylmercury	Sulfide	TOC
Chloride	12 / 12	NA	0.615	0.094	-0.090	0.653	0.067	0.067	-0.200	0.467	0.588	0.200	0.069	0.165	-0.318	0.043
Fluoride	12 / 12	0.615	NA	0.351	0.387	0.653	0.067	0.067	-0.200	0.467	0.211	0.200	-0.224	-0.253	-0.171	0.321
Nitrate	10 / 12	0.094	0.351	NA	-0.015	0.159	-0.031	0.000	-0.092	0.061	0.122	0.061	0.000	-0.565	-0.063	-0.137
Sulfate	12 / 12	-0.090	0.387	-0.015	NA	0.000	-0.200	-0.467	0.067	0.467	-0.565	-0.333	0.009	0.228	0.388	0.453
Aluminum	12 / 12	0.179	0.737	0.412	0.523	0.653	-0.467	-0.467	-0.467	0.200	0.048	-0.067	-0.159	-0.353	0.109	0.650
Arsenic	12 / 12	0.389	0.022	-0.046	-0.486	0.163	-0.067	-0.067	-0.067	-0.200	0.261	0.333	-0.045	-0.260	-0.109	-0.214
Barium	12 / 12	0.212	0.610	0.443	0.307	0.653	-0.067	-0.067	-0.333	0.333	0.540	0.067	-0.046	-0.288	-0.078	0.823
Beryllium	12 / 12	-0.304	0.294	0.321	0.474	0.064	-0.067	-0.333	-0.067	0.067	-0.198	0.333	-0.141	-0.358	0.295	0.658
Boron	8 / 12	-0.277	-0.032	0.272	0.381	0.163	-0.061	-0.122	-0.061	0.031	-0.031	0.031	-0.031	-0.381	0.358	0.064
Cadmium	12 / 12	-0.005	0.501	0.382	0.359	0.653	-0.067	-0.067	-0.333	0.333	0.495	0.067	-0.069	-0.283	-0.047	0.873
Calcium	12 / 12	0.603	0.572	0.382	-0.018	0.490	0.200	0.200	-0.067	0.067	0.598	0.600	0.066	-0.047	-0.326	0.594
Chromium	12 / 12	0.058	-0.079	-0.015	-0.057	0.327	-0.200	-0.200	-0.200	-0.067	0.122	0.200	-0.271	-0.311	-0.047	-0.288
Cobalt	12 / 12	0.167	0.661	0.473	0.392	0.490	-0.200	-0.200	-0.200	0.467	0.356	-0.067	-0.166	-0.352	-0.047	0.792
Copper	12 / 12	0.373	0.688	0.504	0.371	0.490	0.200	0.200	-0.067	0.600	0.449	0.067	0.311	-0.170	-0.078	0.744
Iron	12 / 12	0.626	0.607	0.351	-0.019	0.653	-0.067	-0.067	-0.333	0.333	0.433	0.067	-0.221	-0.388	-0.295	-0.132
Lead	12 / 12	0.739	0.758	0.382	0.089	0.653	-0.067	-0.067	-0.333	0.333	0.535	0.067	-0.026	-0.262	-0.140	0.330
Lithium	12 / 12	0.026	0.308	0.202	0.359	0.000	0.414	0.138	0.276	0.414	-0.165	0.552	0.312	0.043	0.236	0.749
Magnesium	12 / 12	-0.156	0.046	0.369	0.024	0.163	0.200	-0.067	-0.067	0.067	0.163	0.600	-0.154	-0.275	0.094	0.479
Manganese	12 / 12	0.066	0.530	0.504	0.336	0.653	-0.067	-0.067	-0.333	0.333	0.512	0.067	-0.143	-0.301	-0.140	0.855
Mercury	12 / 12	0.593	0.309	-0.046	0.052	-0.490	1.000	0.733	0.733	0.333	0.367	0.600	0.983	0.537	-0.140	0.630
Molybdenum	6 / 12	-0.282	-0.121	0.434	-0.086	0.738	-0.258	-0.258	-0.603	-0.086	0.430	0.258	-0.086	-0.362	-0.088	-0.155
Nickel	12 / 12	-0.053	0.463	0.473	0.365	0.653	-0.067	-0.067	-0.333	0.333	0.494	0.067	0.041	-0.295	-0.078	0.864
Phosphorous	12 / 12	-0.371	-0.449	0.076	-0.166	-0.163	0.200	0.200	0.200	-0.200	-0.024	0.333	-0.081	-0.321	0.140	-0.107
Potassium	12 / 12	-0.068	0.528	0.565	0.514	0.490	-0.333	-0.600	-0.333	0.333	-0.311	0.067	-0.434	-0.492	0.295	0.317
Selenium	12 / 12	0.588	0.631	0.382	-0.050	0.327	0.200	0.200	-0.067	0.067	0.852	0.333	0.103	-0.269	-0.202	0.598
Silicon	12 / 12	-0.342	0.315	0.443	0.668	0.653	-0.200	-0.467	-0.467	0.200	-0.283	0.200	-0.413	-0.285	0.140	0.206
Sodium	10 / 12	0.594	0.748	0.215	0.199	0.327	0.200	-0.067	-0.067	0.600	0.067	0.333	0.333	-0.015	0.000	0.382
Strontium	12 / 12	0.531	0.587	0.473	-0.132	0.327	0.200	0.200	-0.067	0.067	0.880	0.333	0.155	-0.321	-0.233	0.462
Uranium	12 / 12	0.633	0.619	0.000	0.339	0.163	-0.067	-0.333	-0.067	0.333	-0.237	0.067	-0.158	0.107	0.109	0.106
Vanadium	12 / 12	0.249	0.070	0.015	-0.321	0.163	-0.067	-0.067	-0.067	-0.200	0.412	0.333	-0.068	-0.388	-0.109	-0.346
Zinc	12 / 12	0.308	0.478	0.321	0.052	0.327	-0.067	-0.067	-0.067	0.333	0.913	0.067	0.486	-0.216	-0.171	0.582
Mercury (F0)	3 / 6	0.653	0.653	0.159	0.000	NA	-0.490	-0.327	-0.817	0.000	0.327	0.000	-0.327	-0.817	-0.254	-0.653
Mercury (F1)	5 / 6	0.067	0.067	-0.031	-0.200	-0.490	NA	0.733	0.733	0.333	0.333	0.600	0.867	0.467	-0.138	0.600
Mercury (F2)	5 / 6	0.067	0.067	0.000	-0.467	-0.327	0.733	NA	0.467	0.067	0.333	0.333	0.600	0.467	-0.414	0.600
Mercury (F3)	5 / 6	-0.200	-0.200	-0.092	0.067	-0.817	0.733	0.467	NA	0.333	0.067	0.333	0.600	0.733	0.138	0.600
Mercury (F4)	5 / 6	0.467	0.467	0.061	0.467	0.000	0.333	0.067	0.333	NA	0.200	0.200	0.467	0.067	0.276	-0.067
Mercury (F5)	6 / 6	0.588	0.211	0.122	-0.565	0.327	0.333	0.333	0.067	0.200	NA	0.467	0.478	-0.214	-0.552	-0.026
Mercury (F6)	5 / 6	0.200	0.200	0.061	-0.333	0.000	0.600	0.333	0.333	0.200	0.467	NA	0.733	0.067	-0.138	0.200
Mercury (FS)	6 / 6	0.069	-0.224	0.000	0.009	-0.327	0.867	0.600	0.600	0.467	0.478	0.733	NA	0.728	0.000	0.753
Methylmercury	12 / 12	0.165	-0.253	-0.565	0.228	-0.817	0.467	0.467	0.733	0.067	-0.214	0.067	0.728	NA	-0.264	0.160
Sulfide	9 / 12	-0.318	-0.171	-0.063	0.388	-0.254	-0.138	-0.414	0.138	0.276	-0.552	-0.138	0.000	-0.264	NA	-0.078
TOC	12 / 12	0.043	0.321	-0.137	0.453	-0.653	0.600	0.600	0.600	-0.067	-0.026	0.200	0.753	0.160	-0.078	NA

Table E.3. Analyte pairwise correlation matrix for Bear Creek channel sediment (cont.)

	Freq. of detection Alice Description Chicago C														
Chemical	detection	Aluminum	Arsenic	Barium	Beryllium	Boron	Cadmium	Calcium	Chromium	Cobalt	Copper	Iron	Lead	Lithium	Magnesium
Chloride	12 / 12	0.179	0.389	0.212	-0.304	-0.277	-0.005	0.603	0.058	0.167	0.373	0.626	0.739	0.026	-0.156
Fluoride	12 / 12	0.737	0.022	0.610	0.294	-0.032	0.501	0.572	-0.079	0.661	0.688	0.607	0.758	0.308	0.046
Nitrate	10 / 12	0.412	-0.046	0.443	0.321	0.272	0.382	0.382	-0.015	0.473	0.504	0.351	0.382	0.202	0.369
Sulfate	12 / 12	0.523	-0.486	0.307	0.474	0.381	0.359	-0.018	-0.057	0.392	0.371	-0.019	0.089	0.359	0.024
Aluminum	12 / 12	NA	0.120	0.874	0.842	0.350	0.859	0.541	0.104	0.943	0.902	0.440	0.702	0.733	0.405
Arsenic	12 / 12	0.120	NA	0.044	0.022	-0.064	-0.063	0.254	0.597	0.054	0.137	0.544	0.483	0.174	0.102
Barium	12 / 12	0.874	0.044	NA	0.729	0.127	0.963	0.640	-0.138	0.973	0.929	0.231	0.664	0.713	0.410
Beryllium	12 / 12	0.842	0.022	0.729	NA	0.826	0.824	0.331	0.195	0.816	0.708	0.151	0.344	0.751	0.630
Boron	8 / 12	0.350	-0.064	0.127	0.826	NA	0.095	0.032	0.286	0.286	0.064	0.032	0.127	0.226	0.432
Cadmium	12 / 12	0.859	-0.063	0.963	0.824	0.095	NA	0.568	-0.195	0.960	0.850	0.064	0.483	0.767	0.494
Calcium	12 / 12	0.541	0.254	0.640	0.331	0.032	0.568	NA	-0.049	0.617	0.716	0.430	0.717	0.587	0.630
Chromium	12 / 12	0.104	0.597	-0.138	0.195	0.286	-0.195	-0.049	NA	-0.010	0.040	0.670	0.326	-0.141	0.182
Cobalt	12 / 12	0.943	0.054	0.973	0.816	0.286	0.960	0.617	-0.010	NA	0.928	0.328	0.677	0.705	0.449
Copper	12 / 12	0.902	0.137	0.929	0.708	0.064	0.850	0.716	0.040	0.928	NA	0.457	0.820	0.719	0.432
Iron	12 / 12	0.440	0.544	0.231	0.151	0.032	0.064	0.430	0.670	0.328	0.457	NA	0.830	-0.017	0.125
Lead	12 / 12	0.702	0.483	0.664	0.344	0.127	0.483	0.717	0.326	0.677	0.820	0.830	NA	0.386	0.224
Lithium	12 / 12	0.733	0.174	0.713	0.751	0.226	0.767	0.587	-0.141	0.705	0.719	-0.017	0.386	NA	0.586
Magnesium	12 / 12	0.405	0.102	0.410	0.630	0.432	0.494	0.630	0.182	0.449	0.432	0.125	0.224	0.586	NA
Manganese	12 / 12	0.867	-0.018	0.987	0.798	0.127	0.993	0.584	-0.158	0.972	0.885	0.129	0.555	0.742	0.464
Mercury	12 / 12	0.300	0.147	0.503	0.076	-0.159	0.420	0.648	-0.308	0.407	0.570	0.062	0.492	0.524	0.059
Molybdenum	6 / 12	0.052	0.052	0.017	0.569	0.452	-0.017	0.017	0.224	0.121	0.052	0.190	0.017	-0.053	0.417
Nickel	12 / 12	0.857	-0.061	0.954	0.849	0.191	0.997	0.525	-0.159	0.957	0.840	0.058	0.464	0.758	0.490
Phosphorous	12 / 12	0.047	0.549	-0.055	0.369	0.350	-0.021	-0.204	0.812	0.019	-0.022	0.216	0.018	0.057	0.298
Potassium	12 / 12	0.842	0.108	0.611	0.793	0.509	0.616	0.252	0.175	0.684	0.648	0.341	0.471	0.640	0.405
Selenium	12 / 12	0.704	0.405	0.830	0.453	0.064	0.714	0.857	0.093	0.804	0.866	0.565	0.886	0.549	0.423
Silicon	12 / 12	0.524	-0.357	0.206	0.660	0.572	0.317	0.057	0.250	0.375	0.289	0.218	0.073	0.258	0.468
Sodium	10 / 12	0.565	0.199	0.657	0.015	-0.016	0.687	0.657	-0.076	0.473	0.595	0.229	0.565	0.605	0.417
Strontium	12 / 12	0.582	0.358	0.716	0.326	0.000	0.587	0.822	0.004	0.661	0.769	0.494	0.790	0.488	0.475
Uranium	12 / 12	0.479	0.475	0.221	0.125	0.127	0.128	0.384	0.231	0.285	0.366	0.523	0.573	0.389	-0.059
Vanadium	12 / 12	0.165	0.801	-0.051	0.142	0.127	-0.147	0.122	0.882	0.050	0.148	0.788	0.508	-0.055	0.149
Zinc	12 / 12	0.620	0.273	0.695	0.456	0.191	0.654	0.586	0.112	0.721	0.716	0.409	0.654	0.429	0.279
Mercury (F0)	3 / 6	0.653	0.163	0.653	0.163	0.064	0.653	0.490	0.327	0.490	0.490	0.653	0.653	0.000	0.163
Mercury (F1)	5 / 6	-0.467	-0.067	-0.067	-0.067	-0.061	-0.067	0.200	-0.200	-0.200	0.200	-0.067	-0.067	0.414	0.200
Mercury (F2)	5 / 6	-0.467	-0.067	-0.067	-0.333	-0.122	-0.067	0.200	-0.200	-0.200	0.200	-0.067	-0.067	0.138	-0.067
Mercury (F3)	5 / 6	-0.467	-0.067	-0.333	-0.067	-0.061	-0.333	-0.067	-0.200	-0.200	-0.067	-0.333	-0.333	0.276	-0.067
Mercury (F4)	5 / 6	0.200	-0.200	0.333	0.067	0.031	0.333	0.067	-0.067	0.467	0.600	0.333	0.333	0.414	0.067
Mercury (F5)	6 / 6	0.048	0.261	0.540	-0.198	-0.031	0.495	0.598	0.122	0.356	0.449	0.433	0.535	-0.165	0.163
Mercury (F6)	5 / 6	-0.067	0.333	0.067	0.333	0.031	0.067	0.600	0.200	-0.067	0.067	0.067	0.067	0.552	0.600
Mercury (FS)	6 / 6	-0.159	-0.045	-0.046	-0.141	-0.031	-0.069	0.066	-0.271	-0.166	0.311	-0.221	-0.026	0.312	-0.154
Methylmercury	12 / 12	-0.353	-0.260	-0.288	-0.358	-0.381	-0.283	-0.047	-0.311	-0.352	-0.170	-0.388	-0.262	0.043	-0.275
Sulfide	9 / 12	0.109	-0.109	-0.078	0.295	0.358	-0.047	-0.326	-0.047	-0.047	-0.078	-0.295	-0.140	0.236	0.094
TOC	12 / 12	0.650	-0.214	0.823	0.658	0.064	0.873	0.594	-0.288	0.792	0.744	-0.132	0.330	0.749	0.479
	12/12	0.050	0.217	0.025	0.050	0.004	0.075	0.374	0.200	0.172	0.744	0.152	0.550	0.749	0.477

Table E.3. Analyte pairwise correlation matrix for Bear Creek channel sediment (cont.)

Character 1	Freq. of		Metals													
Chemical	detection	Manganese	Mercury	Molybdenum	Nickel	Phosphorous	Potassium	Selenium	Silicon	Sodium	Strontium	Uranium	Vanadium	Zinc		
Chloride	12 / 12	0.066	0.593	-0.282	-0.053	-0.371	-0.068	0.588	-0.342	0.594	0.531	0.633	0.249	0.308		
Fluoride	12 / 12	0.530	0.309	-0.121	0.463	-0.449	0.528	0.631	0.315	0.748	0.587	0.619	0.070	0.478		
Nitrate	10 / 12	0.504	-0.046	0.434	0.473	0.076	0.565	0.382	0.443	0.215	0.473	0.000	0.015	0.321		
Sulfate	12 / 12	0.336	0.052	-0.086	0.365	-0.166	0.514	-0.050	0.668	0.199	-0.132	0.339	-0.321	0.052		
Aluminum	12 / 12	0.867	0.300	0.052	0.857	0.047	0.842	0.704	0.524	0.565	0.582	0.479	0.165	0.620		
Arsenic	12 / 12	-0.018	0.147	0.052	-0.061	0.549	0.108	0.405	-0.357	0.199	0.358	0.475	0.801	0.273		
Barium	12 / 12	0.987	0.503	0.017	0.954	-0.055	0.611	0.830	0.206	0.657	0.716	0.221	-0.051	0.695		
Beryllium	12 / 12	0.798	0.076	0.569	0.849	0.369	0.793	0.453	0.660	0.015	0.326	0.125	0.142	0.456		
Boron	8 / 12	0.127	-0.159	0.452	0.191	0.350	0.509	0.064	0.572	-0.016	0.000	0.127	0.127	0.191		
Cadmium	12 / 12	0.993	0.420	-0.017	0.997	-0.021	0.616	0.714	0.317	0.687	0.587	0.128	-0.147	0.654		
Calcium	12 / 12	0.584	0.648	0.017	0.525	-0.204	0.252	0.857	0.057	0.657	0.822	0.384	0.122	0.586		
Chromium	12 / 12	-0.158	-0.308	0.224	-0.159	0.812	0.175	0.093	0.250	-0.076	0.004	0.231	0.882	0.112		
Cobalt	12 / 12	0.972	0.407	0.121	0.957	0.019	0.684	0.804	0.375	0.473	0.661	0.285	0.050	0.721		
Copper	12 / 12	0.885	0.570	0.052	0.840	-0.022	0.648	0.866	0.289	0.595	0.769	0.366	0.148	0.716		
Iron	12 / 12	0.129	0.062	0.190	0.058	0.216	0.341	0.565	0.218	0.229	0.494	0.523	0.788	0.409		
Lead	12 / 12	0.555	0.492	0.017	0.464	0.018	0.471	0.886	0.073	0.565	0.790	0.573	0.508	0.654		
Lithium	12 / 12	0.742	0.524	-0.053	0.758	0.057	0.640	0.549	0.258	0.605	0.488	0.389	-0.055	0.429		
Magnesium	12 / 12	0.464	0.059	0.417	0.490	0.298	0.405	0.423	0.468	0.308	0.475	-0.059	0.149	0.279		
Manganese	12 / 12	NA	0.441	0.086	0.989	-0.008	0.622	0.762	0.273	0.595	0.638	0.146	-0.101	0.669		
Mercury	12 / 12	0.441	NA	-0.328	0.391	-0.327	-0.092	0.635	-0.377	0.657	0.477	0.343	-0.124	0.501		
Molybdenum	6 / 12	0.086	-0.328	NA	0.086	0.362	0.328	0.086	0.534	-0.174	0.086	-0.259	0.155	0.017		
Nickel	12 / 12	0.989	0.391	0.086	NA	0.034	0.623	0.692	0.334	0.534	0.555	0.097	-0.122	0.653		
Phosphorous	12 / 12	-0.008	-0.327	0.362	0.034	NA	0.161	-0.009	0.122	-0.107	-0.074	-0.086	0.675	0.098		
Potassium	12 / 12	0.622	-0.092	0.328	0.623	0.161	NA	0.371	0.650	0.229	0.358	0.381	0.191	0.222		
Selenium	12 / 12	0.762	0.635	0.086	0.692	-0.009	0.371	NA	-0.037	0.565	0.884	0.353	0.303	0.794		
Silicon	12 / 12	0.273	-0.377	0.534	0.334	0.122	0.650	-0.037	NA	-0.015	-0.065	0.086	0.056	0.007		
Sodium	10 / 12	0.595	0.657	-0.174	0.534	-0.107	0.229	0.565	-0.015	NA	0.443	0.534	-0.107	0.382		
Strontium	12 / 12	0.638	0.477	0.086	0.555	-0.074	0.358	0.884	-0.065	0.443	NA	0.250	0.228	0.732		
Uranium	12 / 12	0.146	0.343	-0.259	0.097	-0.086	0.381	0.353	0.086	0.534	0.250	NA	0.304	0.185		
Vanadium	12 / 12	-0.101	-0.124	0.155	-0.122	0.675	0.191	0.303	0.056	-0.107	0.228	0.304	NA	0.258		
Zinc	12 / 12	0.669	0.501	0.017	0.653	0.098	0.222	0.794	0.007	0.382	0.732	0.185	0.258	NA		
Mercury (F0)	3 / 6	0.653	-0.490	0.738	0.653	-0.163	0.490	0.327	0.653	0.327	0.327	0.163	0.163	0.327		
Mercury (F1)	5 / 6	-0.067	1.000	-0.258	-0.067	0.200	-0.333	0.200	-0.200	0.200	0.200	-0.067	-0.067	-0.067		
Mercury (F2)	5 / 6	-0.067	0.733	-0.258	-0.067	0.200	-0.600	0.200	-0.467	-0.067	0.200	-0.333	-0.067	-0.067		
Mercury (F3)	5 / 6	-0.333	0.733	-0.603	-0.333	0.200	-0.333	-0.067	-0.467	-0.067	-0.067	-0.067	-0.067	-0.067		
Mercury (F4)	5 / 6	0.333	0.333	-0.086	0.333	-0.200	0.333	0.067	0.200	0.600	0.067	0.333	-0.200	0.333		
Mercury (F5)	6 / 6	0.512	0.367	0.430	0.494	-0.024	-0.311	0.852	-0.283	0.067	0.880	-0.237	0.412	0.913		
Mercury (F6)	5 / 6	0.067	0.600	0.258	0.067	0.333	0.067	0.333	0.200	0.333	0.333	0.067	0.333	0.067		
Mercury (FS)	6 / 6	-0.143	0.983	-0.086	0.041	-0.081	-0.434	0.103	-0.413	0.333	0.155	-0.158	-0.068	0.486		
Methylmercury	12 / 12	-0.301	0.537	-0.362	-0.295	-0.321	-0.492	-0.269	-0.285	-0.015	-0.321	0.107	-0.388	-0.216		
Sulfide	9 / 12	-0.140	-0.140	-0.088	-0.078	0.140	0.295	-0.202	0.140	0.000	-0.233	0.109	-0.109	-0.171		
TOC	12 / 12	0.855	0.630	-0.155	0.864	-0.107	0.317	0.598	0.206	0.382	0.462	0.106	-0.346	0.582		

# Correlation is significant at the 0.05 significance level where $0.01 \le \text{two-sided } p\text{-value} \le 0.05$ .

Correlation is significant at the 0.01 significance level where  $0.001 \le \text{two-sided } p\text{-value} \le 0.01$ .

Correlation is significant at the 0.001 significance level where  $0.0001 \le \text{two-sided } p\text{-value} \le 0.001$ .

Pearson correlations are calculated when both variables have all detections.

Kendall tau correlations are calculated when there is at least one non-detection in one or both variables, but at least half detections in each variable.

Freq. = frequency
NA = not applicable
TOC = Total organic carbon average

Table E.4. Correlation matrix for sediment analytes for BCV RSE for locations with particle size #10

	E of		Anio	ons					Sequentia	al extraction				Other inorganics	Wet chemistry	
Chemical	Freq. of detection	Chloride	Fluoride	Nitrate	Sulfate	Mercury (F0)	Mercury (F1)	Mercury (F2)	Mercury (F3)	Mercury (F4)	Mercury (F5)	Mercury (F6)	Mercury (FS)	Methylmercury	Sulfide	тос
Chloride	7 / 7	NA	0.783	0.000	0.347	0.522	-0.333	-0.333	-0.333	0.959	0.467	-0.334	0.057	0.190	-0.309	0.105
Fluoride	7 / 7	0.783	NA	0.143	0.578	0.817	-0.333	-0.333	-0.333	0.978	-0.012	-0.744	-0.427	-0.388	-0.143	0.387
Nitrate	6 / 7	0.000	0.143	NA	0.333	0.817	-0.333	-0.333	-0.333	1.000	0.333	-0.333	-0.333	-0.619	0.143	0.048
Sulfate	7 / 7	0.347	0.578	0.333	NA	0.817	-1.000	-1.000	-1.000	0.521	-0.735	-0.997	-0.950	-0.103	0.048	0.470
Aluminum	7 / 7	0.292	0.733	0.238	0.700	0.817	-1.000	-1.000	-1.000	0.678	-0.587	-0.993	-0.871	-0.476	0.143	0.827
Antimony	4 / 7	-0.444	-0.514	0.206	-0.206	1.000	-0.817	-0.817	-0.817	0.817	0.000	-0.817	-0.817	-0.103	0.103	-0.617
Arsenic	7 / 7	-0.107	-0.402	-0.714	-0.412	-0.817	0.333	0.333	0.333	-0.991	-0.323	0.478	0.100	0.329	0.143	-0.201
Barium	7 / 7	0.134	0.555	0.333	0.454	0.817	-0.333	-0.333	-0.333	0.982	0.378	-0.425	-0.042	-0.419	-0.143	0.938
Beryllium	7 / 7	-0.214	0.280	0.143	0.580	0.000	-0.333	-0.333	-0.333	-0.101	-0.995	-0.745	-0.945	-0.402	0.429	0.854
Boron	6 / 7	0.000	0.143	0.238	0.524	0.000	-0.333	-0.333	-0.333	-0.333	-1.000	-0.333	-0.333	-0.238	0.333	0.619
Cadmium	7 / 7	0.056	0.487	0.143	0.436	0.817	-0.333	-0.333	-0.333	1.000	0.209	-0.578	-0.217	-0.417	0.048	0.944
Calcium	7 / 7	0.575	0.730	0.238	0.544	0.000	0.333	0.333	0.333	0.814	0.730	-0.009	0.379	-0.046	-0.238	0.867
Chromium	7 / 7	-0.207	-0.466	-0.524	0.108	0.000	-0.333	-0.333	-0.333	-0.687	-0.847	-0.183	-0.549	0.606	-0.048	-0.282
Cobalt	7 / 7	0.198	0.617	0.333	0.545	0.817	-0.333	-0.333	-0.333	0.974	-0.031	-0.757	-0.444	-0.395	0.048	0.937
Copper	7 / 7	0.260	0.650	0.619	0.581	0.817	-0.333	-0.333	-0.333	0.996	0.279	-0.518	-0.146	-0.343	-0.238	0.932
Iron	7 / 7	0.733	0.635	0.048	0.501	0.817	-0.333	-0.333	-0.333	0.973	-0.033	-0.758	-0.446	0.167	-0.429	-0.056
Lead	7 / 7	0.695	0.892	0.238	0.656	0.817	-0.333	-0.333	-0.333	0.999	0.244	-0.548	-0.182	-0.165	-0.048	0.722
Lithium	7 / 7	-0.011	0.349	0.195	0.479	0.000	-0.333	-0.333	-0.333	-0.726	-0.817	-0.130	-0.503	-0.345	0.390	0.844
Magnesium	7 / 7	-0.130	0.169	0.238	0.609	0.000	-0.333	-0.333	-0.333	-0.663	-0.864	-0.215	-0.577	-0.007	0.333	0.897
Manganese	7 / 7	0.055	0.489	0.333	0.439	0.817	-0.333	-0.333	-0.333	0.996	0.285	-0.512	-0.140	-0.419	-0.143	0.945
Mercury	7 / 7	0.349	0.411	-0.048	0.139	-0.817	1.000	1.000	1.000	-0.141	0.943	0.883	0.996	0.047	-0.143	0.855
Molybdenum	5 / 7	-0.369	-0.293	0.195	0.000	1.000	-0.817	-0.817	-0.817	0.817	0.000	-0.817	-0.817	-0.098	0.000	0.000
Nickel	7 / 7	0.002	0.441	0.238	0.422	0.817	-0.333	-0.333	-0.333	1.000	0.198	-0.587	-0.228	-0.413	-0.048	0.946
Phosphorous	7 / 7	-0.689	-0.799	-0.333	-0.198	-0.817	0.333	0.333	0.333	-0.978	-0.395	0.409	0.023	0.423	0.333	-0.035
Potassium	7 / 7	0.003	0.525	0.619	0.742	0.817	-1.000	-1.000	-1.000	0.303	-0.875	-0.949	-0.997	-0.676	0.333	0.465
Selenium	7 / 7	0.377	0.612	0.143	0.363	0.000	0.333	0.333	0.333	0.541	0.931	0.362	0.695	-0.170	-0.143	0.923
Silicon	7 / 7	0.137	0.484	0.524	0.870	0.817	-1.000	-1.000	-1.000	0.696	-0.567	-0.990	-0.858	-0.275	-0.143	0.282
Silver	4 / 7	0.252	0.467	0.817	0.467	1.000	-0.817	-0.817	-0.817	0.817	0.000	-0.817	-0.817	-0.700	0.233	0.117
Sodium	5 / 7	0.580	0.781	0.098	0.586	0.817	-1.000	-1.000	-1.000	0.333	-0.333	-1.000	-1.000	-0.390	0.098	0.390
Strontium	7/7	0.519	0.641	0.333	0.233	0.000	0.333	0.333	0.333	0.466	0.959	0.442	0.755	-0.114	-0.333	0.785
Uranium	7 / 7	0.642	0.592	-0.238	0.525	0.000	-0.333	-0.333	-0.333	0.034	-0.973	-0.828	-0.981	-0.003	0.238	0.102
Vanadium	7 / 7	-0.194	-0.530	-0.429	-0.321	-0.817	0.333	0.333	0.333	-0.897	-0.610	0.170	-0.224	0.521	-0.143	-0.562
Zinc	7 / 7	0.432	0.379	-0.048	0.070	0.000	0.333	0.333	0.333	0.004	0.981	0.806	0.972	0.215	-0.333	0.767
Mercury (F0)	2/3	0.522	0.817	0.817	0.817	NA	-0.817	-0.817	-0.817	0.817	0.000	-0.817	-0.817	-0.817	0.000	-0.817
Mercury (F1)	2/3	-0.333	-0.333	-0.333	-1.000	-0.817	NA	1.000	1.000	-0.333	0.333	1.000	1.000	1.000	-0.333	1.000
Mercury (F2)	2/3	-0.333	-0.333	-0.333	-1.000	-0.817	1.000	NA	1.000	-0.333	0.333	1.000	1.000	1.000	-0.333	1.000
Mercury (F3)	2/3	-0.333	-0.333	-0.333	-1.000	-0.817	1.000	1.000	NA	-0.333	0.333	1.000	1.000	1.000	-0.333	1.000
Mercury (F4)	3 / 3	0.959	0.978	1.000	0.521	0.817	-0.333	-0.333	-0.333	NA	0.197	-0.588	-0.229	-0.697	-0.333	-0.271
Mercury (F5)	3 / 3	0.467	-0.012	0.333	-0.735	0.000	0.333	0.333	0.333	0.197	NA	0.677	0.909	0.567	-1.000	0.891
Mercury (F6)	3 / 3	-0.334	-0.744	-0.333	-0.997	-0.817	1.000	1.000	1.000	-0.588	0.677	NA	0.922	0.990	-0.333	0.938
Mercury (FS)	3 / 3	0.057	-0.427	-0.333	-0.950	-0.817	1.000	1.000	1.000	-0.229	0.909	0.922	NA	0.858	-0.333	0.999
Methylmercury	7 / 7	0.190	-0.388	-0.619	-0.103	-0.817	1.000	1.000	1.000	-0.697	0.567	0.990	0.858	NA	-0.333	-0.103
Sulfide	6/7	-0.309	-0.143	0.143	0.048	0.000	-0.333	-0.333	-0.333	-0.333	-1.000	-0.333	-0.333	-0.333	NA	-0.048
TOC	7 / 7	0.105	0.387	0.048	0.470	-0.817	1.000	1.000	1.000	-0.271	0.891	0.938	0.999	-0.103	-0.048	NA

Table E.4. Correlation matrix for sediment analytes for BCV RSE for locations with particle size #10 (cont.)

	Freq. of								Metals							
Chemical	detection	Aluminum	Antimony	Arsenic	Barium	Beryllium	Boron	Cadmium	Calcium	Chromium	Cobalt	Copper	Iron	Lead	Lithium	Magnesium
Chloride	7 / 7	0.292	-0.444	-0.107	0.134	-0.214	0.000	0.056	0.575	-0.207	0.198	0.260	0.733	0.695	-0.011	-0.130
Fluoride	7 / 7	0.733	-0.514	-0.402	0.555	0.280	0.143	0.487	0.730	-0.466	0.617	0.650	0.635	0.892	0.349	0.169
Nitrate	6 / 7	0.238	0.206	-0.714	0.333	0.143	0.238	0.143	0.238	-0.524	0.333	0.619	0.048	0.238	0.195	0.238
Sulfate	7 / 7	0.700	-0.206	-0.412	0.454	0.580	0.524	0.436	0.544	0.108	0.545	0.581	0.501	0.656	0.479	0.609
Aluminum	7 / 7	NA	-0.514	-0.297	0.920	0.847	0.429	0.907	0.827	-0.398	0.955	0.935	0.191	0.847	0.847	0.706
Antimony	4 / 7	-0.514	NA	-0.309	-0.514	-0.206	-0.309	-0.514	-0.617	0.103	-0.309	-0.206	0.103	-0.412	-0.474	-0.309
Arsenic	7 / 7	-0.297	-0.309	NA	-0.299	-0.229	-0.143	-0.252	-0.302	0.276	-0.316	-0.431	-0.345	-0.409	0.061	-0.304
Barium	7 / 7	0.920	-0.514	-0.299	NA	0.859	0.333	0.996	0.852	-0.524	0.990	0.974	-0.062	0.767	0.871	0.771
Beryllium	7 / 7	0.847	-0.206	-0.229	0.859	NA	0.905	0.890	0.591	-0.139	0.874	0.831	-0.128	0.522	0.856	0.895
Boron	6 / 7	0.429	-0.309	-0.143	0.333	0.905	NA	0.333	0.238	0.238	0.524	0.238	0.048	0.429	0.390	0.810
Cadmium	7 / 7	0.907	-0.514	-0.252	0.996	0.890	0.333	NA	0.814	-0.491	0.982	0.957	-0.127	0.714	0.896	0.794
Calcium	7 / 7	0.827	-0.617	-0.302	0.852	0.591	0.238	0.814	NA	-0.391	0.878	0.913	0.321	0.949	0.663	0.662
Chromium	7 / 7	-0.398	0.103	0.276	-0.524	-0.139	0.238	-0.491	-0.391	NA	-0.439	-0.445	0.291	-0.358	-0.394	0.052
Cobalt	7 / 7	0.955	-0.309	-0.316	0.990	0.874	0.524	0.982	0.878	-0.439	NA	0.988	0.063	0.824	0.853	0.792
Copper	7 / 7	0.935	-0.206	-0.431	0.974	0.831	0.238	0.957	0.913	-0.445	0.988	NA	0.135	0.865	0.791	0.800
Iron	7 / 7	0.191	0.103	-0.345	-0.062	-0.128	0.048	-0.127	0.321	0.291	0.063	0.135	NA	0.547	-0.290	-0.024
Lead	7 / 7	0.847	-0.412	-0.409	0.767	0.522	0.429	0.714	0.949	-0.358	0.824	0.865	0.547	NA	0.544	0.541
Lithium	7 / 7	0.847	-0.474	0.061	0.871	0.856	0.390	0.896	0.663	-0.394	0.853	0.791	-0.290	0.544	NA	0.718
Magnesium	7 / 7	0.706	-0.309	-0.304	0.771	0.895	0.810	0.794	0.662	0.052	0.792	0.800	-0.024	0.541	0.718	NA
Manganese	7 / 7	0.906	-0.514	-0.273	0.996	0.890	0.333	1.000	0.816	-0.491	0.982	0.960	-0.121	0.717	0.888	0.798
Mercury	7 / 7	0.620	-0.926	-0.030	0.807	0.491	0.333	0.792	0.871	-0.505	0.772	0.778	-0.097	0.689	0.682	0.569
Molybdenum	5 / 7	-0.098	0.422	-0.098	-0.098	0.390	0.293	-0.098	-0.195	0.098	0.098	0.195	0.195	0.000	-0.250	0.293
Nickel	7 / 7	0.891	-0.412	-0.252	0.990	0.905	0.429	0.998	0.789	-0.467	0.973	0.947	-0.160	0.680	0.895	0.813
Phosphorous	7 / 7	-0.345	0.000	0.502	-0.269	0.133	0.429	-0.191	-0.429	0.739	-0.268	-0.335	-0.385	-0.573	-0.032	0.231
Potassium	7 / 7	0.810	-0.103	-0.302	0.626	0.743	0.429	0.633	0.389	-0.315	0.656	0.619	0.026	0.481	0.721	0.493
Selenium	7 / 7	0.821	-0.514	-0.259	0.933	0.666	0.333	0.909	0.956	-0.507	0.927	0.939	0.090	0.854	0.735	0.685
Silicon	7 / 7	0.557	0.206	-0.629	0.316	0.529	0.333	0.298	0.317	0.219	0.418	0.462	0.581	0.499	0.203	0.515
Silver	4 / 7	0.583	-0.126	-0.467	0.583	0.117	0.233	0.467	0.467	-0.583	0.467	0.583	0.000	0.467	0.538	0.233
Sodium	5 / 7	0.878	-0.580	0.000	0.781	0.293	0.390	0.878	0.683	-0.195	0.683	0.488	0.195	0.781	0.650	0.293
Strontium	7 / 7	0.684	-0.514	-0.267	0.818	0.429	0.143	0.777	0.922	-0.651	0.794	0.828	0.092	0.822	0.598	0.474
Uranium	7 / 7	0.439	-0.514	0.365	0.136	0.104	0.333	0.115	0.340	0.073	0.222	0.174	0.487	0.471	0.329	-0.020
Vanadium	7 / 7	-0.645	0.206	0.478	-0.727	-0.455	-0.048	-0.701	-0.606	0.868	-0.668	-0.700	0.211	-0.556	-0.631	-0.356
Zinc	7 / 7	0.476	-0.617	-0.082	0.682	0.328	0.333	0.655	0.845	-0.420	0.653	0.687	0.020	0.668	0.490	0.494
Mercury (F0)	2/3	0.817	1.000	-0.817	0.817	0.000	0.000	0.817	0.000	0.000	0.817	0.817	0.817	0.817	0.000	0.000
Mercury (F1)	2/3	-1.000	-0.817	0.333	-0.333	-0.333	-0.333	-0.333	0.333	-0.333	-0.333	-0.333	-0.333	-0.333	-0.333	-0.333
Mercury (F2)	2/3	-1.000	-0.817	0.333	-0.333	-0.333	-0.333	-0.333	0.333	-0.333	-0.333	-0.333	-0.333	-0.333	-0.333	-0.333
Mercury (F3)	2/3	-1.000	-0.817	0.333	-0.333	-0.333	-0.333	-0.333	0.333	-0.333	-0.333	-0.333	-0.333	-0.333	-0.333	-0.333
Mercury (F4)	3 / 3	0.678	0.817	-0.991	0.982	-0.101	-0.333	1.000	0.814	-0.687	0.974	0.996	0.973	0.999	-0.726	-0.663
Mercury (F5)	3 / 3	-0.587	0.000	-0.323	0.378	-0.995	-1.000	0.209	0.730	-0.847	-0.031	0.279	-0.033	0.244	-0.817	-0.864
Mercury (F6)	3 / 3	-0.993	-0.817	0.478	-0.425	-0.745	-0.333	-0.578	-0.009	-0.183	-0.757	-0.518	-0.758	-0.548	-0.130	-0.215
Mercury (FS)	3/3	-0.871	-0.817	0.100	-0.042	-0.945	-0.333	-0.217	0.379	-0.549	-0.444	-0.146	-0.446	-0.182	-0.503	-0.577
Methylmercury	7 / 7	-0.476	-0.103	0.329	-0.419	-0.402	-0.238	-0.417	-0.046	0.606	-0.395	-0.343	0.167	-0.165	-0.345	-0.007
Sulfide	6 / 7 7 / 7	0.143 0.827	0.103	0.143	-0.143	0.429	0.333	0.048	-0.238	-0.048	0.048	-0.238	-0.429	-0.048	0.390	0.333 0.897
TOC	7/7	0.827	-0.617	-0.201	0.938	0.854	0.619	0.944	0.867	-0.282	0.937	0.932	-0.056	0.722	0.844	0.897

Table E.4. Correlation matrix for sediment analytes for BCV RSE for locations with particle size #10 (cont.)

	Freq. of							Me	tals						
Chemical	detection	Manganese	Mercury	Molybdenum	Nickel	Phosphorous	Potassium	Selenium	Silicon	Silver	Sodium	Strontium	Uranium	Vanadium	Zinc
Chloride	7 / 7	0.055	0.349	-0.369	0.002	-0.689	0.003	0.377	0.137	0.252	0.580	0.519	0.642	-0.194	0.432
Fluoride	7 / 7	0.489	0.411	-0.293	0.441	-0.799	0.525	0.612	0.484	0.467	0.781	0.641	0.592	-0.530	0.379
Nitrate	6 / 7	0.333	-0.048	0.195	0.238	-0.333	0.619	0.143	0.524	0.817	0.098	0.333	-0.238	-0.429	-0.048
Sulfate	7/7	0.439	0.139	0.000	0.422	-0.198	0.742	0.363	0.870	0.467	0.586	0.233	0.525	-0.321	0.070
Aluminum	7 / 7	0.906	0.620	-0.098	0.891	-0.345	0.810	0.821	0.557	0.583	0.878	0.684	0.439	-0.645	0.476
Antimony	4 / 7	-0.514	-0.926	0.422	-0.412	0.000	-0.103	-0.514	0.206	-0.126	-0.580	-0.514	-0.514	0.206	-0.617
Arsenic	7 / 7	-0.273	-0.030	-0.098	-0.252	0.502	-0.302	-0.259	-0.629	-0.467	0.000	-0.267	0.365	0.478	-0.082
Barium	7 / 7	0.996	0.807	-0.098	0.990	-0.269	0.626	0.933	0.316	0.583	0.781	0.818	0.136	-0.727	0.682
Beryllium	7 / 7	0.890	0.491	0.390	0.905	0.133	0.743	0.666	0.529	0.117	0.293	0.429	0.104	-0.455	0.328
Boron	6/7	0.333	0.333	0.293	0.429	0.429	0.429	0.333	0.333	0.233	0.390	0.143	0.333	-0.048	0.333
Cadmium	7 / 7	1.000	0.792	-0.098	0.998	-0.191	0.633	0.909	0.298	0.467	0.878	0.777	0.115	-0.701	0.655
Calcium	7/7	0.816	0.871	-0.195	0.789	-0.429	0.389	0.956	0.317	0.467	0.683	0.922	0.340	-0.606	0.845
Chromium	7/7	-0.491	-0.505	0.098	-0.467	0.739	-0.315	-0.507	0.219	-0.583	-0.195	-0.651	0.073	0.868	-0.420
Cobalt	7/7	0.982	0.772	0.098	0.973	-0.268	0.656	0.927	0.418	0.467	0.683	0.794	0.222	-0.668	0.653
Copper	7/7	0.960	0.778	0.195	0.947	-0.335	0.619	0.939	0.462	0.583	0.488	0.828	0.174	-0.700	0.687
Iron	7/7	-0.121	-0.097	0.195	-0.160	-0.385	0.026	0.090	0.581	0.000	0.195	0.092	0.487	0.211	0.020
Lead	7 / 7	0.717	0.689	0.000	0.680	-0.573	0.481	0.854	0.499	0.467	0.781	0.822	0.471	-0.556	0.668
Lithium	7 / 7	0.888	0.682	-0.250	0.895	-0.032	0.721	0.735	0.203	0.538	0.650	0.598	0.329	-0.631	0.490
Magnesium	7 / 7	0.798	0.569	0.293	0.813	0.231	0.493	0.685	0.515	0.233	0.293	0.474	-0.020	-0.356	0.494
Manganese	7 / 7	NA	0.789	-0.098	0.998	-0.196	0.631	0.910	0.308	0.583	0.781	0.779	0.101	-0.704	0.655
Mercury	7 / 7	0.789	NA	-0.293	0.777	-0.252	0.133	0.936	-0.143	0.233	0.586	0.943	0.103	-0.597	0.969
Molybdenum	5 / 7	-0.098	-0.293	NA	0.000	0.488	0.098	-0.098	0.390	-0.120	-0.150	-0.098	-0.390	0.000	-0.098
Nickel	7 / 7	0.998	0.777	0.000	NA	-0.145	0.626	0.893	0.297	0.467	0.781	0.751	0.075	-0.684	0.639
Phosphorous	7 / 7	-0.196	-0.252	0.488	-0.145	NA	-0.262	-0.356	-0.143	-0.467	-0.195	-0.542	-0.235	0.651	-0.266
Potassium	7 / 7	0.631	0.133	0.098	0.626	-0.262	NA	0.370	0.665	0.700	0.488	0.226	0.407	-0.585	-0.066
Selenium	7 / 7	0.910	0.936	-0.098	0.893	-0.356	0.370	NA	0.174	0.350	0.683	0.951	0.166	-0.661	0.883
Silicon	7 / 7	0.308	-0.143	0.390	0.297	-0.143	0.665	0.174	NA	0.350	0.195	0.005	0.256	-0.141	-0.181
Silver	4 / 7	0.583	0.233	-0.120	0.467	-0.467	0.700	0.350	0.350	NA	0.538	0.467	0.117	-0.583	0.117
Sodium	5 / 7	0.781	0.586	-0.150	0.781	-0.195	0.488	0.683	0.195	0.538	NA	0.586	0.683	-0.488	0.488
Strontium	7 / 7	0.779	0.943	-0.098	0.751	-0.542	0.226	0.951	0.005	0.467	0.586	NA	0.140	-0.727	0.926
Uranium	7 / 7	0.101	0.103	-0.390	0.075	-0.235	0.407	0.166	0.256	0.117	0.683	0.140	NA	0.016	0.029
Vanadium	7 / 7	-0.704	-0.597	0.000	-0.684	0.651	-0.585	-0.661	-0.141	-0.583	-0.488	-0.727	0.016	NA	-0.490
Zinc	7 / 7	0.655	0.969	-0.098	0.639	-0.266	-0.066	0.883	-0.181	0.117	0.488	0.926	0.029	-0.490	NA
Mercury (F0)	2/3	0.817	-0.817	1.000	0.817	-0.817	0.817	0.000	0.817	1.000	0.817	0.000	0.000	-0.817	0.000
Mercury (F1)	2/3	-0.333	1.000	-0.817	-0.333	0.333	-1.000	0.333	-1.000	-0.817	-1.000	0.333	-0.333	0.333	0.333
Mercury (F2)	2/3	-0.333	1.000	-0.817	-0.333	0.333	-1.000	0.333	-1.000	-0.817	-1.000	0.333	-0.333	0.333	0.333
Mercury (F3)	2/3	-0.333	1.000	-0.817	-0.333	0.333	-1.000	0.333	-1.000	-0.817	-1.000	0.333	-0.333	0.333	0.333
Mercury (F4)	3 / 3	0.996	-0.141	0.817	1.000	-0.978	0.303	0.541	0.696	0.817	0.333	0.466	0.034	-0.897	0.004
Mercury (F5)	3 / 3	0.285	0.943	0.000	0.198	-0.395	-0.875	0.931	-0.567	0.000	-0.333	0.959	-0.973	-0.610	0.981
Mercury (F6)	3 / 3	-0.512	0.883	-0.817	-0.587	0.409	-0.949	0.362	-0.990	-0.817	-1.000	0.442	-0.828	0.170	0.806
Mercury (FS)	3 / 3	-0.140	0.996	-0.817	-0.228	0.023	-0.997	0.695	-0.858	-0.817	-1.000	0.755	-0.981	-0.224	0.972
Methylmercury	7 / 7	-0.419	0.047	-0.098	-0.413	0.423	-0.676	-0.170	-0.275	-0.700	-0.390	-0.114	-0.003	0.521	0.215
Sulfide	6 / 7	-0.143	-0.143	0.000	-0.048	0.333	0.333	-0.143	-0.143	0.233	0.098	-0.333	0.238	-0.143	-0.333
TOC	7 / 7	0.945	0.855	0.000	0.946	-0.035	0.465	0.923	0.282	0.117	0.390	0.785	0.102	-0.562	0.767

Table E.4. Correlation matrix for sediment analytes for BCV RSE for locations with particle size #10 (cont.)

# Correlation is significant at the 0.05 significance level where $0.01 \le \text{two-sided } p\text{-value} \le 0.05$ .

Correlation is significant at the 0.01 significance level where  $0.001 \le \text{two-sided } p\text{-value} \le 0.01$ .

Correlation is significant at the 0.001 significance level where  $0.0001 \le \text{two-sided } p\text{-value} \le 0.001$ .

Pearson correlations are calculated when both variables have all detections.

Kendall tau correlations are calculated when there is at least one non-detection in one or both variables, but at least half detections in each variable. BCV = Bear Creek Valley

Freq. = frequency
NA = not applicable
RSE = remedial site evaluation
TOC = Total organic carbon average

The pairwise correlation tables were used to identify analytes that are significantly correlated with mercury in soil and sediment. For example, Table E.1 for all soil combined shows the variables most highly correlated with mercury and having all detected concentrations were fluoride, cadmium, copper, lead, lithium, zinc, aluminum, arsenic, calcium, nickel, phosphorous, selenium, silicon, strontium, uranium, vanadium, and total organic carbon (TOC) (exclusive of the sequential extraction results that are expected to be correlated with mercury and have only n = 18 samples). The correlation analysis was used as an exploratory tool to identify analytes that would likely be good predictors for mercury. Because there were so many analytes and the dataset has a relatively small number of samples, the number of probable predictors needed to be subset before using them in the multivariate models.

Four multivariate model selection methods using information criteria were used to evaluate all possible  $2^k - 1$  subset models simultaneously for k independent variables in the model. These four criteria are described in more detail in Section E.2. SAS® version 9.4 was used to evaluate all possible subset models using the four criteria (Beal 2007). For example, if k = 10 variables were selected as the most correlated (and detected) with mercury from Tables E.1 and E.3, and then all possible  $2^{10} - 1 = 1023$  subset models were evaluated simultaneously. Four information criteria were calculated for each subset model and were sorted in ascending order from smallest to largest for each criterion. The four criteria may not agree which model is best. In that case, professional judgment is used to select the overall best model using the guiding statistical principle of parsimony, which of the four criteria agree, and historical experience in Bear Creek Valley soil and sediment. From a statistical perspective, a more parsimonious (fewer independent parameters) is preferred over models with more independent parameters. Multicollinearity was considered and examined for each of the final models and found not to be a problem because the purpose of the modeling is exploratory and not predictive given the relatively small number of samples.

# **E.2. INFORMATION CRITERIA METHODS**

The following sections describe various criteria methods.

#### E.2.1 AKAIKE'S INFORMATION CRITERION

Akaike (1973) introduced the concept of information criteria as a tool for optimal model selection. Other authors who use the Akaike's Information Criterion (AIC) for model selection include Akaike (1987) and Bozdogan (1987, 2000). AIC is a function of the number of observations n, the sum of squared errors (SSE), and the number of independent variables k, as shown in Equation 1.

$$AIC = n \cdot \ln\left(\frac{SSE}{n}\right) + 2k$$
 (Equation 1)

The first term in Equation 1 is a measure of the model lack of fit, while the second term—2k—is a penalty term for additional parameters in the model. Therefore, as the number of independent variables k included in the model increases, the lack of fit term decreases while the penalty term increases. Conversely, as variables are dropped from the model, the lack of fit term increases while the penalty term decreases. The model with the smallest AIC is deemed the "best" model because it minimizes the difference from the given model to the "true" model.

#### E.2.2 AKAIKE'S INFORMATION CRITERION CORRECTED

When the sample size n is small, AIC tends to select models that have too many parameters (i.e., AIC tends to overfit). To address such potential overfitting, AIC corrected (AICc) was developed. AICc is AIC with a correction for small sample sizes, as shown in Equation 2.

$$AICc = AIC + \frac{2k(k+1)}{n-k-1}$$
 (Equation 2)

The second term in Equation 2 is an additional penalty term that discourages AIC from overfitting the model. Thus, AICc is essentially AIC with an extra penalty term for the number of parameters. Note that, as  $n \to \infty$ , the extra penalty term converges to 0, and thus AICc converges to AIC. The formula for AIC includes k but not  $k^2$ . In other words, AIC is a first-order estimate of the information loss; whereas, AICc is a second-order estimate. The model with the smallest AICc is deemed the "best" model because it minimizes the difference from the given model to the "true" model.

#### E.2.3 BAYESIAN INFORMATION CRITERION

Sawa (1978) developed a model selection criterion that was derived from a Bayesian modification of the AIC criterion. Bayesian Information Criterion (BIC) is a function of the number of observations (n), the SSE, the pure error variance fitting the full model  $(\sigma^2)$ , and the number of independent variables (k), as shown in Equation 3.

$$BIC = n \cdot \ln\left(\frac{SSE}{n}\right) + \frac{2(k+2)n\sigma^2}{SSE} - \frac{2n^2\sigma^4}{SSE^2}$$
 (Equation 3)

The penalty term for BIC is more complex than the AIC penalty term and is a function of n, the SSE, and  $\sigma^2$ , in addition to k. The model with the smallest BIC is deemed the "best" model because it minimizes the difference from the given model to the "true" model.

## E.2.4 SCHWARZ BAYESIAN CRITERION

Schwarz (1978) developed a model selection criterion that was derived from a Bayesian modification of the AIC criterion. Schwarz Bayesian Criterion (SBC) is a function of the number of observations (n), the SSE, and the number of independent variables (k), as shown in Equation 4.

$$SBC = n \cdot \ln \left( \frac{SSE}{n} \right) + k \ln n$$
 (Equation 4)

The penalty term for SBC is similar to AIC in Equation 4, but it uses a multiplier of  $\ln n$  for k instead of a constant 2, by incorporating the sample size n. The model with the smallest SBC is deemed the "best" model because it minimizes the difference from the given model to the "true" model.

# **E.3. SOIL RESULTS**

The modeling began as a descriptive process because the dataset is too small to establish definitive predictive models. For exploratory purposes, four different selection criteria were used for model evaluation to see which models were the best for each of the four information criteria using mercury as the dependent variable. The scores from these four criteria cannot be compared with each other because they score the models differently.

#### E.3.1 FLOODPLAIN AND CREEK BANK SOIL COMBINED

From Table E.1, the following analytes were selected that were significantly correlated with mercury and had all detected concentrations: fluoride, cadmium, copper, lead, lithium, zinc, aluminum, arsenic, calcium, nickel, phosphorous, selenium, silicon, strontium, uranium, vanadium, and TOC average. The sequential extraction analytes were excluded from consideration because there were only 18 samples, compared to 48 samples for the other analytes. Including even 1 sequential extraction analyte would reduce the dataset to 18 records, which would further limit the number of regressors that could be considered in the model because the number of regressors could exceed the number of observations. The best model with the lowest AIC score using all floodplain and creek bank soil data has an AIC = -71.4113, an adjusted  $R^2 = 0.8864$ , a root mean square error (RMSE) = 0.4274, an F = 34.35, an F *p*-value = < 0.0001, and *k* = 11 variables (not counting the intercept).

Table E.5 shows the parameter estimates, standard errors, student's t-statistics, and two-sided *p*-values for the best AIC model.

Table E.5. Model statistics for all soil for best AIC model

Independent variable	Parameter estimate (mg/kg)	Standard error (mg/kg)	Student's t-statistic	Two-sided <i>p</i> -value
Intercept	-2.31956	0.43593	-5.32	< 0.0001
Fluoride	-0.09391	0.02596	-3.62	0.0009
Cadmium	0.1079	0.06458	1.67	0.1034
Copper	0.19591	0.04088	4.79	< 0.0001
Lead	0.09284	0.03024	3.07	0.0041
Lithium	0.12433	0.03412	3.64	0.0008
Zinc	-0.03543	0.01313	-2.70	0.0105
Calcium	5.68E-05	2.09E-05	2.72	0.0101
Nickel	-0.02166	0.00983	-2.20	0.0341
Silicon	1.27E-04	8.67E-05	1.46	0.1524
Uranium	-0.03002	0.00564	-5.32	< 0.0001
Vanadium	-0.04043	0.0154	-2.63	0.0127

AIC = Akaike Information Criterion

Table E.5 shows cadmium and silicon do not significantly contribute to the model with *p*-values of 0.1034 and 0.1524, respectively. Figure E.1 is a plot that shows the multivariate-predicted concentrations and 95% prediction limits compared with the actual mercury concentrations. Different symbols for the bank soil, bank soil lower, bank soil upper, and floodplain soil are shown to see if there are any patterns by soil type. Figure E.1 shows the soil types are scattered fairly uniformly throughout the distribution. All concentrations, except BCT13-BSU, are within the 95% prediction limits.





Figure E.1. Best AIC model for all soil.

Figure E.2 shows a normal probability plot of the best AIC model residuals. Model residuals were tested for normality using the Shapiro-Wilk (SW) test, which showed the residuals are approximately normally distributed at the 0.05 significance level, with an SW p-value = 0.0572. All residuals, except BCT13-BSU, plot approximately linearly along the regression line.

The best model with the lowest AICc, BIC, and SBC scores using all floodplain and creek bank soil data has an AICc = -65.9029, a BIC = -63.6272, an SBC = -52.7544, an adjusted  $R^2 = 0.8766$ , an RMSE = 0.44547, an F = 42.74, an F p-value = < 0.0001, and k = 8 variables (not counting the intercept). Both the BIC and SBC scores are more than 1 lower than the BIC and SBC scores of the model with the next lowest scores. This indicates the model is considerably better than the model with the next lowest BIC and SBC scores.

Table E.6 shows the parameter estimates, standard errors, student's t-statistics, and two-sided *p*-values for the best AICc, BIC, and SBC model.



Deviates from the Standard Normal Cumulative Distribution Function

AIC = Akaike Information Criterion

BSU = upper creek bank soil

Figure E.2. AIC model residuals for all soil.

Table E.6. Model statistics for all soil for best AICc, BIC, and SBC model

Independent variable	Parameter estimate (mg/kg)	Standard error (mg/kg)	Student's t-statistic	Two-sided <i>p</i> -value
Intercept	-2.5047	0.3638	-6.88	< 0.0001
Fluoride	-0.0841	0.02264	-3.71	0.0006
Copper	0.22741	0.03789	6.00	< 0.0001
Lead	0.08844	0.03037	2.91	0.0059
Lithium	0.14225	0.03163	4.50	< 0.0001
Zinc	-0.05004	0.00974	-5.14	< 0.0001
Calcium	6.53E-05	2.14E-05	3.05	0.0041
Uranium	-0.02853	0.00568	-5.02	< 0.0001
Vanadium	-0.03417	0.01455	-2.35	0.024

AICc = Akaike's Information Criterion

BIC = Bayesian Information Criterion

SBC = Schwarz Bayesian Criterion

Table E.6 shows all variables significantly contribute to the model at the 0.05 significance level. Figure E.3 is a plot that shows the multivariate-predicted concentrations and 95% prediction limits compared with the actual mercury concentrations. Different symbols for the bank soil, bank soil lower, bank soil upper, and floodplain soil are shown to see if there are any patterns by soil type. Figure E.3 shows the soil types are scattered fairly uniformly throughout the distribution. All concentrations, except BCT13-BSU, are within the 95% prediction limits.




Figure E.3. Best AICc, BIC, and SBC model for all soil.

Figure E.4 shows a normal probability plot of the best AICc, BIC, and SBC model residuals. Model residuals were tested for normality using the SW test, which showed the residuals are approximately normally distributed at the 0.05 significance level, with an SW p-value = 0.0618. All residuals, except BCT13-BSU, plot approximately linearly along the regression line.




Figure E.4. AICc, BIC, and SBC model residuals for all soil.

A comparison of the best AIC model (Table E.5) with the best model determined by AICc, BIC, and SBC (Table E.6) shows the preferred model to be the AICc, BIC, and SBC model because it is more parsimonious (k = 8 variables instead of k = 11 for AIC). AIC tends to overfit the data by selecting more variables compared to other information criteria. The AIC model also selected two variables that did not significantly contribute to the model. Because AICc, BIC, and SBC agree on the same model and it is more parsimonious than the AIC model, the model shown in Table E.6 was selected as the best overall model for predicting mercury for all soil data combined.

#### E.3.2 FLOODPLAIN SOIL

There are n = 17 samples from the floodplain soil. Pearson correlations were calculated between all 29 analytes detected in all 17 samples. The following analytes were significantly correlated with mercury at the 0.05 significance level and had all detected concentrations: fluoride, aluminum, cadmium, copper, lead, lithium, phosphorous, selenium, silicon, uranium, zinc, methylmercury, and TOC. The sequential extraction analytes were excluded from consideration because there were only 9 samples, compared to 17 samples for the other analytes. Including even one sequential extraction analyte would reduce the dataset to nine records, which would further limit the number of regressors that could be considered in the model because the number of regressors could exceed the number of observations.

The best model with the lowest AIC score using all floodplain soil data has an AIC = -44.8449, an adjusted  $R^2 = 0.9627$ , an RMSE = 0.2296, an F = 60.03, an F p-value = < 0.0001, and k = 7 variables (not counting the intercept). Table E.7 shows the parameter estimates, standard errors, student's t-statistics, and two-sided p-values for the best AIC model for floodplain soil.

Table E.7. Model statistics for floodplain soil for AIC model

Independent variable	Parameter estimate (mg/kg)	Standard error (mg/kg)	Student's t-statistic	Two-sided <i>p</i> -value
Intercept	-0.33472	0.44576	-0.75	0.4719
Fluoride	0.2668	0.03355	7.95	< 0.0001
Copper	-0.05141	0.03447	-1.49	0.17
Lead	0.08922	0.02337	3.82	0.0041
Phosphorous	-0.00428	0.00368	-1.16	0.2749
Selenium	-0.4963	0.20929	-2.37	0.0418
Uranium	0.01332	0.00516	2.58	0.0297
TOC	2.89E-05	1.69E-05	1.71	0.1218

AIC = Akaike's Information Criterion

TOC = total organic carbon

Table E.7 shows the intercept, copper, phosphorous, and TOC do not significantly contribute to the model at the 0.05 significance level. Figure E.5 is a plot that shows the multivariate-predicted concentrations and 95% prediction limits compared with the actual mercury concentrations. Figure E.5 shows all concentrations are within the 95% prediction limits.





Figure E.5. Best AIC model for floodplain soil.

Figure E.6 shows a normal probability plot of the best AIC model residuals. Model residuals were tested for normality using the SW test, which showed the residuals are approximately normally distributed at the 0.05 significance level, with an SW p-value = 0.8977. All residuals plot approximately linearly along the regression line.



Deviates from the Standard Normal Cumulative Distribution Function

AIC = Alaike Information Criterion

Figure E.6. AIC model residuals for floodplain soil.

The best model with the lowest AICc, BIC, and SBC scores using all floodplain soil data has an AICc = -40.4186, a BIC = -29.6802, an SBC = -39.5859, an adjusted R<sup>2</sup> = 0.9576, an RMSE = 0.2449, an F = 91.27, an F p-value = <0.0001, and k = 4 variables (not counting the intercept). Table E.8 shows the parameter estimates, standard errors, student's t-statistics, and two-sided p-values for the best AICc, BIC, and SBC model for floodplain soil.

Table E.8. Model statistics for floodplain soil for AICc, BIC, and SBC

Independent variable	Parameter estimate (mg/kg)	Standard error (mg/kg)	Student's t-statistic	Two-sided <i>p</i> -value
Intercept	-0.76039	0.2266	-3.36	0.0057
Fluoride	0.23247	0.02512	9.25	< 0.0001
Lead	0.06446	0.01847	3.49	0.0045
Selenium	-0.46983	0.20547	-2.29	0.0412
Uranium	0.0115	0.00482	2.39	0.0344

AICc = Akaike's Information Criterion corrected

BIC = Bayesian Information Criterion

SBC = Schwarz Bayesian Criterion

Table E.8 shows all variables significantly contribute to the model at the 0.05 significance level. Copper, phosphorous, and TOC have been removed from the AICc, BIC, and SBC model compared to the AIC model in Table E.7. Figure E.7 is a plot that shows the multivariate-predicted concentrations and 95% prediction limits compared with the actual mercury concentrations. Figure E.7 shows all concentrations are within the 95% prediction limits.




Figure E.7. Best AICc, BIC, and SBC model for floodplain soil.

Figure E.8 shows a normal probability plot of the best AICc, BIC, and SBC model residuals. Model residuals were tested for normality using the SW test, which showed the residuals are approximately normally distributed at the 0.05 significance level, with an SW p-value = 0.298. All residuals, except the smallest residual of -0.5485 from BCT9-FP (observation 10 in Figure E.7.), plot approximately linearly along the regression line.




Figure E.8. AICc, BIC, and SBC model residuals for floodplain soil.

A comparison of the best AIC model (Table E.7) with the best model determined by AICc, BIC, and SBC (Table E.8) shows the preferred model to be the AICc, BIC, and SBC model because it is more parsimonious (k = 4 variables instead of k = 7 for AIC). AIC tends to overfit the data by selecting more variables compared to other information criteria. The AIC model also selected copper, phosphorous, and TOC, which did not significantly contribute to the model. Because AICc, BIC, and SBC agree on the same model and it is more parsimonious than the AIC model, the model shown in Table E.8 was selected as the best overall model for predictING MERCURY FOR FLOODPLAIN SOIL DATA.

### E.3.3 CREEK BANK SOIL

There are n = 31 samples from creek bank soil. Pearson correlations were calculated between all 27 analytes detected in all 31 samples. The following analytes were significantly correlated with mercury at the 0.05 significance level and had all detected concentrations: fluoride, arsenic, cadmium, calcium, copper, lead, lithium, phosphorous, selenium, strontium, zinc, and TOC. The sequential extraction analytes were excluded from consideration because there were only 9 samples, compared to 31 samples for the other analytes. Including even one sequential extraction analyte would reduce the dataset to nine records, which would further limit the number of regressors that could be considered in the model because the number of regressors could exceed the number of observations.


The best model with the lowest AIC score using all creek bank soil data has an AIC = -31.6709, an adjusted  $R^2 = 0.8337$ , an RMSE = 0.5381, an F = 22.49, an F p-value = < 0.0001, and k = 7 variables (not counting the intercept). Table E.9 shows the parameter estimates, standard errors, student's t-statistics, and two-sided p-values for the best AIC model for creek bank soil.

Table E.9. Model statistics for creek bank soil for AIC model

Independent variable	Parameter estimate (mg/kg)	Standard error (mg/kg)	Student's t-statistic	Two-sided <i>p</i> -value
Intercept	-1.7876	0.4639	-3.85	0.0008
Fluoride	-0.10065	0.0426	-2.36	0.027
Cadmium	0.13534	0.06649	2.04	0.0535
Calcium	3.93E-04	1.17E-04	3.36	0.0027
Copper	0.33347	0.05754	5.80	< 0.0001
Lead	0.09004	0.04748	1.90	0.0705
Selenium	-0.13023	0.04575	-2.85	0.0091
Zinc	-0.07549	0.01944	-3.88	0.0008

AIC = Akaike's Information Criterion


Table E.9 shows cadmium and lead do not significantly contribute to the model at the 0.05 significance level. Figure E.9 is a plot that shows the multivariate-predicted concentrations and 95% prediction limits compared with the actual mercury concentrations. Figure E.9 shows all concentrations are within the 95% prediction limits.



AIC = Akaike's Information Criterion BCV = Bear Creek Valley BSU = upper creek bank soil

Figure E.9. Best AIC model for creek bank soil.

Figure E.10 shows a normal probability plot of the best AIC model residuals. Model residuals were tested for normality using the SW test, which showed the residuals are approximately normally distributed at the 0.05 significance level, with an SW p-value = 0.653. All residuals plot approximately linearly along the regression line.



Deviates from the Standard Normal Cumulative Distribution Function

AIC = Alaike Information Criterion

Figure E.10. AIC model residuals for creek bank soil.

The best model with the lowest AICc, BIC, and SBC scores using all creek bank soil data has an AICc = -29.2201, a BIC = -25.808, an SBC = -22.0501, an adjusted  $R^2 = 0.8068$ , an RMSE = 0.58005, an F = 32.33, an F p-value = < 0.0001, and k = 4 variables (not counting the intercept). Table E.10 shows the parameter estimates, standard errors, student's t-statistics, and two-sided p-values for the best AICc, BIC, and SBC model for creek bank soil.

Table E.10. Model statistics for creek bank soil for AICc, BIC, and SBC


Independent variable	Parameter estimate (mg/kg)	Standard error (mg/kg)	Student's t-statistic	Two-sided <i>p</i> -value
Intercept	-1.37462	0.23	-5.98	< 0.0001
Calcium	3.42E-04	1.17E-04	2.93	0.007
Copper	0.31536	0.04581	6.88	< 0.0001
Strontium	-0.09753	0.04185	-2.33	0.0278
Zinc	-0.04884	0.0169	-2.89	0.0077

AICc = Akaike's Information Criterion corrected

BIC = Bayesian Information Criterion

SBC = Schwarz Bayesian Criterion

Table E.10 shows all variables significantly contribute to the model at the 0.05 significance level. Cadmium and lead were eliminated from the AIC model in Table E.9 because they did not significantly contribute to the model at the 0.05 significance level. In addition, compared to the AIC model, fluoride and selenium were eliminated and replaced with strontium. Figure E.11 is a plot that shows the multivariate-predicted concentrations and 95% prediction limits compared with the actual mercury concentrations. Figure E.11 shows all concentrations, except BCT13-BSU, are within the 95% prediction limits.



AICc = Akaike's Information Criterion corrected

BCV = Bear Creek Valley

BIC = Bayesian Information Criterion

BSU = upper creek bank soil

SBC = Schwarz Bayesian Criterion

Figure E.11. Best AICc, BIC, and SBC model for creek bank soil.

Figure E.12 shows a normal probability plot of the best AICc, BIC, and SBC model residuals. Model residuals were tested for normality using the SW test, which showed the residuals are approximately normally distributed at the 0.04 significance level, with an SW p-value = 0.0457. The two smallest residuals (BCT4-BSL and BCT12B-BSL) and the largest residual (BCT13-BSU) cause the SW p-value to be slightly below 0.05. All other residuals plot approximately linearly along the regression line.

A comparison of the best AIC model (Table E.9) with the best model determined by AICc, BIC, and SBC (Table E.10) shows the preferred model to be the AICc, BIC, and SBC model because it is more parsimonious (k = 4 variables instead of k = 7 for AIC). AIC tends to overfit the data by selecting more variables compared to other information criteria. The AIC model also selected lead and cadmium, which did not significantly contribute to the model. The AICc, BIC, and SBC scores are more than 1 lower than the AICc, BIC, and SBC scores of the model with the next lowest scores. This indicates the model is considerably better than the model with the next lowest AICc, BIC, and SBC scores. Because AICc, BIC, and SBC agree on the same model and it is more parsimonious than the AIC model, the model shown in Table E.10 was selected as the best overall model for predicting mercury for creek bank soil data.




Figure E.12. AICc, BIC, and SBC model residuals for creek bank soil.

A comparison of the overall best models for floodplain soil with creek bank soil shows fluoride, lead, selenium, and uranium best predict mercury concentrations in floodplain soil, while calcium, copper, strontium, and zinc best predict mercury concentrations in creek bank soil. This may indicate differences in mercury characteristics in the floodplain and creek bank soils.

#### E.3.4 FLOODPLAIN AND CREEK BANK SOIL BY PARTICLE SIZE

The particle sizes of the floodplain and creek bank soil data were also a part of the database. Pearson correlations were calculated for all pairs of variables with all detected concentrations for particle size #10 using data from locations BCT14-BSL, BCT14-BSU, BCT14-FP, BCT3-BSL, BCT4-BSU, and BCT4-FP. Kendall's tau correlations were calculated for pairs of variables with at least one non-detect, and at least one-half of the six samples were detected. There were only two sequential extraction samples collected with particle size #10, so these were excluded. Table E.2 summarizes the correlations of soil for particle size #10. Correlations that test the null hypothesis of zero correlation and are significant at the 0.05 significance level are highlighted by significance. Correlations that are not highlighted are not significant at the 0.05 significance level.

For particle size #10 soil, arsenic, copper, lithium, manganese, phosphorous, zinc, and TOC were significantly positively correlated with mercury at the 0.05 significance level. All these analytes, except manganese, are also significantly correlated with mercury in all soil (Table E.1).

## **E.4. SEDIMENT RESULTS**

As with soil, the modeling for sediment began as a descriptive process because the dataset is too small (n = 12) to establish definitive predictive models. For exploratory purposes, four different selection criteria were used for model evaluation to see which models were the best for each of the four information criteria using mercury as the dependent variable. The scores from these four criteria cannot be compared with each other because they score the models differently.

From Table E.3, the following analytes were selected that were significantly correlated with mercury and had all detected concentrations: chloride, calcium, selenium, and TOC. The sequential extraction analytes were excluded from consideration because there were only 6 samples, compared to 12 samples for the other analytes. Including even one sequential extraction analyte would reduce the dataset to six observations, which would further limit the number of regressors that could be considered in the model. The best model with the lowest AIC, AICc, BIC, and SBC scores using all sediment data has an AIC = -51.3739, an AICc = -50.0405, a BIC = -46.3792, an SBC = -49.9191, an adjusted  $R^2 = 0.6557$ , an RMSE = 0.10574, an F = 11.47, an F = 11.47, an F = 11.47, an F = 11.47, and F = 11.47,

Table E.11. Model statistics for all sediment

Independent variable	Parameter estimate (mg/kg)	Standard error (mg/kg)	Student's t-statistic	Two-sided <i>p</i> -value
Intercept	-0.11258	0.07631	-1.48	0.1742
Chloride	0.01064	0.00332	3.20	0.0108
TOC	2.21E-05	6.47E-06	3.42	0.0076


TOC = total organic carbon

Table E.11 shows chloride and TOC significantly contribute to the model at the 0.05 significance level, while the intercept does not. Figure E.13 is a plot that shows the multivariate-predicted concentrations and 95% prediction limits compared with the actual mercury concentrations. Figure E.13 shows all concentrations are within the 95% prediction limits.

Figure E.14 shows a normal probability plot of the best AIC, AICc, BIC, and SBC model residuals. Model residuals were tested for normality using the SW test, which showed the residuals are approximately normally distributed at the 0.05 significance level, with an SW p-value = 0.9965.

#### E.4.1 SEDIMENT BY PARTICLE SIZE

The particle sizes of the sediment data were used for a correlation analysis. Of the 12 sediment locations, 7 had particle size #10, 1 had particle size #20, 1 had particle size #4, 1 had particle size 3/4 in., and 2 had particle size 3/8 in. Therefore, only particle size #10 had enough samples to perform a correlation analysis. Table E.4 summarizes the correlations of sediment for particle size #10. Combinations that result in correlations that are significant at the 0.05 significance level are highlighted by significance. Correlations that are not highlighted are not significant at the 0.05 significance level.




AICc = Akaike's Information Criterion corrected

SBC = Schwarz Bayesian Criterion

BCV = Bear Creek Valley

Figure E.13. Best AIC, AICc, BIC, and SBC model for all sediment.



Deviates from the Standard Normal Cumulative Distribution Function

AIC Akaike's Information Criterion AICc = Akaike's Information Criterion corrected BIC = Bayesian Information Criterion SBC = Schwarz Bayesian Criterion

Figure E.14. AIC, AICc, BIC, and SBC model residuals for all sediment.

For particle size #10 sediment, barium, cadmium, calcium, cobalt, copper, manganese, nickel, selenium, strontium, zinc, and TOC were significantly positively correlated with mercury at the 0.05 significance level, while antimony was significantly negatively correlated with mercury at the 0.05 significance level. Table E.3 shows chloride, calcium, selenium, sodium, mercury (F1), mercury (FS), and TOC were all significantly positively correlated with mercury for all sediment.

### E.5. CONCLUSIONS

Correlation analysis was performed for all pairs of analytes for soil and sediment where at least one-half of the samples were detected and  $n \ge 3$ . Those analytes with significant correlation with mercury were used in multivariate regression models. Four information criteria methods were used to determine the best multivariate models to predict mercury in soil, sediment, creek bank soil, and floodplain soil.

Correlations for particle size #10 show different analytes significantly correlate with mercury compared to all soil and all sediment.


Fluoride, copper, lead, lithium, zinc, calcium, uranium, and vanadium are significant predictors of mercury for all soil combined. Fluoride, lead, selenium, and uranium are significant predictors of mercury in floodplain soil. Calcium, copper, strontium, and zinc are significant predictors of mercury in creek bank soil. Chloride and TOC are significant predictors of mercury in sediment.


### **E.6. REFERENCES**


- Akaike, H. 1973. "Information theory and an extension of the maximum likelihood principle." In B.N. Petrov and F. Csaki (Eds.), *Second international symposium on information theory*, 267-281. Budapest: Academiai Kiado.
- Akaike, H. 1987. "Factor analysis and AIC," Psychometrika, 52:317-332.
- Beal, D.J. 2007. "Information criteria methods in SAS® for multiple linear regression models," *Proceedings of the 15th Annual Conference of the SouthEast SAS Users Group.*
- Bozdogan, H. 1987. "Model selection and Akaike's information criterion (AIC): the general theory and its analytical extensions," *Psychometrika*, 52, No. 3, 345-370.
- Bozdogan, H. 2000. "Akaike's information criterion and recent developments in informational complexity," *Journal of Mathematical Psychology*, 44:62-91.
- Helsel, D.R. 2005. Nondetects and Data Analysis: Statistics for Censored Environmental Data, John Wiley & Sons, Inc.
- Sawa, T. 1978. "Information criteria for discriminating among alternative regression models," *Econometrica*, 46:1273-1282.
- Schwarz, G. 1978. "Estimating the dimension of a model," *Annals of Statistics*, 6:461-464.


# APPENDIX F. WATER RESOURCES RESTORATION PROGRAM SURFACE WATER TOTAL MERCURY AND METHYLMERCURY LONGITUDINAL DATA PLOTS FOR BEAR CREEK


	ROGRAM SUR FA PLOTS FOR	FACE WATER BEAR CREEK
















WATER RESOURCES RESTORATION PROGRAM SURFACE WATER TOTAL METHYLMERCURY LONGITUDINAL DATA PLOTS FOR BEAR CREEK











# APPENDIX G. BEAR CREEK FIELD OBSERVATIONS




Figure G.1. BCV transect locations and 100-year floodplain.




Figure G.2. BCV transect area 1 and 100-year floodplain.




Figure G.3. BCV transect area 2 and 100-year floodplain.




Figure G.4. BCV transect area 3 and 100-year floodplain.




Figure G.5. BCV transect area 4 and 100-year floodplain.

Table G.1. Field observations of Bear Creek during transect sampling

Site identifier	Stream morphology
BCT1	<ul> <li>Stream is 384 in. wide</li> <li>Stream depth is 14.5 in.</li> <li>Sandbar extends 138 in. into the stream</li> <li>Bank is 30 in. above the water's surface</li> </ul>
BCT2	- Stream is 248 in. wide - Banks are 360 in. apart - Gravel bars are present along the edges of the water - Bank is 50 in. above the water's surface - Stream is straight and has moderate flow
ВСТ3	-Stream is 192 in. wide - Stream depth is 13 to 36 in Banks are extremely steep - Channel is straight and has moderate flow - Soil/sediment samples are collected approximately 50 yd downstream of culvert due to banks being riprap/exposed bedrock
BCT4	<ul> <li>Stream is 264 in. wide</li> <li>Stream depth is 20 to 42 in.</li> <li>South bank is 12 in. above the water's surface; north bank is 4 to 6 in. above the water's surface</li> <li>Stream is meandering and has slow flow</li> <li>Due to ponding at beaver dam, samples collected approximately 150 yd upstream of the planned location</li> </ul>
BCT5	<ul> <li>Stream is 216 in. wide</li> <li>Stream depth is 27 in.</li> <li>Bank is 24 in. above the water's surface</li> <li>Gravel bar extends 36 in. into the channel at the sample point</li> <li>Stream has slight curves with high flow</li> <li>Beaver dam near this location was blown out/destroyed, presumably from extreme flow from recent storm event</li> </ul>
ВСТ6	<ul> <li>Stream is 348 in. wide</li> <li>Stream depth is 18 in.</li> <li>Bank is 24 in. above the water's surface</li> <li>Debris bank (gravel/rock/sediment, sticks/wood, or leaf litter) extends 96 in. into the channel at the sample point</li> <li>Stream has slight curves with high flow</li> <li>Beaver dam near this location was blown out/destroyed, presumably from extreme flow from recent storm event</li> </ul>

Table G.1. Field observations of Bear Creek during transect sampling (cont.)

Site identifier	Stream morphology
BCT7	<ul> <li>Stream is 324 in. wide</li> <li>Stream depth at middle of channel is unobtainable due to ponding/depth/safety (estimated to be at least 60 in. deep)</li> <li>No true bank, only floodplain, due to ponding</li> <li>Stream is meandering and has slow flow</li> <li>Site is directly below beaver dam</li> </ul>
ВСТ8	<ul> <li>Stream is 396 in. wide</li> <li>No defined bank, only floodplain, due to ponding</li> <li>Edges of channel are 24 in. deep, and the deepest point is at least 60 in. (too deep to safely measure)</li> <li>Channel is straight and has slow flow</li> <li>Samples collected upstream of beaver dam at Haul Road intersection</li> </ul>
ВСТ9	<ul> <li>Stream is 456 in. wide</li> <li>Edges of channel are 6 in. deep. Stream depth at middle of channel is unobtainable due to depth/safety concerns</li> <li>No defined bank due to ponding</li> <li>Channel is straight and has slow flow</li> <li>Site is above beaver dam</li> </ul>
BCT10	<ul> <li>Stream is 216 in. wide</li> <li>Banks are 304 in. apart</li> <li>Stream depth is 10 in.</li> <li>Debris (gravel/rock/sediment, sticks/wood, or leaf litter) buildup is 82 in. wide</li> <li>Banks are 30 in. above the water's surface</li> <li>Channel is straight and has high flow</li> <li>Flow had decreased (post-storm event) and stream depth was 6 in. during a second sampling event</li> </ul>
BCT11	<ul> <li>Stream is 180 in. wide</li> <li>Stream depth at sample point is 8 in., but pools that reach 42 in. deep are present</li> <li>Bank is 12 in. above the water's surface</li> <li>Debris (gravel/rock/sediment, sticks/wood, or leaf litter) buildup is 48 in. wide</li> <li>Channel is curvy and has slow flow due to debris buildup</li> </ul>
BCT12A	<ul> <li>Stream is 60 in. wide</li> <li>Banks are 132 in. apart</li> <li>Stream depth of sample point is 6 in.; however, deeper pockets exist upstream and downstream that are 12 in. deep</li> <li>Gravel/sand bar is 72 in. wide</li> <li>Bank is 50 in. above the water's surface</li> <li>Channel is straight and has high flow</li> <li>Flow had decreased (post-storm event) and water depth was 4 in. during a second sampling event</li> </ul>

Table G.1. Field observations of Bear Creek during transect sampling (cont.)

Site identifier	Stream morphology
BCT12B	<ul> <li>Stream is 96 in. wide</li> <li>Banks are 168 in. apart</li> <li>Stream depth at sample point is 10 in.</li> <li>Gravel/sand bar is 72 in. wide</li> <li>Channel is straight and has high flow</li> <li>Bank is 60 in. above the water's surface</li> <li>Flow had decreased (post-storm event) and water depth was 6 in. during a second sampling event</li> </ul>
BCT13	<ul> <li>Stream is 36 in. wide</li> <li>Banks are 96 in. apart</li> <li>Stream depth is 8 in.</li> <li>Banks are 30 in. above the water's surface</li> <li>Channel is curvy and has high flow</li> <li>Flow had decreased (post-storm event) and water depth was 4 in. during a second sampling event</li> </ul>
BCT14	<ul> <li>Stream is 120 in. wide</li> <li>Water is 16 in. deep at the sample point directly following a dropoff in the channel</li> <li>Bank is 72 in. above the water's surface</li> <li>Channel is straight and has moderate flow</li> </ul>
BCT15	<ul> <li>Stream is 36 in. wide</li> <li>Stream depth is 4 in.</li> <li>Bank is 24 in. above the water's surface</li> <li>Stream is meandering with a moderate/average flow rate for that location</li> </ul>
HCTREF (Hinds Creek transect reference site)	<ul> <li>Stream is 372 in. wide</li> <li>Stream depth is 13 in.</li> <li>Bank is 24 in. above the water's surface</li> <li>Stream is straight and has fast flow</li> <li>Bottom of the channel is exposed bedrock with small pockets of gravel</li> </ul>

BCT = Bear Creek transect

This page intentionally left blank.

# DOE/OR/01-2977&D2

### RECORD COPY DISTRIBUTION

File—DMC—RC



Document Number:	Document Title:		
DOE/OR/01-2977&D1	Bear Creek Valley Mercury Sources Remedial Site Evaluati	ion for the U.S. Department of Energy Oak Ridge Site Oak Ridge, Tennessee	
Name of Reviewer:	Organization:	Date Comments Transmitted:	
John W. Sayer	U.S. Environmental Protection Agency	December 6, 2024	

John W. Sa	yer	U.S. Environmental Protection Agency	December 6, 2024	
Comment No.	Sect./Page	Comment	Response (02/04/2025)	Revised Response (04/10/2025) – Changes highlighted in yellow are based on the 04/22/2025 comment response meeting and subsequent email correspondence revisions are highlighted in green
			General Comments	
		The purpose of conducting the RSE as part of the Mercury Management Approach is stated in the Record of Decision (ROD) for the Environmental Management Disposal Facility (EMDF) as part of the Mercury Management Approach for Bear Creek. The ROD states that "DOE shall make efforts to restore Bear Creek to attain full compliance with recreational use designation, including conducting a Remedial Site Evaluation (RSE) (40 CFR 300.420) to evaluate mercury methylation in Bear Creek and conduct pilot or treatability studies as needed." However, the RSE provides little evaluation of methylation, or how the results shed more light on the methylation process in Bear Creek. Discussion on the study of methylation is limited to referencing the Bear Creek Studies Report 2021 (ORNL/SPR-2021/2162), without sharing the results of the studies or how those results affect the interpretation of the data in the RSE. Please revise the RSE to address this data gap.	Clarification. As stated in the DQO meeting held on June 29, 2023, and as described in Section 3.1 of the BCV mercury sources RSE, the overall problem that would be addressed by the RSE is that there are insufficient data along Bear Creek and its tributaries to determine if there are potential sources of mercury and methylmercury in channel sediment and creek bank and floodplain soils that may be contributing to exceedances of fish tissue criterion in prior years. As such, the two main goals of the BCV mercury sources RSE (Section 3.2) were to:  • Determine if there are areas (channel sediment and creek bank and floodplain soils) along Bear Creek and its tributaries that are potential sources of mercury and methylmercury that may affect fish.  • Obtain data from various hydrologic settings (pools, beaver ponds, etc.) that may contribute to mercury methylation and its bioaccumulation in the environment of Bear Creek and a reference location (e.g., Hinds Creek).  The BCV mercury sources RSE fulfills the scope included in the project DQOs and adds significant data to the evolving conceptual model of mercury in the Bear Creek floodplain and aquatic system. The understanding of mercury methylation in Bear Creek is incomplete and will require an ongoing long-term investigation to also inform development of the mercury bioaccumulation conceptual model for Bear Creek.  It is recognized that additional investigation is needed to better understand the mercury methylation in Bear Creek. These investigations will continue through the mercury remediation technology development program; however, given the significant scope of understanding the mercury methylation, additional studies on mercury methylation in Bear Creek will be a long-term investigation.  No change to the document proposed.	Clarification. The last paragraph of Section 5.1 (Conclusions) was revised as follows:  "Overall, mercury contamination of surface water, floodplain and creek bank soil, and channel sediment is highest in the upper reaches of Bear Creek, with decreasing concentrations downstream. The increasing mercury flux in the downstream flow direction is dominated by the increasing flow volume, as concentration data tend to gradually decrease. This characteristic suggests mercury entering Bear Creek downstream of the known source associated with the S-3 Ponds area is derived from dispersed mercury that may be associated with secondary contamination of creek bank and floodplain soil. Although mercury is detected at low levels in sampled media, concentrations are much lower than other ORR mercury-contaminated sites and fish concentrations continue to decrease. This BCV mercury sources RSE did not identify a source of mercury in the media collected in and along Bear Creek that indicates active remediation or an RI of mercury sources is needed. However, the presence of mercury in fish tissue indicates additional studies to understand methylation in Bear Creek are necessary to further refine the CSM."  In addition, Section 5.2 (Recommendations) was revised as follows:  "While the historical water quality and discharge measurement dataset is extensive in Bear Creek, there are far fewer coordinated water-quality-plus-discharge-measurement campaigns with a specific emphasis on total versus dissolved mercury and methylmercury. Although the baseflow analysis of total versus dissolved mercury and methylmercury. Although the baseflow analysis of total versus dissolved mercury and methylmercury were particle-associated, there are no event-based (i.e., precipitation-driven flows), coordinated, discharge and water quality sampling campaigns. To assess total and dissolved fluxes of mercury and other solutes in addition to those available at established stations in Zone 3, the following additional coordinated flow measurement and sampling



Comment No.	Sect./Page	Comment	Response (02/04/2025)	Revised Response (04/10/2025) – Changes highlighted in yellow are based on the 04/22/2025 comment response meeting and subsequent email correspondence revisions are highlighted in green
				also the base of the food web, serving to concentrate methylmercury concentrations in the dietary pathways leading to fish, future monitoring should focus on periphyton dynamics.
				Although periphyton are understood to play an important role in mercury methylation in stream systems, how the periphyton community changes spatially and temporally in Bear Creek and how these changes relate to methylmercury concentrations in the stream are yet to be understood. Recommendations to evaluate periphyton in Bear Creek in support of the CSM under the mercury remediation technology development program follow:
				<ul> <li>Based on guidance provided by Bravo et al. (2017), evaluating organic material characteristics that may affect mercury methylation and biouptake is required to understand the mixes of organic compound origins and molecular weights and to inform attempts to discern potential aquatic habitat characteristics that influence mercury methylation, accumulation in media, and bioaccumulation. This information is critical to any best management watershed-scale practices aimed at reducing methylmercury concentrations in fish.</li> </ul>
				<ul> <li>To gain a better understanding of mercury methylation for decision-making, data from the aforementioned continuous monitoring instruments should also be supplemented with manual sampling data for measures of periphyton abundance and community structure.</li> </ul>
				• To understand spatial and temporal periphyton dynamics over time, deploying nutrient (e.g., the nitrate sensor at the BCK 1 monitoring station), light, and turbidity sensors to augment the discharge and water quality measurements collected at additional future monitoring station installations is required. The continuous measurement of periphyton- and mercury-relevant water quality parameters may also be enriched by installing relatively low-cost field cameras capable of estimating aquatic vegetation distribution and abundance in proximity to the monitoring station on a daily basis. Nutrients and light are both important drivers of periphyton distribution and abundance, along with temperature, depth, velocity, and substrate type. Integrated telemetry and web-based data portals to allow these contemporaneous measurements, continuous monitoring stations, field cameras, and stream conditions to be viewed in near real-time would also improve data accessibility and usability and add a new capability to the toolset used to assess the BCV Watershed.
				Additional BCV investigations under the mercury remediation technology development program will be documented in a report that is updated annually. The initial document will be issued as a secondary DOE document with an associated FFA Appendix E milestone that presents the results of historical studies, the current CSM for mercury methylation in BCV, and the data gaps that will be addressed in future studies to support further development of the CSM. An annual addendum to that report (also with an FFA Appendix E milestone [i.e., the ORNL Technical Memorandum]) will be issued and will summarize completed activities and additional investigation recommendations based on the results of those activities. The report will be appended to the annual EMDF Phased Construction Completion Report. An annual roundtable meeting will be held with the FFA parties (DOE, EPA, and TDEC) to provide the status of the investigations and to discuss recommendations for additional studies. While no remedial action is required at this time, the scientific investigations proposed may indicate an opportunity to reduce mercury in Bear Creek in the future. DOE will determine if Bear Creek is in compliance with applicable water quality standards prior to EMDF operations. If Bear Creek is determined to not be in compliance, then DOE commits to a schedule of actions as approved by the FFA parties (DOE, EPA, and TDEC)."
				The last paragraph of the Executive Summary was revised as follows:
				"Although this BCV mercury sources RSE determines no remedial investigations or actions are required at this time, enhanced monitoring of instream mercury and methylmercury concentrations and flux rates and further development of the mercury bioaccumulation conceptual model continue under the mercury remediation technology development program. Recommendations for future technology development program investigations to enhance the understanding of the mercury bioaccumulation conceptual model are provided. While no remedial action is required at this time, the



Comment No.	Sect./Page	Comment	Response (02/04/2025)	Revised Response (04/10/2025) – Changes highlighted in yellow are based on the 04/22/2025 comment response meeting and subsequent email correspondence revisions are highlighted in green
				scientific investigations proposed may indicate an opportunity to reduce mercury in Bear Creek in the future, and CERCLA actions will be taken at that time as appropriate."
				The BCV mercury sources RSE focused on key data gaps that are not addressed by the mercury remediation technology development program. However, some additional detail of these long-term investigations that are conducted by ESD were added to Section 1.2.1.2 as follows:
				"A 2022 data compilation report for mercury in Bear Creek (ORNL/TM-2023/3069) summarized data from compliance and investigatory studies to begin building a conceptual model to understand the processes affecting mercury transport and transformation in the Bear Creek Watershed and to highlight key knowledge gaps in understanding these processes. This report summarizes historical data from compliance and investigatory studies relevant to mercury contamination and transformation in the Bear Creek Watershed and highlights the relative paucity of water quality data in Bear Creek with respect to mercury concentrations and transformation processes. A recommendation from this study was to install a long-term monitoring station to provide needed data to establish functional linkages among identified components of a conceptual model of mercury transport with measured mercury values from the watershed.
				The mercury remediation technology development program at ORNL, which began in 2014, originally focused on understanding and addressing mercury concentrations, flux, and bioaccumulation in EFPC. Research from this program highlighted the importance of periphyton in mercury transformation in stream systems, and in 2021, a task specifically focused on periphyton dynamics and distribution was added to the study plan. Because the aqueous methylmercury concentrations in EFPC are similar to those in nearby Bear Creek despite significantly higher aqueous total mercury concentrations in EFPC, comparing mercury methylation and demethylation processes between the two streams was of particular interest, and Bear Creek became (and remains) a point of focus in the mercury remediation technology development program (ORNL/SPR-2023/3178).
				Future work will build"
				Note that the following reference was added to Chapter 6:
				"ORNL/SPR-2023/3178. Mercury Remediation Technology Development for Lower East Fork Poplar Creek FY 2023 Update, 2023, Oak Ridge National Laboratory, Oak Ridge, TN."
2		There is a misalignment between the stated objective of the RSE and the focus of the report. Section 1.1 (Remedial Site Evaluation Objective) indicates that the primary goal of the RSE is to "evaluate potential sources of mercury and methylmercury within the Bear Creek Valley (BCV) Watershed." However, much of the report focuses on the results of water, soil, and sediment sampling and does not provide sufficient discussion on how these data contribute to the identification and evaluation of specific source areas for mercury and methylmercury. For instance, while the RSE details elevated mercury concentrations in certain transects, the RSE lacks a clear analysis that explains whether and how these concentrations correlate with known or suspected source areas. Please revise the RSE to more explicitly discuss how the data relates to the evaluation of mercury and methylmercury sources within BCV.	Clarification. Please see response to EPA general comment 1. No change to the document proposed.	Clarification. Please see the revised response to EPA general comment 1.
3		The RSE states in Section 1.2.1.2 (BCV Watershed Hydrogeological Conceptual Model), that a 2022 data compilation report for mercury in Bear Creek (ORNL/TM-2023/3069) summarized data from data collection studies in order to understand the	Clarification. The BCV mercury sources RSE is a key part of understanding the mercury bioaccumulation conceptual model for Bear Creek. As stated in Section 1.2.3 of the BCV mercury sources RSE, prior to this investigation, there were limited sediment data in Bear Creek and no sediment methylmercury data. The BCV mercury	



Comment No. Sect./Page	Comment	Response (02/04/2025)	Revised Response (04/10/2025) – Changes highlighted in yellow are based on the 04/22/2025 comment response meeting and subsequent email correspondence revisions are highlighted in green
	processes affecting mercury transport and transformation in the Bear Creek Watershed and to highlight key knowledge gaps in understanding these processes. This section also states future work was intended to build on the summary presented in the data compilation report to develop a conceptual model that outlines the key environmental parameters that correlate with methylmercury concentrations and bioaccumulation in Bear Creek. The conceptual model was to be used to provide the technical basis for prioritizing new data collection and optimizing potential mitigation actions or best management practices, with the goal of lowering fish tissue mercury concentrations. However, the RSE does not state how, or if previous data was used to develop the sampling strategy for the RSE, and does not provide the conceptual site model (CSM) that was described as being in development. Please revise the RSE to include a CSM and to state what remaining data gaps still exist based on an evaluation of all previously collected data, including this most recent data collected as part of the RSE.	sources RSE has provided sediment data that were previously unavailable.  A key finding of the BCV mercury sources RSE comes from the data provided through the mercury sequential extraction analyses that document the majority of the mercury in BCV floodplain soil, creek bank soil, and channel sediment is associated with organic components of the media.  Refer to the response to EPA general comment 1. No change to the document proposed.	
4	The RSE notes that Bear Creek has the highest concentrations of mercury near NT-3, where several historical land disposal sites are located. These land disposal areas include the Bone Yard/Burn Yard, the Oil Landfarm, S-3 Ponds, Sanitary Landfill 1, Bear Creek Road Debris Burial site, and Creekside Debris Burial. The RSE outlines previous remedial investigation/remedial activities conducted for these sites where mercury was found to be present in the soils and/or groundwater. For example, Section 1.2.2 (Summary of Potential Mercury Source Areas), states that the previously issued Remedial Investigation Report on Bear Creek Valley Operable Unit 2 (Rust Spoil Area, Spoil Area 1, and SY-200 Yard) at the Oak Ridge Y-12 Plant identified mercury as a contaminant of concern (COC) at the SY 200 Yard, and states that a soil cover was placed over this area. This section also states that free-phase mercury was identified in soil borings in this area. The RSE does not discuss, however, whether the SY 200 Yard or any of these other historical waste disposal areas/sites near NT-3 still only a partial or complete soil cover rather than a Resource Conservation and Recovery Act (RCRA)-like cap. This information is useful for identifying which areas may still be contributing mercury to Bear Creek (either through runoff, erosion, or infiltration/groundwater recharge). Please revise the RSE to describe which of the land disposal areas may still have a partial or complete soil cover as the remedy for the land disposal units.	Clarification. New Table 1.1 has been added to Section 1.2.2 to describe the caps and soil covers at associated BCV disposal areas. A new paragraph was added to the text at the end of Section 1.2.2 as follows:  "Many of the disposal sites have caps or soil covers that are maintained as described in Table 1.1."  New Table 1.1 is included in Attachment 1 at the end of these comment responses.	



Comment No.	Sect./Page	Comment	Response (02/04/2025)	Revised Response (04/10/2025) – Changes highlighted in yellow are based on the 04/22/2025 comment response meeting and subsequent email correspondence revisions are highlighted in green
5		The RSE does not discuss the correlation between methylation of mercury and the presence of phytoplankton. Previous mercury source investigation project team presentations proposed that phytoplankton was a major contributor to the generation of methyl mercury. Please revise the RSE to state whether the RSE data provided any further evidence to support the idea that phytoplankton contributed to the methylation of mercury in Bear Creek and whether higher concentrations of phytoplankton correlated with higher levels of methyl mercury in Bear Creek.	Clarification. Please refer to the responses to EPA general comments 1 and 3. No change to the document proposed.	
6		It is unclear whether additional source area soil investigations are warranted. Section 1.2.2 (Summary of Potential Mercury Source Areas) notes that free mercury was observed in some borings during the BCV Operable Unit (OU) 2 Remedial Investigation (RI) at SY-200 Yard; Section 3.2.2 (Soil and Sediment Chemical Characteristics) reports a maximum mercury concentration of 3500 micrograms per kilogram (µg/kg) in floodplain soil at BCT12A and 7100 µg/kg in creek bank soil at BCT13 (near suspected source areas). Also, Section 5.1 (Conclusions) indicates that floodplain and creek bank mercury concentrations were highest in the stretch downstream from BCK 12.34 and upstream of NT-3 (proximal to source areas). Considering the detections of mercury in soils near, but not directly within, known source areas, it may be necessary to conduct additional soil investigations within source areas to determine the potential migration of contaminants and provide a complete understanding of how mercury is being transported and whether additional mitigation or remediation efforts are necessary. Please revise the RSE to discuss whether further soil sampling and evaluation to fully assess the risk and extent of contamination from these source areas is warranted.	Clarification. The BCV mercury sources RSE SAP identified the SY-200 Yard as a potential source of mercury in BCV, as also described in Section 1.2.2 of the BCV mercury sources RSE. Based on this potential source of mercury to Bear Creek, transect locations were selected along Bear Creek in the vicinity of the SY-200 Yard (transects BCT14 and BCT15). Prior to the BCV OU 2 ROD that included the SY-200 Yard, a soil cover of 3 to 5 ft was placed at SY-200. The BCV OU 2 ROD did not require implementation of additional RAs for SY-200. The selected remedy from the BCV OU 2 ROD relies on maintaining the soil cover and LUCs for SY-200.  No change to the document proposed.	
7		Potential remaining source areas are the S-3 Ponds, the SY-200 Yard, and the Boneyard/Burnyard (BYBY). Although groundwater was not included in the scope of the Remedial Site Evaluation (RSE), the EPA considers groundwater a potential source. Potential sources are the groundwater at the BYBY, the groundwater southwest of the S-3 Ponds, and the groundwater northwest of the SY-200 Yard. The groundwater at the BYBY was historically considered a potential source to North tributary 3 (NT-3) for which a hydraulic isolation remedy was implemented. OREIS data shows concentrations were orders of magnitude above the surface water standard for mercury at BY-06, BY-11, BY-29, and BY-39 (Comment Figure 1). Groundwater northwest of the SY-200 Yard ranged from 11 µg/L at GW-835 to 32	Clarification. Please see response to EPA general comment 1. No change to the document proposed.	



Comment No.	Sect./Page	Comment	Response (02/04/2025)	Revised Response (04/10/2025) – Changes highlighted in yellow are based on the 04/22/2025 comment response meeting and subsequent email correspondence revisions are highlighted in green
		μg/L mercury at GW-834 (Comment Figure 2). In addition to groundwater at the BYBY, southwest of the S-3 Ponds historical data show up to 110 ug/L of mercury at GW-243 (Comment Figure 3). Although the RSE detected the highest mercury concentrations in surface water downstream of Station BCT15 at stations BCT-13 and BCT-12A, groundwater plumes from S-3 Ponds could be a lingering source. Consider developing a work plan to investigate mercury in the groundwater as a potential source to the karst system of subsurface channels and seeps.		
8		It is unclear whether any investigation of lower trophic organisms has been conducted as the RSE provides fish tissue biological data only. Since mercury methylation often occurs in sediments, benthic organisms that live in or near these sediments are particularly susceptible to bioaccumulation and may contribute to mercury concentrations in higher trophic organisms. These organisms can also serve as early warning bioindicators to determine whether additional mitigation measures are necessary. Additionally, because benthic organisms are spatially distributed and more abundant than fish, they may provide better insight into source area contamination and allow for a more targeted remediation approach, if needed. Please revise the RSE to discuss whether mercury and methylmercury investigations focused on lower trophic organisms have been conducted, and whether this type of study could be beneficial for the identification of source areas.	Clarification. Please see response to EPA general comment 1. No change to the document proposed.	Clarification. Please see the revised response to EPA general comment 1.
9		Upstream to downstream mercury concentrations in fish reflect similar distributions in surface water. For example, central stonerollers at upstream monitoring station BCK 12.4 have higher mercury than fish at BCK 9.9. Surface water monitoring shows an upstream to downstream decreasing trend in total mercury concentrations from BCK 12.34 to BCK 11.54 (Figure 1.4) with typically higher concentrations at SS-4 and upstream (Appendix F). The April RSE sampling event was considered the most representative of upstream to downstream trends due to variable flows during the January event. The highest mercury concentration in unfiltered surface water (22 ng/L) in January 2024 sampling event was collected upstream of NT-3 at BCT13 and not located immediately downstream of a known source. The highest concentration of mercury in filtered surface water (4.3 ng/L) was collected both upstream and at the confluence of NT-3 and Bear Creek in the January 2024 event. Channel sediments in the RSE had relatively high concentrations of mercury at upstream station BCT15 (430 ug/kg) but did not show as great of an upstream to downstream decrease as did surface water. The upstream to downstream pattern	Comment noted. No change to the document proposed.	



Comment No.	Sect./Page	Comment	Response (02/04/2025)	Revised Response (04/10/2025) – Changes highlighted in yellow are based on the 04/22/2025 comment response meeting and subsequent email correspondence revisions are highlighted in green
		suggests potential sources in Zone 3 including surface soils adjacent to the creek.		
10		The pattern of having relatively high mercury and methylmercury concentrations in surface water at BCT13 suggests the persistence of a mercury source upstream of NT-3. Also, the creek bed sediments, creek bank soils, and floodplain soils had higher mercury concentrations at NT-3 and BCT13. The maximum detected mercury concentration in sediment was downstream of the hazardous chemical disposal area (HCDA) at BCT13. Excavated soils within the footprint of the BYBY had mercury concentrations above approximately 31 mg/kg. The remaining BYBY soils were covered with 6 inches of fill. Some soils adjacent to Bear Creek might not have been covered (Comment Figure 4). Additional characterization of mercury in floodplain soils that were not covered by 6 inches of clean fill is recommended particularly in the vicinity of the BYBY.	Clarification. Mercury concentrations in surface water in NT-3 are influenced by sampling event stream turbidity levels and probably by suspended solids levels. DOE has updated the WRRP BCV annual sampling to collect field parameters along with field-filtered and unfiltered surface water samples in future sampling events to allow interpretation of dissolved versus particle-associated mercury transport. The current BCV mercury sources RSE document provides an important data addition to the evolving BCV mercury conceptual model. No change to the document proposed.	
11		A soil sample upstream of the BYBY in OREIS had a mercury concentration of 165 mg/kg in the top 2 inches at location K-MAN-YS1454 near Building 9721-73 northeast of the Y-200 Scrap Yard (Comment Figure 5). The soils around this location were not delineated towards the creek and could present a source.	Comment noted. No change to the document proposed.	
12		Soils upstream of BCT15 and southeast of the former S-3 Ponds in the vicinity of Bear Creek measured 141 mg/kg in the top 2 inches at sampling location K-MAN-YB0325 and 31 mg/kg in the top 2 inches at sampling location K-MAN-YB0334. Two other S-3 Pond samples with elevated mercury in soils were sampling locations S3-03 and S3-08, with 82.9 mg/kg and 107 mg/kg, respectively (Comment Figure 6). It should be noted that the concentrations in S3-03 and S3-08 were composites over the top 4 feet and might have underrepresented the concentrations in surface soils. Please revise the document to support conclusions regarding the absence of specific sources in floodplain soils upstream of NT-3.	Clarification. Long-term performance and baseline monitoring of groundwater and surface water is evaluated on an annual basis in the RER. Known residual mercury in the vicinity of the S-3 Ponds plume may contribute to mercury detections in Bear Creek (see Attachment 2 at the end of these comment responses).  Note that response actions for the S-3 Ponds and for BCV groundwater have yet to be implemented and will be included in the future design considerations for Pathway 3 under the BCV Phase I ROD or in the final groundwater decision for BCV.  No change to the document proposed.	
13		The RSE does not provide benchmark sediment screening levels to enhance the understanding of potential risks to aquatic receptors from mercury. While the RSE employs a site-specific risk assessment approach and does not rely on initial benchmark screening levels for a direct comparison, it would be beneficial to include sediment-specific screening levels for general comparison. For instance, the Chemical Aquatic Fate and Effects (CAFÉ) database may provide relevant screening criteria for mercury in sediments which could serve as a useful comparison for understanding risk to aquatic receptors. Please revise the RSE to include	Clarification. The BCV mercury sources RSE is a source investigation based on the mercury management approach outlined in the EMDF ROD and is not a risk assessment. The mercury management approach in the EMDF ROD specifies mercury levels in water and fish to be met. Thus, the BCV mercury sources RSE focuses on these benchmarks. To provide an understanding of the magnitude of contamination in the sediment and associated risk, the BCV mercury sources RSE provides levels of mercury in Bear Creek sediment and discusses mercury partitioning in comparison to LEFPC. No change to the document proposed.	



Comment	Coat /D	Comment	Decisiones (00/04/0005)	Revised Response (04/10/2025) – Changes highlighted in yellow are based on the 04/22/2025
No.	Sect./Page	Comment	Response (02/04/2025)	comment response meeting and subsequent email correspondence revisions are highlighted in green
		benchmark screening levels for sediment in order to provide a better understanding of the risk associated with mercury and methylmercury to aquatic receptors.		5
14		The RSE should summarize the results of the data validation and verification and attach the data validation packages as supporting documentation. Section 6.1 (Data Verification and Validation) of Appendix A (Bear Creek Valley Mercury Sources Remedial Site Evaluation Sampling and Analysis Plan) indicates all data will be verified and Level 4 data packages will be required for all analyses. However, data validation is not discussed within the RSE, and data validation packages are not provided as supporting documentation. Please revise the RSE to summarize the data validation findings and include the data validation packages as an appendix.	Clarification. Data validation is performed in accordance with procedures PROC-ES-5005, Sample Management Office Laboratory Data Validation for Inorganic and Organic Analyses, and PROC-ES-5006, Sample Management Office Laboratory Data Validation for Radiochemical Analyses. All data were verified and validated as required per the BCV mercury sources RSE SAP. OREIS provides the data validation qualifiers. No change to the document proposed.	
			Specific Comments	
1	Section 1.2.3, Page 1-8	The mercury concentrations in surface water at NT-3 appear to be increasing or remaining stable in the last five years (Figure 1.4), although the Man-Kendall test did not identify a significant trend. If more recent data are considered, 28 ng/L was detected in May 2023 and 26 ng/L in June 2024. Please revise text on Page 1-8 to explain that while other locations are declining, trends at NT-3 have been stable. Stable trends may indicate a persistent ongoing source. It may be worth investigating the groundwater beneath the BYBY to ensure that there is not an expanding plume towards the creek or tributary (See General Comment 1.).	Clarification. Please see response to EPA general comment 10. No change to the document proposed.	
2	Section 2.1.1, Page 2-6	It is unclear whether the Hinds Creek kilometer (HCK) 20.6 reference site (HCKREF) can be used as a true baseline for comparison. The text indicates sediment sampling or sequential extraction was not conducted at the HCKREF site due to a lack of fine-grained sediment; however, since fine-grained sediment was present at most other transects, it is unclear whether the HCKREF conditions represent downstream conditions. Additionally, fine-grained sediments have a higher capacity for binding mercury, therefore, it is unclear how an accurate assessment of mercury and methylmercury levels relative to baseline conditions can be conducted. Please revise the text to discuss whether HCKREF is suitable for use as a background site if the conditions are not similar to the BCV transects and discuss whether the lack of fine-grained sediment and sequential extraction data impact the assessment conclusions.	Clarification. Hinds Creek is a suitable reference site for Bear Creek. Like Hinds Creek, Bear Creek has portions of the creek (see locations BCT5 and BCT6) that have limited fine-grained sediment. No change to the document proposed.	
3	Section 2.1.2, Page 2-7	The last paragraph of this section states sequential extraction of mercury was conducted at select transect locations, including at BCT5 and BCT6 and at reference site HCTREF; however the previous section 2.1.1 states sequential extraction of mercury	Clarification. Section 2.1.1 is about channel sediment sampling. Note that Section 2.1.2 is about creek bank soil samples where sequential extraction was conducted at locations BCT5, BCT6, and HCTREF. No change to the document proposed.	



Comment No.	Sect./Page	Comment	Response (02/04/2025)	Revised Response (04/10/2025) – Changes highlighted in yellow are based on the 04/22/2025 comment response meeting and subsequent email correspondence revisions are highlighted in green
		was not performed for samples at these locations. Please revise Section 2.1.2 as appropriate to provide consistent information regarding whether mercury extraction analysis was performed at sample locations BCT5, BCT6, and HCTREF.		
4	Section 2.1.5, Page 2-7 and Figure 2.1 Page 2-3	The text states, "Eight transect locations (BCT5, BCT6, BCT7, BCT8, BCT9, BCT11, BCT12B, and BCT14), as well as a reference site (HCTREF [HCK 20.6]), were sampled for mercury speciation/sequential extraction analysis;" however, since BCT5, BCT6 and HCTREF were not sampled the statement should be revised to state, "Eight transect locations (BCT5, BCT6, BCT7, BCT8, BCT9, BCT11, BCT12B, and BCT14), as well as a reference site (HCTREF [HCK 20.6]), were selected for mercury speciation/sequential extraction analysis." Additionally, Figure 2.1 (BCV transect locations) should be updated to clarify BCT5, BCT6 and HCTREF were not sampled for mercury speciation/sequential extraction analysis (i.e., represented by pink dashed lines instead of purple dashed lines). Please revise the quoted statement for clarity and revise Figure 2.1 to use pink dashed lines for each transect that did not have mercury speciation/sequential extraction.	Clarification. Only channel sediment at locations BCT5, BCT6, and HCTREF were not sampled for speciation/sequential extraction analysis. All other planned media (creek bank and floodplain soils) at those locations were sampled for speciation/sequential extraction analysis.  The text in Section 2.1.5 was revised as follows:  "Sufficient mass of solid material from the channel sediment (where available),"	
5	Section 2.2, Page 2-8	The text should discuss whether any efforts were made to retrieve a sample upstream or downstream of the transect when conditions did not allow for sample collection. The text states, "No channel sediment was sampled at BCT1, BCT2, BCT5, BCT6, or the reference site HCTREF (HCK 20.6) because no fine-grained sediment was observed at these locations;" however, it is unclear whether attempts were made to collect the sample from a step-out location (e.g., 20 feet) upstream or downstream. Please clarify whether attempts were made to collect samples from proximate areas.	Clarification. As stated in Section 4.5 of the BCV mercury sources RSE SAP, sample locations were adjusted based on field conditions and sampling viability. Attempts were made to collect these samples 20 to 30 ft in both the upstream and downstream directions as possible based on field conditions. However, sample collection was not feasible at these locations due to limited fine-grained material. No change to the document proposed.	
6	Section 2.2, Page 2-8 and Table 3.6, Page 3-16	The text should elaborate on the decision-making process for not collecting samples at certain locations. Specifically, it states that "no fine-grained sediment was encountered" (Section 2.2) at BCT1, BCT2, BCT5, BCT6, and the reference site HCTREF (HCK 20.6). However, Table 3.6 indicates that fine-grained sediment was present at some of these locations. For example, the description of BCT2-CH reports a combined 28 percent (%) fine sand, silt, and clay. Please revise the text in Section 2.2 to clarify the decision-making process for not collecting sediment samples.	Clarification. The third bullet in Section 2.2 was revised as follows:  "Only particle size analysis samples were collected from channel sediments at BCT1, BCT2, BCT5, BCT6, and the reference site HCTREF (HCK 20.6) due to limited fine-grained sediment observed at these locations. Planned sequential extraction of channel sediment at locations BCT5 and BCT6 and the reference site HCTREF (HCK 20.6) was not performed due to these fine sediment limitations."  Table 3.6 represents the relative proportion of size classes for the sediment that was encountered but does not represent the volume of material available to sample. The substrate for the PSA-only locations was primarily exposed bedrock and/or gravel with fines present as thin veneers or within the void spaces.	



Comment				Revised Response (04/10/2025) – Changes highlighted in yellow are based on the 04/22/2025
No.	Sect./Page	Comment	Response (02/04/2025)	comment response meeting and subsequent email correspondence revisions are highlighted in green
7	Figure 3.1 Page 3-7	Table 3.4 indicates the concentration of UF methylmercury was 0.023J nanograms per liter (ng/L) and filtered methylmercury was 0.025J ng/L at HCTREF; however, Figure 3.1 indicates dissolved mercury was approximately 0.05 ng/L. Please check Table 3.4 and confirm the HCTREF methylmercury total and dissolved concentrations for accuracy and revise Figure 3.1 as needed.	Agree. Figure 3.1 was revised to show the correct dissolved methylmercury for the original sample data at HCTREF. The revised figure is included in Attachment 1 at the end of these comment responses.	
8	Section 3.31, Page 3-41	The text should provide a more thorough evaluation of the correlation analysis presented in Appendix E (Bear Creek Valley Remedial Site Evaluation Analyte Correlations). While the text notes that "pairwise correlations showed many strong and significant correlations among metals, including mercury," it does not explain whether these correlations could be used to predict mercury concentrations or further refine source areas. Please revise the section to discuss whether the strong correlations among metals could be used to predict mercury concentrations and, in turn, help identify potential source areas for mercury contamination.	Clarification. Calculating all pairwise correlations was an exploratory data analysis tool to see which other constituents were strongly correlated with mercury in soil and sediment. Attempting predictions of mercury concentrations in media absent analytical results for mercury was not a goal of the BCV mercury sources RSE. Since the dataset often had more variables (constituents) than concentrations (records), examining these correlations helped narrow down the constituents that might be good predictors for mercury in a multivariate model. Only constituents that were highly correlated with mercury were considered for a multivariate model, which works best when there are many more records than variables. Once these highly correlated constituents were identified, the multivariate model could focus on a select few constituents (typically 6 or fewer) instead of 25 or more.  These pairwise correlations by themselves were not meant to infer that individual constituents could predict mercury alone. But when combined with a few other highly correlated constituents in a multivariate model, there is a goodness of fit of the resulting equation to the modeled dataset in most cases. One location in particular (BCT13) was an exception to the multivariate statistical fit to the soil dataset. No change to the document proposed.	
9	Section 3.2.2, Page 3-13	Mercury concentrations in floodplain soil decrease in the downstream direction. The floodplain soil results are below the soil sampling results available in OREIS downstream of Sanitary Landfill 1, indicating the probable origin was transport from upstream. However, few soil samples near the creek are available between BCT12B and BCT11 to provide additional information on a source area. Additional characterization of mercury in floodplain soils that were not covered by clean fill by previous actions is recommended, particularly in the vicinity of the BYBY to address a potential upstream source (see General Comment 7).	Clarification. Please see response to EPA general comment 1 that describes the problem to be addressed by the BCV mercury sources RSE and the two main goals of the study. Note that transect locations were discussed with EPA and TDEC in the DQO meeting. No change to the document proposed.	Clarification. Please see the revised response to EPA general comment 1.
10	Section 4.1, Page 4-7	Much of the mercury flux is already present in Bear Creek upstream of NT-3 associated with the S-3 Ponds groundwater plume. However, Bear Creek is a losing stream at NT-3 with the waters lost resurfacing at SS-4 and SS-5. Groundwater mercury plumes from the BYBY could travel through the Maynardville Limestone and surface at SS-4 and SS-5. The text indicated flux measured at SS-5 was contributed to by several upstream sources but did not narrow it down to anything specific such as sources in the upstream portion of Zone 3. Please discuss whether the BYBY	Clarification. Please see response to EPA general comment 1. No change to the document proposed.	



Comment No.	Sect./Page	Comment	Response (02/04/2025)	Revised Response (04/10/2025) – Changes highlighted in yellow are based on the 04/22/2025 comment response meeting and subsequent email correspondence revisions are highlighted in green
		and the other waste management areas are likely sources.		
11	Section 5.1, Page 5-1	The text indicates that dissolved mercury levels were highest (1.2 ng/L to 1.9 ng/L) in the stream reach encompassing transect locations BCT14, BCT13, BCT12A, BCT12B, and BCT11, with decreasing concentrations downstream; however, conclusions should be discussed in context to the source areas investigated. These transects are located near known contamination sites, including the Boneyard/Burnyard (BYBY), Hazardous Chemical Disposal Area (HCDA), S-3 Ponds Site, SY-200, and Spoil Area 1; however, there is no discussion of their proximity to the suspected source areas. Additionally, the text should provide conclusions (including context related to source areas investigated) for total mercury and methylmercury, and dissolved methylmercury concentrations.	Clarification. As stated in the comment, the highest dissolved mercury concentration in surface water was 1.9 ng/L, which is well below the AWQC of 51 ng/L.  Please see response to EPA general comment 1 that describes the problem to be addressed by the BCV mercury sources RSE and the two main goals of the study. As stated, the goal was to determine if there are areas (channel sediment and creek bank and floodplain soils) along Bear Creek and its tributaries that are potential sources of mercury and methylmercury that may affect fish.  During the DQO and development of the BCV mercury sources RSE, the transect locations were selected to bias the samples to areas where there may be mercury input into Bear Creek. However, the BCV mercury sources RSE did not identify a source of mercury that significantly contributes to the Bear Creek mercury contamination or that would warrant active remediation. Note that transect locations were discussed with EPA and TDEC at the DQO meeting.  In addition, please see the response to EPA general comment 12. No	
12	Section 5.1, Page 5-1	The text states: "A significant secondary peak of methylmercury was measured in surface water at sample locations BCT9, BCT8, and BCT7, all of which are associated with beaver ponds." Please revise the text to explain the higher methylmercury concentrations surrounding the beaver ponds, and any observable influence from the potential source area sites.	change to the document proposed.  Clarification. RSE transects BCT11 through BCT15 are located in Zone 3, as designated in the BCV Phase I ROD. Zone 3 encompasses the CERCLA legacy waste disposal units and contamination associated with secondary contaminated media, including groundwater (and associated bedrock and soils below the water table), surface water, stream sediment, and overbank floodplain soils. RSE transects BCT01 through BCT10 are located downstream of the Zone 3 contaminant source areas where only secondary mercury contamination is known to occur in and adjacent to the aquatic system. Within Zone 3, there exist extensive Bear Creek stream reaches that experience interflow between the stream surface channel and epikarst groundwater flow pathways that re-emerge to surface flow via spring discharges as well as small macropores and via streambed seepage. As shown in Figure 3.10, methylmercury concentrations in Bear Creek surface water generally increase downstream of Zone 3. Also, as evident in Figure 3.10, there seems to be relatively higher methylmercury in sample locations that are associated with active or past beaver pond habitats. Studies by others (see references below) demonstrate the habitats created by impounded water associated with beaver ponds are conducive to enhance mercury methylation. Precise mechanisms of this habitat-related mercury methylation are not uniquely identified, although higher concentrations of TOC in the quiescent, ponded habitat than in the rapidly flowing stream habitats are cited as contributors. The data obtained in this BCV mercury sources RSE are consistent with conditions documented in other studies. Further long-term investigations are needed to gain additional insights into the biological processes extant in Bear Creek that promote the enhanced mercury methylation several kilometers removed from primary mercury inputs into the aquatic system.	



Comment No.	Sect./Page	Comment	Response (02/04/2025)	Revised Response (04/10/2025) – Changes highlighted in yellow are based on the 04/22/2025 comment response meeting and subsequent email correspondence revisions are highlighted in green
			downstream#:~:text=Scientists%20in%20Sweden%20have%20shown,b ackground%20levels%20during%20summer%20months. https://bioone.org/journals/wildlife-biology/volume-2020/issue-	
			3/wlb.00678/The-effect-of-beaver-dams-on-organic-carbon-nutrients- and/10.1111/wlb.00678.full	
			The following text was added to Section 5.1:	
			"A significant secondary peak of methylmercury was measured in surface water at sample locations BCT9, BCT8, and BCT7, all of which are associated with beaver ponds. Data collected in June and August 2017 and in August 2018 (ORNL/SPR-2018/902) suggest the beaver dams located along Bear Creek are promoting conditions favorable for methylmercury generation."	
13	Section 5.2, page 5-2	The text states: "Despite low aqueous mercury concentrations in Bear Creek, the fish fillet concentrations are relatively high, likely due to high methylation efficiency and efficient trophic transfer." Please revise the text in Section 5.3 to discuss what additional studies are proposed to understand the methylation efficacy and processes trophic transfer accounting for the high fish fillet concentrations. Please provide an explanation if no such studies are planned.	Clarification. Please see response to EPA general comment 1. No change to the document proposed.	Clarification. Please see the revised response to EPA general comment 1.
14	Section 5.2, Page 5-2	The text states: "Heavy sampling in this creek has led to a decline in fish numbers and sizes in 2023; the fish collected in 2023 were smaller than average, which could affect mercury concentrations. If populations recover and/or habitat changes in the creek to facilitate mercury methylation, concentrations in fish may increase above AWQC. Future monitoring will determine whether fish mercury concentrations remain low as fish populations recover and whether the overall decreasing trend observed throughout the stream continues." As stated, smaller fish are less likely to bioaccumulate mercury and so are less representative of tissue mercury levels then when the fish population returns to a more natural size distribution. Please revise the text in Section 5.3 to discuss what alternative biological studies other than additional fish sampling, are planned to supplement the fish sampling data. Please provide an explanation if no such studies are planned.	Clarification. Please see response to EPA general comment 1. No change to the document proposed.	Clarification. Please see the revised response to EPA general comment 1.
15	Section 5.2, Recommendati ons Page 5-2	The RSE does not discuss whether there are additional actions or monitoring measures that should be taken to further facilitate the mercury-management approach for Bear Creek. For example, the Executive Summary states, "Although this BCV mercury sources RSE determines no action is required, enhanced monitoring of instream mercury and methylmercury concentrations and flux rates and further development of the mercury bioaccumulation conceptual model continue under the mercury technology development program;" however, the	Clarification. Please see response to EPA general comment 1. No change to the document proposed.	Clarification. Please see the revised response to EPA general comment 1.






Comment No.	Sect./Page	Comment	Response (02/04/2025)	Revised Response (04/10/2025) – Changes highlighted in yellow are based on the 04/22/2025 comment response meeting and subsequent email correspondence revisions are highlighted in green
		recommendations provided only address enhanced monitoring of flux rates (e.g., continuous monitoring stations) without discussing whether changes to the current monitoring program are recommended. Please revise the RSE to discuss recommendations for the ongoing monitoring and mitigation of mercury and methylmercury in BCV.		

REDLINED figures found at end of EPA comment letter dated 12/6/24.










































## **ATTACHMENT 1**



Table 1.1 BCV cap and soil cover areas (New table based on EPA general comment 4)

Potential mercury source areas	Cap/soil cover details				
BYBY	BYBY was covered with a 2-ft layer of low-permeability clay and a 6-in. layer of vegetative cover				
Oil Landfarm and HCDA	HCDA was covered with a low-permeability RCRA-type cap in 1989; however, this was not a RCRA-regulated facility				
	The Oil Landfarm was covered with a multilayer low-permeability cap, consisting of a minimum of 2 ft of compacted clay, a flexible synthetic membrane liner, a geosynthetic drainage net, a geotextile filter fabric, a minimum of 1.5 ft of topsoil, cover vegetation, and cap drains. The final RCRA closure for the Oil Landfarm was certified by TDEC in December 1990				
S-3 Ponds Site	The S-3 Ponds Site was covered with a multilayer low-permeability engineered cap, covered with crushed stone, and paved with asphalt for use as a passenger vehicle parking lot. TDEC accepted the final RCRA closure certification for the S-3 Ponds Site on November 15, 1990				
Sanitary Landfill 1	The landfill was closed in 1985 by grading to promote drainage, capping with 2 ft of clay and topsoil, and establishing a vegetative cover				
Bear Creek Road Debris Burial	The BCV Phase I ROD concluded this area did not contain significant sources of contamination that could pose a risk to health or the environment; therefore, no action was warranted under CERCLA				
Creekside Debris Burial	The BCV Phase I ROD concluded this area did not contain significant sources of contamination that could pose a risk to health or the environment; therefore, no action was warranted under CERCLA				
SY-200 Yard	A compacted soil cover (3 to 5 ft) was installed over the site beginning in 1986				
Spoil Area 1	A 2-ft-minimum vegetative clay soil cover was installed over the site beginning in 1985				
Rust Spoil Area	A minimum of 1.5 ft of compacted clay and a 0.5-ft topsoil soil cover were installed over the site beginning in 1983				

BCV Phase I ROD = DOE/OR/01-1750&D4. Record of Decision for the Phase I Activities in Bear Creek Valley at the Oak Ridge Y-12 Plant, Oak Ridge, Tennessee, 2000, U.S. Department of Energy, Office of Environmental Management, Oak Ridge, TN.

BCV = Bear Creek Valley

BYBY = Boneyard/Burnyard

CERCLA = Comprehensive Environmental Response, Compensation, and Liability Act of 1980


HCDA = Hazardous Chemical Disposal Area

RCRA = Resource Conservation and Recovery Act of 1976

ROD = Record of Decision

SY = scrapyard

TDEC = Tennessee Department of Environment and Conservation






Figure 3.1. Total and dissolved mercury and methylmercury for the initial surface water sampling event, upstream to downstream". (Revised based on EPA specific comment 7)

Page 22 of 32





## **ATTACHMENT 2**



Long-term performance and baseline monitoring of groundwater and surface water is evaluated on an annual basis in the Remediation Effectiveness Report. Based on the annual evaluations, the following summary of mercury and methylmercury data is provided.

Bear Creek Valley mercury and methylmercury data available in the Oak Ridge Environmental Information System in surface water and selected groundwater monitoring locations for the monitoring period of January 1, 2009, through June 30, 2024, are provided in Table 1. The data summary includes the number of total mercury results for analyses with detection limits less than 51 ng/L, the number of detected results available for each location, and the average and maximum concentrations of detected total mercury and total methylmercury for the location. Sample locations are shown on Figure 1. In addition to sample locations, Figure 1 shows the area where the Maynardville Limestone subcrops beneath the soil layer. The Maynardville Limestone is known for its dissolutional weathering and well-developed karst. Interflows between the Bear Creek surface water channel and epikarst conduit flow pathways occur throughout the extent of the Maynardville Limestone beneath Bear Creek.

Data in Table 1 are arranged from upstream sample locations at the top of the table to downstream sample locations near Tennessee Highway 95 at the bottom of the table. It is noted there is a general absence of methylmercury data for groundwater monitoring wells, although there are methylmercury data for groundwater emerging from the springs. Historic research on methylmercury in the environment has largely focused on the aquatic food chain because of the toxicity of methylmercury via fish consumption by humans. Mercury and methylmercury may enter the food chain from sediment and water, and research on biogeochemical factors that enhance mercury methylation have focused on the roles of physicochemical conditions, habitat conditions, and microbial populations in aquatic settings where fish reproduce and live. There has been little research published on the methylation of mercury in the subsurface environment or in groundwater aquifer settings.

As shown in Table 1, groundwater mercury concentrations are highest in wells that monitor conditions in the S-3 Ponds plume area at the headwater of the Bear Creek hydrologic and Comprehensive Environmental Response, Compensation, and Liability Act of 1980 Administrative Watersheds. Surface water in the two headwater sample stations (NT-1 and BCK 12.34) exhibit lower mercury concentrations than are present in the S-3 Ponds vicinity groundwater.

As indicated in Table 1, a stream channel losing flow reach exists seasonally in Bear Creek between the Bear Creek Road crossing and the North Tributary-3 confluence. North Tributary-3 is the long-term surface water monitoring location for the now-remediated Boneyard/Burnyard. Prior to the Boneyard/Burnyard remedial action, there were elevated mercury concentrations in the North Tributary-3 surface water. The Boneyard/Burnyard remedial action included excavation of source areas in the Boneyard/Burnyard and contaminated floodplain soils and sediments. Completion of the remedial action included backfill of the excavated area with clean soil and construction of a sinuous stream channel with riparian vegetation planting. There are areas of marshy conditions adjacent to the restored stream channel that appear to be conducive to mercury methylation, as indicated by the relatively higher methylmercury concentration in the North Tributary-3 surface water. North Tributary-3 exhibits the highest average and the highest maximum measured methylmercury concentration in the Bear Creek monitoring network. At BCK 11.54/BCK 11.54A, a short distance downstream from the North Tributary-3 confluence to Bear Creek, the second highest average and maximum methylmercury concentrations are observed.

A short distance downstream of BCK 11.54/BCK 11.54A, the flow of Bear Creek sinks into the epikarst subsurface flow system. During significant periods of the year, the Bear Creek channel is dry from the sink



point downstream to the North Tributary-5 confluence. The North Tributary-5 total and methylmercury concentration data demonstrate relatively low inputs to Bear Creek.

Dye tracing conducted as part of the Bear Creek Valley remedial investigation demonstrated epikarst transport from this losing reach of Bear Creek emerges at a groundwater resurgence point at spring SS-4. The average and maximum total mercury concentrations at SS-4 and the maximum observed methylmercury concentration support the probable connection between the monitoring locations NT-3 and BCK 11.54/BCK 11.54A reach of higher concentrations.

Deep bedrock groundwater monitoring wells GW-704 (samples at 244.5 to 256 ft below ground surface) and GW-706 (samples at 156.1 to 182.5 ft below ground surface) lie between the Bear Creek sink point and the SS-4 spring and exhibit very low total mercury concentrations.

Downstream of the SS-4 spring discharge confluence, Bear Creek enters another significant losing flow reach that extends seasonally to the Bear Creek Road crossing adjacent to spring SS-5. Groundwater monitoring wells in this reach include wells GW-683 (sampling interval at 146 to 196.8 ft below ground surface) and GW-684 (sampling interval at 113.8 to 128.8 ft below ground surface). Total mercury concentrations in this well pair are higher than in the deeper wells, GW-704 and GW-706. Total mercury concentrations in wells GW-683 and GW-684 are similar to those measured at nearby spring SS-5. At spring SS-5, perennial flow exists that forms the head of continuous flow that extends downstream to Bear Creek kilometer 9.2 and beyond. Data from spring SS-6 may be somewhat misleading because during periods of high flow stage of Bear Creek, the spring area is sometimes inundated with main stem flow. Also, beaver ponding a short distance downstream of spring SS-6 caused inundation of the spring for a number of years and caused surface water backup into the epikarst of the area.

As previously noted, epikarst groundwater transport has been documented from the Bear Creek reach near BC-NT3 and BCK 11.54/BCK 11.54A and solids transport from Bear Creek to spring SS-4 has been observed.

Table 2 summarizes dissolved oxygen and redox potential measurements for most of the mercury and methylmercury monitoring locations. Surface water and spring discharges are well aerated, while groundwater from monitoring wells exhibits lower dissolved oxygen levels and wells GW-704 and GW-706 exhibit anoxic conditions consistent with their deep sample intervals. Electrochemically, most of the sample locations exhibit oxidizing conditions. Standard deviation of the average redox measurements is provided as an indicator of the relative variability of measurements at each location. Some, but not all of the groundwater monitoring well data exhibit electrochemically reducing conditions. Electrochemically reductive conditions in groundwater at the Oak Ridge Reservation are not uncommon.

Past mercury and methylmercury sampling events generally have not included field-filtered and unfiltered aliquots collected for analysis and have rarely had contemporaneous field turbidity or total suspended solids analyses performed. The primary sampling objectives to date have been the screening of total mercury concentrations for ambient water quality criteria (51 ng/L) comparisons, and ambient concentrations are usually much lower than the criterion, such that focus on partitioning of mercury and methylmercury between dissolved versus particle-associated was not a focus of investigation.

Although contemporaneous mercury and methylmercury data are not available for direct correlation with solids in the water samples, there are turbidity data from other sampling events during January 1, 2009, through June 30, 2024. Figure 2 is a graph that shows daily average flow rate versus turbidity data from



locations BC-NT3, SS-5, and BCK9.2. At locations SS-5 and BCK9.2, data are amenable to separating turbidity versus flow values into lower versus higher flow conditions; however, this is not the case for BC-NT3. Figure 3 includes box and whisker plots for turbidity at locations SS-5, SS-4 (where no flow data are available), BC-NT3, and BCK 9.2. Monitoring location BCK 11.54/BCK 11.54A has too few turbidity measurements to provide useful interpretive value. Comparison of the methylmercury values with the box plots in Figure 3 suggests turbidity/suspended solids may be an important factor in aqueous transport.

Components of turbidity may include fine mineral sediment and/or organic materials that could contain significant mercury and/or methylmercury.

Although groundwater from monitoring wells along the Bear Creek floodplain have not been analyzed for methylmercury, the methylmercury distribution and apparent connections from demonstrated primary surface area aquatic sources (North Tributary-3) through the epikarst transport systems provide a coherent conceptual model of the available data.



Table 1. Summary of BCV surface water and groundwater detected mercury and methylmercury data, 2009–2024

Station	Mercury (ng/L)			Meth	Methylmercury (ng/L)			
name	n/ndet	Average	Max	n/ndet	Average	Max	Medium	Comment
GW-276	9/9	101.74	440	-	-	-	GW	
GW-246	12/12	345.36	741	-	-	-	GW	
GW-815	2/2	310.50	410	-	-	-	GW	S-3 Area
NT-1	21/15	13.14	55.7	-	-	-	SW	S 3 med
BCK 12.34	43/43	14.68	127.2	32/32	0.11	0.258	BC SW - IP	
S07 (NT-2)	4/4	6.85	16.5	-	-	-	Trib SW	
		Bear	Creek los	ing reach	- sometimes	no flow		
BC-NT3	38/38	19.62	147	27/27	0.31	2.698	Trib SW	
BCK 11.54 / 11.54A	36/36	7.15	22.2	32/32	0.11	0.754	BC SW	BYBY residual
NT-4	60/2	84.50	108	-	-	-	Trib SW	EMWMF office area
EM-VWEIR	12/8	27.14	96	-	-	_	SW	EMWMF non- contact water
NT-5	7/7	2.64	6.1	7/7	0.10	0.14	Trib SW	Receives V-Weir flow
GW-706	19/19	0.99	1.94	-	-	-	GW	S-3 nitrate +
GW-704	19/19	0.43	1.82	-	-	-	GW	uranium
SS-4	52/32	15.61	87.7	28	0.10	0.39	Trib GW	Epikarst resurgence from BCK 11.54 sink
BC-NT8	53/53	3.88	10.7	31/30	0.11	0.266	Trib SW	BCBG discharge
GW-683	31/24	1.51	6.4	-	-	-	GW	S-3 nitrate +
GW-684	31/27	1.15	8	_	<u>-</u>	_	GW	uranium
GW-077	19/4	0.97	2.2	-	_	_	GW	
GW-078	19/18	0.60	2.1	-	-	-	GW	Zone 3 IP GW
SS-5	52/36	2.80	6.64	32/16	0.04	0.15	Trib GW	



Table 1. Summary of BCV surface water and groundwater detected mercury and methylmercury data, 2009–2024 (cont.)

Station Mercury (ng/I			L)	Methylmercury (ng/L)			Medium	Comment
SS-5 flow provides continuous baseflow in BC downstream to the IP								
BCK9.2	44/44	4.66	39.4	32/32	0.07	0.22	SW - IP	Zone 3 IP
BCK-7.87	15/15	3.39	7.45	-	-	-	BC SW	Zone 2/Zone 1
GW-712	19/2	1.64	3.13	-	-	-	GW	
GW-714	19/4	1.06	3.19	-	-	-	GW	
GW-713	31/8	5.02	38.7	-	-	-	GW	Zone 1 GW
SS-6	12/12	13.89	145	-	-	-	Trib GW	
SS-7	4/4	1.05	1.6	-	-	-	Trib GW	
BCK 4.6	30/30	4.62	61	30/26	0.11	0.419	BC SW	Near
BCK-4.55	15/15	2.92	7.6	-	-	-	BC SW	Highway 95
SS-8	8/8	2.79	9.8	-	-	-	Trib GW	

<sup>-</sup> Denotes no data available.

BC = Bear Creek

BCBG = Bear Creek Burial Grounds

BCK = Bear Creek kilometer

BCV = Bear Creek Valley

BYBY = Boneyard/Burnyard

EMWMF = Environmental Management Waste Management Facility

GW = groundwater

IP = integration point

max = maximum

n = number of analytical results with detection limits < 51 ng/L

n/det = number of detected results

NT = North Tributary

SS = surface spring

SW = surface water

Trib = Bear Creek tributary sample location






Figure 1. Locations of selected surface water and groundwater monitoring sites in BCV.



Table 2. Summary of BCV surface water and groundwater dissolved oxygen and redox, 2009-2024

Station name	Diss	Dissolved oxygen (ppm)			Redox (mV	Comment	
	n	Average	St dev	n	Average	St dev	<del>_</del>
GW-276	26	2.63	2.89	24	261	134	
GW-246	12	1.15	1.13	12	274	43	
GW-815	2	8.48	0.57	2	112	81	S-3 Area
NT-1	20	8.06	3.11	20	164	103	
BCK 12.34	217	7.74	2.08	189	165	83	
S07 (NT-2)	7	6.92	3.28	7	160	60	
BC-NT3	28	8.40	3.66	19	136	99	DANDAY '1 1
BCK 11.54	4	6.49	2.70	4	135	54	BYBY residual
NT-4	160	6.64	1.94	154	149	77	EMWMF office area
NT-5	7	7.91	2.31	7	179	50	Receives V-Weir flow
GW-706	31	0.31	0.38	29	112	90	S-3 nitrate + uranium
GW-704	32	0.61	0.34	30	59	134	uramum
SS-4	39	7.39	1.31	39	173	70	Epikarst resurgence
BC-NT8	41	7.98	2.40	39	146	96	BCBG discharge
GW-683	34	4.49	2.16	30	94	101	S-3 nitrate +
GW-684	31	2.48	2.74	29	91	102	uranium
GW-077	31	1.04	0.69	29	7	86	
GW-078	31	2.52	1.93	31	68	94	Zone 3 IP GW
SS-5	55	5.86	1.52	55	166	93	
BCK9.2	31	8.76	2.89	23	131	90	Zone 3 IP
BCK7.87	37	8.01	2.03	37	142	116	Zone 2/Zone 1
GW-712	31	1.48	3.84	29	-19	81	
GW-714	32	1.72	6.02	32	100	90	7 1 0 1
GW-713	31	2.31	4.32	29	-76	82	Zone 1 GW
SS-6	31	6.63	1.75	29	176	205	
SS-7	29	6.52	1.33	29	162	106	
BCK04.55	38	7.60	1.90	38	214	415	Near Highway 95
SS-8	29	7.07	1.91	29	197	170	

BC = Bear Creek

BCBG = Bear Creek Burial Grounds

BCK = Bear Creek kilometer

BCV = Bear Creek Valley BYBY = Boneyard/Burnyard

EMWMF = Environmental Management Waste Management

Facility

GW = groundwater

IP = integration point

n = number of dissolved oxygen results

NT = North Tributary ppm = parts per million

SS = surface spring

St dev = standard deviation






Figure 2. Daily average flow rate (Q) versus turbidity plots.




Figure 3. Box and whisker plots of turbidity data available for the data evaluation period 01/01/09-06/30/24.



Comment

Document Number:	Document Title:			
DOE/OR/01-2977&D1	Bear Creek Valley Mercury Sources Remedial Site Evaluation for the U.S. Department of Energy Oak Ridge Site Oak Ridge, Tennessee			
Name of Reviewer:	Organization:	Date Comments Transmitted:		
Randy C. Young	Tennessee Department of Environment and Conservation	November 22, 2024		

**General Comments** 

No.	Sect./Page	Comment	
1		The Mercury Management Approach section of the EMDF ROD states, "Unless the conclusion in the RSE accepted by all parties is for no further action, the RSE shall lead to other milestones for removal or remedial actions, including developing the substantive equivalent to developing load allocations and waste load allocations under 40 CFR 130.7(c)(2) and 130.2(g)(h) and (i)." TDEC does not believe this RSE meets the requirements of the Mercury Management Approach as outlined in the EMDF ROD and does not accept the conclusion that no further action is needed. Please explain how mercury methylation will be evaluated in accordance with the goal as defined by the EMDF ROD considering the following issues:	
		a. The goal of the RSE and defined in the EMDF ROD was to "evaluate mercury methylation in Bear Creek" which includes but is not limited to looking for sources of methylmercury. For example, the first sentence of Section 1.1 (p. 1-1) in the D1 RSE states the RSE objective is to evaluate potential sources of mercury and methylmercury in the BCV watershed in accordance with the EMDF ROD. However, the EMDF ROD (p. 2-64, Paragraph #4) states DOE shall conduct a RSE to evaluate mercury methylation in Bear Creek and conduct pilot or treatability studies as needed. The italicized words highlight an important difference between the objectives of this RSE and the EMDF ROD.	
		b. The U.S. Department of Energy's (DOE) response to TDEC Comment 2 on the Bear Creek Valley Mercury Sources Remedial Site Evaluation Sampling and Analysis Plan, Oak Ridge, Tennessee (DOE/OR/01-2958&D1) states an objective of the RSE is to obtain data from hydrologic settings that may contribute to mercury methylation. It also states, "parameters will be collected that may contribute to a better understanding of mercury methylation" The methylation objective is also included as a data quality objective (DQO) in Section 3.2 (p. 11). However, the report presents no information produced by the RSE regarding the mechanisms for mercury methylation.	
		c. If a conclusion of this RSE is that the concentration of mercury in fish tissue has decreased below the maximum allowable 0.3 $\mu$ g/g criterion, then at a minimum the environmental factors that have contributed to the decrease in methylmercury must be	

Clarification. In the recommendations section of the BCV mercury sources RSE, the text indicates the following: "In addition to obtaining discharge measurements and mass loading in Bear Creek, it is important to understand the biological factors controlling mercury methylation and trophic transfer in this creek. Although periphyton play an important role in mercury methylation in stream systems, how the periphyton community changes spatially and temporally in Bear Creek, and how these changes relate to methylmercury concentrations in the stream, are yet to be understood."

Response (02/04/2025)

It is recognized that additional investigation may be required to better understand the mercury methylation in Bear Creek. These investigations will continue through the mercury remediation technology development program; however, given the significant scope of understanding the mercury methylation, additional studies on mercury methylation in Bear Creek will be a long-term investigation. No change to the document proposed.

- a. Clarification. As stated in the DQO meeting held on June 29, 2023, and as described in Section 3.1 of the BCV mercury sources RSE, the overall problem to be addressed by the RSE is that there are insufficient data along Bear Creek and its tributaries to determine if there are potential sources of mercury and methylmercury in channel sediment and creek bank and floodplain soils that may be contributing to exceedances of fish tissue criterion in prior years. As such, the two main goals of the BCV mercury sources RSE (Section 3.2) were to:
  - Determine if there are areas (channel sediment and creek bank and floodplain soils) along Bear Creek and its tributaries that are potential sources of mercury and methylmercury that may affect fish.
  - Obtain data from various hydrologic settings (pools, beaver ponds, etc.) that may contribute to mercury methylation and its bioaccumulation in the environment of Bear Creek and a reference location (e.g., Hinds Creek).

The BCV mercury sources RSE fulfills the scope included in the project DQOs and adds significant data to the evolving conceptual model of mercury in the Bear Creek floodplain and aquatic system. The understanding of mercury methylation in Bear Creek is incomplete and will require an ongoing long-term investigation to inform development of the mercury bioaccumulation conceptual model for Bear Creek.

No change to the document proposed.

**b.** Clarification. DOE conducted sampling and analyses as outlined in the BCV mercury sources RSE SAP. The BCV mercury

Clarification. The last paragraph of Section 5.1 (Conclusions) was revised as follows:

"Overall, mercury contamination of surface water, floodplain and creek bank soil, and channel sediment is highest in the upper reaches of Bear Creek, with decreasing concentrations downstream. The increasing mercury flux in the downstream flow direction is dominated by the increasing flow volume, as concentration data tend to gradually decrease. This characteristic suggests mercury entering Bear Creek downstream of the known source associated with the S-3 Ponds area is derived from dispersed mercury that may be associated with secondary contamination of creek bank and floodplain soil. Although mercury is detected at low levels in sampled media, concentrations are much lower than other ORR mercury-contaminated sites and fish concentrations continue to decrease. This BCV mercury sources RSE did not identify a source of mercury in the media collected in and along Bear Creek that indicates active remediation or an RI of mercury sources is needed. However, the presence of mercury in fish tissue indicates additional studies to understand methylation in Bear Creek are necessary to further refine the CSM."

Revised Response (04/10/2025) - Changes highlighted in yellow are based on the 04/22/2025

in areen

comment response meeting and subsequent email correspondence revisions are highlighted

In addition, Section 5.2 (Recommendations) was revised as follows:

"While the historical water quality and discharge measurement dataset is extensive in Bear Creek, there are far fewer coordinated water-quality-plus-discharge-measurement campaigns with a specific emphasis on total versus dissolved mercury and methylmercury. Although the baseflow analysis of total versus dissolved mercury and methylmercury conducted in this investigation showed approximately 73% of the mercury and 39% of the methylmercury were particle-associated, there are no event-based (i.e., precipitation-driven flows), coordinated, discharge and water quality sampling campaigns. To assess total and dissolved fluxes of mercury and other solutes in addition to those available at established stations in Zone 3, the following additional coordinated flow measurement and sampling locations are needed:

- A BCK 1 monitoring station equipped with a pressure transducer and multiparameter sonde (temperature, pH, specific conductance, dissolved oxygen, and nitrate) was installed in FY 2024. These continuous monitoring instruments may be supplemented with manual or programmed auto-sampling for mercury and methylmercury or other constituents. These measurements can provide flux data at one location.
- Additional gaging stations would be required to fully assess loading at additional locations in Bear Creek. Dedicated, coupled, concentration-discharge measurements under both baseflow conditions and over several flood hydrographs would be ideal to better understand mass loading. Several flow-control structures at road crossings exist along Bear Creek downstream of existing location BCK 9.2. Evaluating potential instrumentation of selected existing structures may identify opportunities for relatively cost-effective flow measurement and sampling stations for additional flux-measurement monitoring.

In addition to obtaining better discharge measurements in Bear Creek, understanding the biological factors controlling mercury methylation and trophic transfer in this creek is important. In the BCV Watershed, where aqueous mercury concentrations are low but fish tissue concentrations are relatively elevated, the path to achieving and maintaining fish tissue guidelines must include more than a simple assessment of aqueous total mercury concentrations. Mercury, especially in the methylmercury form, biomagnifies through the food web, leading to elevated concentrations in fish. The drivers for mercury methylation in the creek, as well as for mercury bioaccumulation, need to be understood. Because periphyton are likely the substrate that enables mercury methylation and are

Date: 06/24/2025 eIRO#: 8183





Comment No. Sect./Page	Comment	Response (02/04/2025)	Revised Response (04/10/2025) – Changes highlighted in yellow are based on the 04/22/2025 comment response meeting and subsequent email correspondence revisions are highlighted in green
	evaluated to verify the levels of methylmercury do not cause future exceedances. Identifying the mechanisms of mercury methylation should then precede an evaluation of potential technologies that can be used to either disrupt or lessen the effectiveness of the methylation in the creek and ensure long-term protectiveness. For example, given DOE's findings that mercury methylation potential varies by organic matter type, evaluating the potential impacts on methylation from different organic matter types may help inform future remedial actions.  d. Based on the RSE's lack of information about methylation, it is not yet clear what additional actions might be warranted, nor is it clear how to develop the substantive equivalent of developing load allocations and waste load allocations.	sources RSE contains and presents various data on water, sediment, and soil chemistry relevant to environmental conditions that may promote mercury methylation. Data are also displayed in the context of the limnologic setting from which samples were obtained. The OREM investigation of mercury distribution in the Bear Creek channel sediment, surface water, and creek bank and floodplain soils provides a spatial snapshot that may be useful to support future, more detailed investigations of specific environmental "compartments" (bed sediment, bank soil, etc.) of interest. No change to the document proposed.  C. Clarification. There is extensive literature in the global scientific community that describes the factors that can affect mercury methylation, calling mercury methylation "a bio-physico-chemical conundrum in which the efficiency of biological HgII methylation appears to depend chiefly on HgII and nutrients availability, the abundance of electron acceptors such as sulfate or iron, the abundance and composition of organic matter as well as the activity and structure of the microbial community." As mentioned in the BCV mercury sources RSE, ORNL's Environmental Sciences Division compiled data to construct a conceptual model for mercury transport and transformation (i.e., methylation) (ORNL/TM-2023/3069) and concluded there were significant data gaps that preclude the ability to identify methylation mechanisms. A key recommendation from this study was to increase knowledge of the microbial community within the creek, as well as water quality parameters with enough spatio-temporal frequency to be able to make correlations and predictions. These data are currently being collected as part of ORNL's mercury remediation technology development program, and the findings will be reported as they are available. No change to the document proposed.  d. Clarification. See initial response above.	also the base of the food web, serving to concentrate methylmercury concentrations in the dietary pathways leading to fish, future monitoring should focus on periphyton dynamics. Although periphyton are understood to play an important role in mercury methylation in stream systems, how the periphyton community changes spatially and temporally in Bear Creek and how these changes relate to methylmercury concentrations in the stream are yet to be understood. Recommendations to evaluate periphyton in Bear Creek in support of the CSM under the mercury remediation technology development program follow:  • Based on guidance provided by Bravo et al. (2017), evaluating organic material characteristics that may affect mercury methylation and biouptake is required to understand the mixes of organic compound origins and molecular weights and to inform attempts to discern potential aquatic habitat characteristics that influence mercury methylation, accumulation in media, and bioaccumulation. This information is critical to any best management watershed-scale practices aimed at reducing methylmercury concentrations in fish.  • To gain a better understanding of mercury methylation for decision-making, data from the aforementioned continuous monitoring instruments should also be supplemented with manual sampling data for measures of periphyton abundance and community structure.  • To understand spatial and temporal periphyton dynamics over time, deploying nutrient (e.g., the nitrate sensor at the BCK1 monitoring station), light, and turbidity sensors to augment the discharge and water quality measurements collected at additional future monitoring station installations is required. The continuous measurement of periphyton- and mercury-relevant water quality perameters may also be enriched by installing relatively low-cost field cameras capable of estimating aquatic vegetation distribution and abundance in proximity to the monitoring station on a daily basis. Nutrients and light are both important drivers of periphyton distributi





Comment No.	Sect./Page	Comment	Response (02/04/2025)	Revised Response (04/10/2025) – Changes highlighted in yellow are based on the 04/22/2025 comment response meeting and subsequent email correspondence revisions are highlighted in green
2		Please include a discussion of efforts conducted or planned for identifying mechanisms of mercury methylation and/or ways in which the methylation of mercury in Bear Creek could be reduced.	Clarification: The understanding of mercury methylation in Bear Creek is incomplete and will require ongoing long-term investigations to inform development of the mercury bioaccumulation conceptual model and hydrologic conceptual model for Bear Creek. ORNL's mercury remediation technology development program was originally developed to investigate watershed-scale approaches to reducing mercury and methylmercury concentrations in EFPC but has increasingly focused on understanding the differences in mercury methylation between EFPC and Bear Creek, while evaluating best management practices or management actions that could reduce methylmercury concentrations in both creeks. These efforts are summarized in annual reports and publications (e.g., ORNL/SPR-2023/3178, ORNL/TM-2021/2207). Rather than reproducing these efforts, the BCV mercury sources RSE focused on key data gaps that are not addressed by the mercury remediation technology development program (i.e., sources of mercury and methylmercury to Bear Creek). No change to the document proposed.  ORNL/SPR-2023/3178. Mercury Remediation Technology Development for Lower East Fork Poplar Creek FY 2023 Update, 2023, Oak Ridge National Laboratory, Oak Ridge, TN.  ORNL/TM-2021/2207. Mercury Remediation Technology Development for Lower East Fork Poplar Creek - FY 2021 Update, 2021, Oak Ridge National Laboratory, Oak Ridge, TN.	Clarification. The BCV mercury sources RSE focused on key data gaps that are not addressed by the mercury remediation technology development program. However, some additional detail of these long-term investigations that are conducted by ESD were added to Section 1.2.1.2 as follows:  "A 2022 data compilation report for mercury in Bear Creek (ORNL/TM-2023/3069) summarized data from compliance and investigatory studies to begin building a conceptual model to understand the processes affecting mercury transport and transformation in the Bear Creek Watershed and to highlight key knowledge gaps in understanding these processes. This report summarizes historical data from compliance and investigatory studies relevant to mercury contamination and transformation in the Bear Creek Watershed and highlights the relative paucity of water quality data in Bear Creek with respect to mercury concentrations and transformation processes. A recommendation from this study was to install a long-term monitoring station to provide needed data to establish functional linkages among identified components of a conceptual model of mercury transport with measured mercury values from the watershed.  The mercury remediation technology development program at ORNL, which began in 2014, originally focused on understanding and addressing mercury concentrations, flux, and bioaccumulation in EFPC. Research from this program highlighted the importance of periphyton in mercury transformation in stream systems, and in 2021, a task specifically focused on periphyton dynamics and distribution was added to the study plan. Because the aqueous methylmercury concentrations in EFPC are similar to those in nearby Bear Creek despite significantly higher aqueous total mercury concentrations in EFPC, comparing mercury methylation and demethylation processes between the two streams was of particular interest, and Bear Creek became (and remains) a point of focus in the mercury remediation technology development program (ORNL/SPR-2023/3178).  Future work will bui
3		According to the U.S. Environmental Agency (EPA) guidance, to properly evaluate trends in bioaccumulation and make decisions regarding risks associated with fish consumption, the size of the fish sampled should be both consumable and consistent with past data collection efforts. DOE should not use data from fish that are abnormally small for decision making, unless additional effort is made to account for the small size such as allowing proper time for population recovery and/or modeling of bioaccumulation. In 2021 when the mercury strategy was developed and placed in the EMDF ROD, the mercury levels in fish tissue were trending steeply downward and projected to continue downward. Instead, the data has been variable as shown in Figure 4.10:	Clarification. EPA guidance for fish sampling (https://www.epa.gov/sites/default/files/2015-06/documents/volume1.pdf) allows for flexibility in terms of species and size classes sampled, depending on availability. The goal of the sampling is to assess potential exposure from eating resident fish, and while every effort was made to sample comparably sized fish over the years, natural changes (e.g., changes in beaver activity and impoundments) as well as anthropogenic actions (e.g., overfishing of fish populations to achieve tissue mass for EMDF radionuclide monitoring) have led to changes in the species of fish available for sampling as well as the size structure of the fish populations. For quality assurance purposes, non-lethal sampling is not possible with the tissue requirements for the COCs and CERCLA requirements. No change to the document proposed.	



Comment No.	Sect./Page	Comment	Response (02/04/2025)	Revised Response (04/10/2025) – Changes highlighted in yellow are based on the 04/22/2025 comment response meeting and subsequent email correspondence revisions are highlighted in green		
		BCK 3.3 Rock bass BCK 9.9 Redbreast BCK 9.9 Rock bass HCK 20.6 Rock bass HCK 20.6 Rock bass HCK 20.6 Rock bass BCK 9.9 Rock bass HCK 20.6 Rock bass HCK 20.6 Rock bass BCK 9.9 Rock bass HCK 20.6 Rock bass HCK 20.6 Rock bass BCK 9.9 Rock bass HCK 20.6 Rock bass HCK 20.6 Rock bass BCK 9.9 Rock bass HCK 20.6 Rock bass HCK 20.6 Rock bass HCK 20.6 Rock bass BCK 9.9 Redbreast HCK 20.6 Rock bass BCK 9.9 Redbreast BCK 20.6 Rock bass HCK 20.6 Rock bass HCK 20.6 Rock bass HCK 20.6 Rock bass HCK 20.6 Rock bass BCK 9.9 Redbreast BCK 9.9 Red				
4		Given both that the last paragraph of <b>Section 5.1</b> of the RSE states that mercury contamination is "widespread", and the mercury speciation studies indicate organically bound mercury dominates much of the Bear Creek Valley watershed, especially in soils, addressing areas with low-level mercury sources may be necessary to effect a change on fish tissue in the watershed. For example, even though Bear Creek mercury is orders of magnitude lower than the remediation goal of 400ppm for Lower East Fork Poplar Creek (LEFPC) (as referenced for comparison in presentations made at multiple project team meetings), LEFPC is characterized by mercury fractions that are less readily available to absorb and bioaccumulate than Bear Creek. These differences suggest that actions targeting lower-level sources may be necessary to address methylmercury cycling in the food chain and restore Bear Creek to meet the recreational use designation, per the goal stated in the EMDF ROD Mercury Management Approach.	Clarification. The text in Section 5.1 was revised as follows to better describe the mercury detected in media associated with the BCV mercury sources RSE:  "Although mercury is detected at low levels in sampled media, concentrations are much lower than other ORR mercury-contaminated sites and fish concentrations in Bear Creek continue to decrease. This BCV mercury sources RSE did not identify a source of mercury in the media collected along Bear Creek that significantly contributes to the Bear Creek mercury contamination or that would warrant active remediation."	Clarification. See revised general comment response 1.		
5		Data presented in this RSE report (including site description in Section 1.2.1) support the hypothesis that shallow groundwater/known contaminant plumes in the subsurface are potential sources of mercury	Clarification. Long-term performance and baseline monitoring of groundwater and surface water is evaluated on an annual basis in the RER. Known residual mercury in the vicinity of the S-3 Ponds plume			





Comment No.	Sect./Page	Comment	Response (02/04/2025)	Revised Response (04/10/2025) – Changes highlighted in yellow are based on the 04/22/2025 comment response meeting and subsequent email correspondence revisions are highlighted in green
		and/or methylation impacting Bear Creek that should be investigated. TDEC requested that shallow groundwater be considered as a source of mercury, methylmercury, or mercury methylation in DQO meetings and in written comments on the related Mercury RSE Sampling and Analysis Plan (SAP) (Appendix A). While typical hyporheic zone/pushpoint sampling may not be appropriate given conditions in Bear Creek (as discussed in the Bear Creek Valley (BCV) Mercury Sources RSE SAP response to comments), TDEC is again recommending that shallow groundwater be investigated as a potential source that could be addressed to positively impact Bear Creek water quality.	may contribute to mercury detections in Bear Creek (see Attachment 1 at the end of these comment responses).  Note that response actions for the S-3 Ponds and for BCV groundwater have yet to be implemented and will be included in the future design considerations for Pathway 3 under the BCV Phase I ROD or in the final groundwater decision for BCV.  No change to the document proposed.	in green
6		TDEC recommends additional cooperation with Oak Ridge National Laboratory's Environmental Sciences Division to evaluate innovative strategies to mitigate Hg impacts and support the restoration of Bear Creek.	Clarification. ORNL ESD is involved in the process of evaluating innovative strategies to mitigate mercury impacts of Bear Creek (see responses to comments 1c and 2); however, these investigations are not in the scope of the BCV mercury sources RSE. No change to the document proposed.	Clarification. See revised general comment response 2.
7		DOE should not use the conclusions of this RSE as a basis for decision-making, and there is not a consensus on future actions related to the water treatment and discharge at EMDF. One sampling event that is atypical due to small fish sizes is not a trend, and the variability of concentrations dictates that a longer period of observation is necessary. The data needed to support the removal of Bear Creek from the 303(d) list would have to be much more than what has been presented in this RSE. In order to maintain compliance with the fish tissue standard, if achieved, the RSE would also need to explain the mechanisms of methylation.	Clarification. See the responses to general comment 1 and specific comment 10. No change to the document proposed.	Clarification. See revised general comment response 1.
	<u> </u>		Specific Comments	
1	Page 1-7, Section 1.2.1.2, second paragraph	Revise the wording in the final sentence of this paragraph that characterizes mercury concentrations in fish tissue as occasionally exceeding the ambient water quality criteria (AWQC). Multiple types of fish have exhibited methylmercury levels in tissue regularly exceeding the AWQC level for the past couple decades.	Clarification. The text in Section 1.2.1.2 was revised as follows:  "but mercury concentrations in fish collected in Bear Creek are relatively high, occasionally exceeding the EPA-recommended ambient water quality criterion (AWQC) for mercury (0.3 µg/g in fish) until recent years (ORNL/TM-2023/3069)."	
2	Page 1-8, Section 1.2.2, last paragraph	This section describes the issues with mercury at the Storage Yard-200 (SY-200) area and states, "free mercury was observed in some of the borings" This text does not align with the conclusions of the report that the RSE did not identify a source of mercury warranting active remediation. Please add an explanation for why the report did not identify the SY-200 area as a potential source of mercury to Bear Creek.	Clarification. The BCV mercury sources RSE SAP identified the SY-200 Yard as a potential source of mercury in BCV, as also described in Section 1.2.2 of the BCV mercury sources RSE. Based on this potential source of mercury to Bear Creek, transect locations were selected along Bear Creek in the vicinity of the SY-200 Yard (transects BCT14 and BCT15). Prior to the BCV OU 2 ROD that included the SY-200 Yard, a soil cover of 3 to 5 ft was placed at SY-200. The BCV OU 2 ROD did not require implementation of additional RAs for SY-200. The selected remedy from the BCV OU 2 ROD relies on maintaining the soil cover and LUCs for SY-200.	





Comment No.	Sect./Page	Comment	Response (02/04/2025)	Revised Response (04/10/2025) – Changes highlighted in yellow are based on the 04/22/2025 comment response meeting and subsequent email correspondence revisions are highlighted in green
			No change to the document proposed.	
3	Page 2-7, Section 2.1.3,	Please include additional discussion on the width of the floodplain around Bear Creek, the lateral extent of floodplain samples, and whether any biased samples were collected from identified depositional areas. Please explain why the floodplain soils further from the creek were excluded from consideration as a potential source of mercury to surface water.	Clarification. Transect locations were selected during the reconnaissance survey based on the location of potential source areas and pool areas in Bear Creek and the mouth of tributaries. Considerations on the selection of the transect locations included potential depositional areas. Figures G.1 through G.5, which are included in Attachment 2 at the end of these comment responses, show Bear Creek and the 100-year floodplain. In the upstream areas of Bear Creek (Zone 3) (Figure G.5), there is very little floodplain identified due to steep banks, and floodplain soil samples were generally collected within 5 ft of Bear Creek. In the downstream areas where the floodplain is slightly wider, soil samples were collected within 5 to 10 ft on each side of Bear Creek, but within the extent of any observed floodplain terrace.	
			Also included in Attachment 2 at the end of these comment responses is Table G.1, which provides observations made during the field sampling event. Figures G.1 through G.5 and Table G.1 are included as new Appendix G to the D2 version of the BCV mercury sources RSE.	
4	Page 2-7, Section 2.1.3, second paragraph	The first sentence states soils samples were collected from "either" side of Bear Creek, should this say that samples were collected from "each" side of Bear Creek? Please provide additional details regarding sample volumes and how samples were collected and composited.	Clarification. The sentence was revised as follows:  "Floodplain soil was collected from the upper 0.5 ft on each side of Bear Creek to generate a single composite sample representing both sides of the floodplain (Figure 2.2)."  As described in the WRRP QAPP, soil and sediment sampling are conducted per UCOR procedures (PROC-ES-2300, Soil Sampling, and PROC-ES-2302, Sediment Sampling, respectively). Sample compositing is per UCOR procedure PROC-ES-2009, Sample Compositing and Homogenization, which outlines collecting equal proportions of solid material and mixing to achieve a homogeneous composite sample. As described in the BCV mercury sources RSE SAP, the volume collected was based on the volume required to fill the sample jars.	
5	Section 3.3, Table 3.9 and Figures 3.11 through 3.22	Several locations where sequential extraction results are presented in Table 3.9 show a cumulative percentage significantly greater than 100%. For example, floodplain soil collected from the Hind's Creek Transect Reference (HCTREF) location has a cumulative percent that adds up to 128%. The graphs in this section normalize the data from each location to 100%. This makes it seem as though the HCREF location had significantly less organically bound mercury (F3) when in fact the percent reported in the table for F3 at HCTREF was 62.1%, which is within the range of F3 percentages measured at the BCV transects. Please verify the data reported is accurate and, if so, please explain what errors in the mercury fraction calculations could result in a sum that exceeds the total amount of mercury.	Clarification. Table 3.9 was revised to be consistent with the method of summation us by the laboratory. The laboratory does not include in the summation of fractions results that were deemed non-detectable mercury. Table 3.9 contains the laboratory-reported result at the non-detect value; however, the percentage contribution to the summed total has been set to 0.0% for non-detected results. There remain some slight differences in the laboratory-reported mercury sums and the manual sums because of significant digit convention used by the laboratory (two significant digits in the sums with exclusion of non-detected fractions).  As a result of this change to use the laboratory reporting convention, Figures 3.11 through 3.22 were revised to remove non-detected fractions from the stacked bar graphs. In those instances, the validation flag "UJ" data label is associated with boundaries between detected fractions.  Revised Table 3.9 and Figures 3.11 through 3.22 are included in Attachment 3 at the end of these comment responses.	





Comment No.	Sect./Page	Comment	Response (02/04/2025)	Revised Response (04/10/2025) – Changes highlighted in yellow are based on the 04/22/2025 comment response meeting and subsequent email correspondence revisions are highlighted
6 6	Section 4.1	Spring SS-5 is identified as a source of significant Hg flux, data in Oak Ridge Environmental Information System indicate that Spring SS-4 exhibits flux of Hg and MeHg comparable to SS-5, and Spring SS-6 also sees a steady flux of Hg. Given the multiple groundwater/surface water connections in Bear Creek, these data suggest that groundwater may be a source of mercury and methylmercury into Bear Creek. Please explain how the Hg fluxes from groundwater along Bear Creek will be evaluated.	Clarification. DOE agrees the extensive interflow between Bear Creek and the underlying epikarst of the Maynardville Limestone is a potential groundwater transport pathway for surface water contaminants. DOE does not agree with the assessment of significant methylmercury formation in the groundwater system presented in the comment. DOE regards flux to be a measure of mass per unit of time (e.g., mg/day). Flow data are not available to calculate mercury flux contributions from the SS-4 spring. A graph included in Attachment 3 at the end of these comment responses shows the actual computed methylmercury flux for the BCK 11.54/BCK 11.54A and SS-5 spring monitoring locations for the period 01/01/09–06/30/24. The calculated methylmercury flux results from upstream at BCK 11.54/BCK 11.54A are also shown for comparison.	in green
			See also the response to TDEC general comment 5. No change to the document proposed.	
7	Page 4-2, Section 4.1,	The text here states "higher flow rates cause increased sediment transport, and with the strong particle retention of mercury, its mass transport can be greatly increased during high flow events". Provided the recognition that high flow events can greatly increase the short-term particle transport and mercury discharge to the stream, please explain how the current sampling strategy is sufficient when the time elapsed since last significant rain event prior to sampling has averaged from between 3 and 24 days as presented in <u>Table 1</u> of the <i>Bear Creek Special Studies Report 2021</i> .	Clarification. The WRRP Bear Creek monitoring results described in Section 4.1 are implemented per the BCV RAR CMP and are based on the BCV Phase I ROD goals of measuring uranium concentrations and flux rates at specified locations and are provided for informational purposes. The monitoring outlined in the BCV RAR CMP fulfills the BCV Phase I ROD performance monitoring requirements. No change to the document proposed.	
8	Page 4-7, Section 4.1, fourth paragraph	The text here emphasizes the lack of dissolved mercury in the surface water "an average of 27% (+/-5%) of mercury in surface water was dissolved, with the remaining 73% being associated with filterable solids". This dominance of particle-bound mercury indicates that the majority of mercury in the system is available to filter feeding organisms in the food web supporting the trophic transfer of mercury to larger organisms and the cycling of methylmercury in the system. Please provide an explanation for how the cycling and transfer of methylmercury was or will be evaluated.		
			The following sentence in Section 5.1 was revised to include this reference:	
			"Despite low aqueous mercury concentrations in Bear Creek, the fish fillet concentrations are relatively high, likely due to high methylation efficiency and efficient trophic transfer (ORNL/TM-2024/3735)."	
			Note that this reference was also added to Chapter 6:	
			"ORNL/TM-2024/3735. Mercury trophic transfer in Bear Creek, 2024, Oak Ridge National Laboratory, Oak Ridge, TN."	
			* Additional references provided in this comment response:	





Comment No.	Sect./Page	Comment	Response (02/04/2025)	Revised Response (04/10/2025) – Changes highlighted in yellow are based on the 04/22/2025 comment response meeting and subsequent email correspondence revisions are highlighted in green		
			ORNL/TM-2011/108. Assessment of Contaminant Bioaccumulation in Invertebrates and Fish in Waters on and Adjacent to the Oak Ridge Reservation-2010, 2016, Oak Ridge National Laboratory, Oak Ridge, TN.			
			ORNL/TM-2016/247. Assessment of Contaminant Bioaccumulation in Aquatic Biota on and Adjacent to the Oak Ridge Reservation-2015, 2016, Oak Ridge National Laboratory, Oak Ridge, TN.			
			ORNL/TM-2021/1891. Assessment of Contaminant Bioaccumulation in Aquatic Biota on and Adjacent to the Oak Ridge Reservation-2020, 2021, Oak Ridge National Laboratory, Oak Ridge, TN.			
9	Page 4-7 Section 4.1,	The text in multiple areas on this page relates mercury in Bear Creek to the S-3 Ponds including the last paragraph of the section that refers to "the known source associated with the S-3 Ponds area." Please provide an explanation for how the conclusion of the report was that no sources of mercury were identified in the context of a known source in the S-3 Ponds Area.	Clarification. The BCV mercury sources RSE investigated soil and sediment along Bear Creek to investigate if there is a source of mercury to fish in the vicinity of Bear Creek. No significant source was identified in the media in the vicinity of Bear Creek. No change to the document proposed.			
10	Section 4.2, Section 5.1, Figure 4.10	It is stated here "fish fillet concentrations were approaching and dropping below the EPA-recommended fish-based AWQC of 0.3 µg/g". As per the guidance document EPA-823-R-01-001 referenced in the EMDF ROD, this EPA recommended value of 0.3 µg/g is a not-to-exceed criterion and is therefore inappropriate to be used to compare to average fish tissue concentrations. While recent fish tissue data from Bear Creek indicate tissue concentrations of mercury/methylmercury have decreased and are hovering around the EPA criterion on average, these data sets continue to include samples with concentrations above the EPA tissue criterion. Also, as indicated in this RSE, populations of rock bass and redbreast sunfish were significantly impacted/overharvested as a result of the intensive sampling effort in 2021 and are not representative of normal conditions. Therefore, these fish tissue data are likely not appropriate for supporting delisting of Bear Creek from the CWA 303(d) list, per the EMDF ROD Mercury Management Approach. Please explain how the size of the fish will be normalized to allow for accurate comparisons to previous sampling events and trends. Please include details on maximum fish tissue concentrations and how many fish tissue samples exceeded the AWQC criterion. Please add distribution boxes to Figure 4.10 similar to Figure 2.19 in the 2024 Remediation Effectiveness Report (DOE/OR/01-2960).	Clarification. The guidance Water Quality Criterion for the Protection of Human Health: Methylmercury, EPA-823-R-01-001, January 2001, has a statement of "The TRC of 0.3 mg methylmercury/kg fish is the concentration in fish tissue that should not be exceeded based on a total consumption of 0.0175 kg fish/day." Implementation guidance for the criterion Guidance for Implementing the January 2001 Methylmercury Water Quality Criterion, EPA-823-R-10-001, April 2010, in Section 4.3 uses an average concentration in fish tissue to compare against the mercury fish tissue WQC. Additionally, ingestion of a contaminant occurs over an exposure time period (duration) and is not represented by a single value.  EPA guidance for fish sampling (https://www.epa.gov/sites/default/files/2015-06/documents/volume1.pdf) allows for flexibility in terms of species and size classes sampled, depending on availability. The goal of the sampling is to assess potential exposure from eating resident fish, and while every effort was made to sample comparably sized fish over the years, natural changes (e.g., changes in beaver activity and impoundments) as well as anthropogenic actions (e.g., overfishing of fish populations to achieve tissue mass for EMDF radionuclide monitoring) have led to changes in the species of fish available for sampling as well as the size structure of the fish populations. For quality assurance purposes, non-lethal sampling is not possible with the tissue requirements for the COCs and CERCLA requirements.  No change to the document proposed.			
11	Page 3-41, Section 3.4, Table 3.10	It is noted there is a distinct difference in partitioning of the forms of mercury in the sequential extraction data between Bear Creek and Lower East Fork Poplar Creek, and that this partitioning may be a key factor in the unsolved mystery of why the conversion of	Clarification. While it is true that the proportion of aqueous methylmercury to total mercury concentrations is higher in Bear Creek than in EFPC, the statement that the "conversion of mercury into methylmercury is much more efficient" is premature. Methylmercury concentrations are the net result of methylation and			





Comment No.	Sect./Page	Comment	Response (02/04/2025)	Revised Response (04/10/2025) – Changes highlighted in yellow are based on the 04/22/2025 comment response meeting and subsequent email correspondence revisions are highlighted in green		
		mercury into methylmercury is much more efficient in the Bear Creek watershed than in East Fork Poplar Creek. This should form the basis of future research opportunities.	demethylation processes, the mechanisms of which are not understood in Bear Creek. Please see the responses to EPA general comments 1, 1c, and 2 for more information on the efforts to understand mercury methylation and speciation in EFPC and Bear Creek. No change to the document proposed.			
12	Section 3.2.2, page 3-13: Table 3.7 page 3-19	Given the earlier general comments and the difference noted in specific comment #11, DOE should consider the idea of removal of the relatively high concentration creek bank soils that have been identified in the general vicinity upstream:  "Mercury concentrations in the sampled BCV soils and channel sediment are comparatively low. (Note the laboratory reported mercury and methylmercury concentrations in soil and sediment in units of µg/kg, equivalent to parts per billion [ppb]). The maximum measured mercury concentration in floodplain soil was 3500 µg/kg at BCT12A, upstream of the NT-3 confluence with Bear Creek. The maximum measured mercury concentration in creek bank soil was 7100 µg/kg at BCT13, located near the former HCDA entrance. The maximum measured mercury concentration in channel sediment was 530 µg/kg at BCK12A, upstream of the confluence of NT-3 with Bear Creek and downstream of the HCDA."  Because of the unique nature of the mercury partitioning in Bear Creek and the efficiency of methylation in the watershed, DOE should consider that the levels for triggering removal of source material may require a lower threshold than in East	Comment noted. For comparison purposes, mercury soil concentrations detected in the BCV mercury sources RSE (maximum mercury concentration of 7.1 mg/kg) were orders of magnitude below the soil cleanup levels established in the LEFPC ROD (400 mg/kg) and the UEPFC Phase II ROD (325 mg/kg). Surface water concentrations for mercury were likewise around an order of magnitude less than the 51-ng/L AWQC. No change to the document proposed.			
13	Section 5.1, page 5.2	Fork Poplar Creek.  The last sentence of this section concludes there is not a source of mercury that would warrant active remediation, but as identified in comments above, multiple statements throughout the report seem contradictory to this conclusion including known sources near the upper reaches of the creek and statements that mercury is being derived from the "widespread" contamination across the creek bank and floodplain soils. Please provide additional supporting detail on how the conclusion of the RSE is for no remedial action, and explain what further plans DOE has to study mercury flux to the creek, investigate the methylation occurring the creek ecosystem, and evaluate potential remedial actions for source removal, stabilization, etc.				





# **ATTACHMENT 1**



Long-term performance and baseline monitoring of groundwater and surface water is evaluated on an Long-term performance and baseline monitoring of groundwater and surface water is evaluated on an annual basis in the Remediation Effectiveness Report. Based on the annual evaluations, the following summary of mercury and methylmercury data is provided.

Bear Creek Valley mercury and methylmercury data available in the Oak Ridge Environmental Information System in surface water and selected groundwater monitoring locations for the monitoring period of January 1, 2009, through June 30, 2024, are provided in Table 1. The data summary includes the number of total mercury results for analyses with detection limits less than 51 ng/L, the number of detected results available for each location, and the average and maximum concentrations of detected total mercury and total methylmercury for the location. Sample locations are shown on Figure 1. In addition to sample locations, Figure 1 shows the area where the Maynardville Limestone subcrops beneath the soil layer. The Maynardville Limestone is known for its dissolutional weathering and well-developed karst. Interflows between the Bear Creek surface water channel and epikarst conduit flow pathways occur throughout the extent of the Maynardville Limestone beneath Bear Creek.

Data in Table 1 are arranged from upstream sample locations at the top of the table to downstream sample locations near Tennessee Highway 95 at the bottom of the table. It is noted there is a general absence of methylmercury data for groundwater monitoring wells, although there are methylmercury data for groundwater emerging from the springs. Historic research on methylmercury in the environment has largely focused on the aquatic food chain because of the toxicity of methylmercury via fish consumption by humans. Mercury and methylmercury may enter the food chain from sediment and water, and research on biogeochemical factors that enhance mercury methylation have focused on the roles of physicochemical conditions, habitat conditions, and microbial populations in aquatic settings where fish reproduce and live. There has been little research published on the methylation of mercury in the subsurface environment or in groundwater aquifer settings.

As shown in Table 1, groundwater mercury concentrations are highest in wells that monitor conditions in the S-3 Ponds plume area at the headwater of the Bear Creek hydrologic and Comprehensive Environmental Response, Compensation, and Liability Act of 1980 Administrative Watersheds. Surface water in the two headwater sample stations (NT-1 and BCK 12.34) exhibit lower mercury concentrations than are present in the S-3 Ponds vicinity groundwater.

As indicated in Table 1, a stream channel losing flow reach exists seasonally in Bear Creek between the Bear Creek Road crossing and the North Tributary-3 confluence. North Tributary-3 is the long-term surface water monitoring location for the now-remediated Boneyard/Burnyard. Prior to the Boneyard/Burnyard remedial action, there were elevated mercury concentrations in the North Tributary-3 surface water. The Boneyard/Burnyard remedial action included excavation of source areas in the Boneyard/Burnyard and contaminated floodplain soils and sediments. Completion of the remedial action included backfill of the excavated area with clean soil and construction of a sinuous stream channel with riparian vegetation planting. There are areas of marshy conditions adjacent to the restored stream channel that appear to be conducive to mercury methylation, as indicated by the relatively higher methylmercury concentration in the North Tributary-3 surface water. North Tributary-3 exhibits the highest average and the highest maximum measured methylmercury concentration in the Bear Creek monitoring network. At BCK 11.54/BCK 11.54A, a short distance downstream from the North Tributary-3 confluence to Bear Creek, the second highest average and maximum methylmercury concentrations are observed.



A short distance downstream of BCK 11.54/BCK 11.54A, the flow of Bear Creek sinks into the epikarst subsurface flow system. During significant periods of the year, the Bear Creek channel is dry from the sink point downstream to the North Tributary-5 confluence. The North Tributary-5 total and methylmercury concentration data demonstrate relatively low inputs to Bear Creek.

Dye tracing conducted as part of the Bear Creek Valley remedial investigation demonstrated epikarst transport from this losing reach of Bear Creek emerges at a groundwater resurgence point at spring SS-4. The average and maximum total mercury concentrations at SS-4 and the maximum observed methylmercury concentration support the probable connection between the monitoring locations NT-3 and BCK 11.54/BCK 11.54A reach of higher concentrations.

Deep bedrock groundwater monitoring wells GW-704 (samples at 244.5 to 256 ft below ground surface) and GW-706 (samples at 156.1 to 182.5 ft below ground surface) lie between the Bear Creek sink point and the SS-4 spring and exhibit very low total mercury concentrations.

Downstream of the SS-4 spring discharge confluence, Bear Creek enters another significant losing flow reach that extends seasonally to the Bear Creek Road crossing adjacent to spring SS-5. Groundwater monitoring wells in this reach include wells GW-683 (sampling interval at 146 to 196.8 ft below ground surface) and GW-684 (sampling interval at 113.8 to 128.8 ft below ground surface). Total mercury concentrations in this well pair are higher than in the deeper wells, GW-704 and GW-706. Total mercury concentrations in wells GW-683 and GW-684 are similar to those measured at nearby spring SS-5. At spring SS-5, perennial flow exists that forms the head of continuous flow that extends downstream to Bear Creek kilometer 9.2 and beyond. Data from spring SS-6 may be somewhat misleading because during periods of high flow stage of Bear Creek, the spring area is sometimes inundated with main stem flow. Also, beaver ponding a short distance downstream of spring SS-6 caused inundation of the spring for a number of years and caused surface water backup into the epikarst of the area.

As previously noted, epikarst groundwater transport has been documented from the Bear Creek reach near BC-NT3 and BCK 11.54/BCK 11.54A and solids transport from Bear Creek to spring SS-4 has been observed.

Table 2 summarizes dissolved oxygen and redox potential measurements for most of the mercury and methylmercury monitoring locations. Surface water and spring discharges are well aerated, while groundwater from monitoring wells exhibits lower dissolved oxygen levels and wells GW-704 and GW-706 exhibit anoxic conditions consistent with their deep sample intervals. Electrochemically, most of the sample locations exhibit oxidizing conditions. Standard deviation of the average redox measurements is provided as an indicator of the relative variability of measurements at each location. Some, but not all of the groundwater monitoring well data exhibit electrochemically reducing conditions. Electrochemically reductive conditions in groundwater at the Oak Ridge Reservation are not uncommon.

Past mercury and methylmercury sampling events generally have not included field-filtered and unfiltered aliquots collected for analysis and have rarely had contemporaneous field turbidity or total suspended solids analyses performed. The primary sampling objectives to date have been the screening of total mercury concentrations for ambient water quality criteria (51 ng/L) comparisons, and ambient concentrations are usually much lower than the criterion, such that focus on partitioning of mercury and methylmercury between dissolved versus particle-associated was not a focus of investigation.



Although contemporaneous mercury and methylmercury data are not available for direct correlation with solids in the water samples, there are turbidity data from other sampling events during January 1, 2009, through June 30, 2024. Figure 2 is a graph that shows daily average flow rate versus turbidity data from locations BC-NT3, SS-5, and BCK9.2. At locations SS-5 and BCK9.2, data are amenable to separating turbidity versus flow values into lower versus higher flow conditions; however, this is not the case for BC-NT3. Figure 3 includes box and whisker plots for turbidity at locations SS-5, SS-4 (where no flow data are available), BC-NT3, and BCK 9.2. Monitoring location BCK 11.54/BCK 11.54A has too few turbidity measurements to provide useful interpretive value. Comparison of the methylmercury values with the box plots in Figure 3 suggests turbidity/suspended solids may be an important factor in aqueous transport.

Components of turbidity may include fine mineral sediment and/or organic materials that could contain significant mercury and/or methylmercury.

Although groundwater from monitoring wells along the Bear Creek floodplain have not been analyzed for methylmercury, the methylmercury distribution and apparent connections from demonstrated primary surface area aquatic sources (North Tributary-3) through the epikarst transport systems provide a coherent conceptual model of the available data.



Table 1. Summary of BCV surface water and groundwater detected mercury and methylmercury data, 2009–2024

Station	N	lercury (ng/	L)	Meth	nylmercury	(ng/L)	3.5.11	Constant	
name	n/ndet	Average	Max	n/ndet	Average	Max	Medium	Comment	
GW-276	9/9	101.74	440	-	-	-	GW		
GW-246	12/12	345.36	741	-	-	-	GW		
GW-815	2/2	310.50	410	-	-	-	GW	S-3 Area	
NT-1	21/15	13.14	55.7	-	-	-	SW	S 3 med	
BCK 12.34	43/43	14.68	127.2	32/32	0.11	0.258	BC SW - IP		
S07 (NT-2)	4/4	6.85	16.5	-	-	-	Trib SW		
		Bear	Creek los	ing reach	- sometimes	no flow			
BC-NT3	38/38	19.62	147	27/27	0.31	2.698	Trib SW		
BCK 11.54 / 11.54A	36/36	7.15	22.2	32/32	0.11	0.754	BC SW	BYBY residual	
NT-4	60/2	84.50	108	-	-	-	Trib SW	EMWMF office area	
EM-VWEIR	12/8	27.14	96	-	-	_	SW	EMWMF non- contact water	
NT-5	7/7	2.64	6.1	7/7	0.10	0.14	Trib SW	Receives V-Weir flow	
GW-706	19/19	0.99	1.94	-	-	-	GW	S-3 nitrate +	
GW-704	19/19	0.43	1.82	-	-	-	GW	uranium	
SS-4	52/32	15.61	87.7	28	0.10	0.39	Trib GW	Epikarst resurgence from BCK 11.54 sink	
BC-NT8	53/53	3.88	10.7	31/30	0.11	0.266	Trib SW	BCBG discharge	
GW-683	31/24	1.51	6.4	-	-	-	GW	S-3 nitrate +	
GW-684	31/27	1.15	8	_	<u>-</u>	_	GW	uranium	
GW-077	19/4	0.97	2.2	-	_	_	GW		
GW-078	19/18	0.60	2.1	-	-	-	GW	Zone 3 IP GW	
SS-5	52/36	2.80	6.64	32/16	0.04	0.15	Trib GW		



Table 1. Summary of BCV surface water and groundwater detected mercury and methylmercury data, 2009–2024 (cont.)

Station name	Mercury (ng/L)			Meth	ylmercury	(ng/L)	Medium	Comment		
	SS-5 f	low provide	es continuo	us baseflo	w in BC do	ownstream	to the IP			
BCK9.2	44/44	4.66	39.4	32/32	0.07	0.22	SW - IP	Zone 3 IP		
BCK-7.87	15/15	3.39	7.45	-			BC SW	Zone 2/Zone 1		
GW-712	19/2	1.64	3.13	-	-	-	GW			
GW-714	19/4	1.06	3.19	-	- -	<u>-</u>	GW	7 1 CW		
GW-713	31/8	5.02	38.7	-	-		GW	Zone 1 GW		
SS-6	12/12	13.89	145	-	-	_	Trib GW			
SS-7	4/4	1.05	1.6	-	-	_	Trib GW	 		
BCK 4.6	30/30	4.62	61	30/26	0.11	0.419	BC SW	Near		
BCK-4.55	15/15	2.92	7.6	-	-	-	BC SW	Highway 95		
SS-8	8/8	2.79	9.8	-	-	-	Trib GW	 		

<sup>-</sup> Denotes no data available.

BC = Bear Creek

BCBG = Bear Creek Burial Grounds

BCK = Bear Creek kilometer

BCV = Bear Creek Valley

BYBY = Boneyard/Burnyard

EMWMF = Environmental Management Waste Management Facility

GW = groundwater

IP = integration point

max = maximum

n = number of analytical results with detection limits < 51 ng/L

n/det = number of detected results

NT = North Tributary

SS = surface spring

SW = surface water

Trib = Bear Creek tributary sample location






Figure 1. Locations of selected surface water and groundwater monitoring sites in BCV.



Table 2. Summary of BCV surface water and groundwater dissolved oxygen and redox, 2009-2024

Station name	Diss	olved oxygen	(ppm)		Redox (mV	Comment			
	n	Average	St dev	n	Average	St dev	<del>-</del>		
GW-276	26	2.63	2.89	24	261	134			
GW-246	12	1.15	1.13	12	274	43			
GW-815	2	8.48	0.57	2	112	81	S-3 Area		
NT-1	20	8.06	3.11	20	164	103			
BCK 12.34	217	7.74	2.08	189	165	83			
S07 (NT-2)	7	6.92	3.28	7	160	60			
BC-NT3	28	8.40	3.66	19	136	99	DYDY 11 1		
BCK 11.54	4	6.49	2.70	4	135	54	BYBY residual		
NT-4	160	6.64	1.94	154	149	77	EMWMF office		
NT-5	7	7.91	2.31	7	179	50	area Receives V-Weir flow		
GW-706	31	0.31	0.38	29	112	90	S-3 nitrate +		
GW-704	32	0.61	0.34	30	59	134	uranium		
SS-4	39	7.39	1.31	39	173	70	Epikarst resurgence		
BC-NT8	41	7.98	2.40	39	146	96	BCBG discharge		
GW-683	34	4.49	2.16	30	94	101	S-3 nitrate +		
GW-684	31	2.48	2.74	29	91	102	uranium		
GW-077	31	1.04	0.69	29	7	86			
GW-078	31	2.52	1.93	31	68	94	Zone 3 IP GW		
SS-5	55	5.86	1.52	55	166	93			
BCK9.2	31	8.76	2.89	23	131	90	Zone 3 IP		
BCK7.87	37	8.01	2.03	37	142	116	Zone 2/Zone 1		
GW-712	31	1.48	3.84	29	-19	81			
GW-714	32	1.72	6.02	32	100	90	7 1 677		
GW-713	31	2.31	4.32	29	-76	82	Zone 1 GW		
SS-6	31	6.63	1.75	29	9 176 205				
SS-7	29	6.52	1.33	29	162	106			
BCK04.55	38	7.60	1.90	38	214	415	Near Highway 95		
SS-8	29	7.07	1.91	29	197	170			

BC = Bear Creek

BCBG = Bear Creek Burial Grounds

BCK = Bear Creek kilometer

BCV = Bear Creek Valley BYBY = Boneyard/Burnyard

EMWMF = Environmental Management Waste Management

Facility

GW = groundwater

IP = integration point

n = number of dissolved oxygen results

NT = North Tributary

ppm = parts per million

SS = surface spring

St dev = standard deviation






Figure 2. Daily average flow rate (Q) versus turbidity plots.




Figure 3. Box and whisker plots of turbidity data available for the data evaluation period 01/01/09-06/30/24.



# ATTACHMENT 2 NEW FIGURES AND TABLE FOR APPENDIX G BASED ON TDEC SPECIFIC COMMENT 3






Figure G.1. BCV transect locations and 100-year floodplain. (New figure based on TDEC specific comment 3)






Figure G.2. BCV transect area 1 and 100-year floodplain. (New figure based on TDEC specific comment 3)






Figure G.3. BCV transect area 2 and 100-year floodplain. (New figure based on TDEC specific comment 3)






Figure G.4. BCV transect area 3 and 100-year floodplain. (New figure based on TDEC specific comment 3)






Figure G.5. BCV transect area 4 and 100-year floodplain. (New figure based on TDEC specific comment 3)



Table G.1. Field observations of Bear Creek during transect sampling (New table based on TDEC specific comment 3)

Site identifier	Stream morphology
	- Stream is 384 in. wide
BCT1	- Stream depth is 14.5 in.
ВСТТ	- Sandbar extends 138 in. into the stream
	- Bank is 30 in. above the water's surface
	- Stream is 248 in. wide
	- Banks are 360 in. apart
BCT2	- Gravel bars are present along the edges of the water
	- Bank is 50 in. above the water's surface
	- Stream is straight and has moderate flow
	-Stream is 192 in. wide
	- Stream depth is 13 to 36 in.
BCT3	- Banks are extremely steep
Ders	- Channel is straight and has moderate flow
	- Soil/sediment samples are collected approximately 50 yd downstream of culvert due to banks being riprap/exposed
	bedrock
	- Stream is 264 in. wide
	- Stream depth is 20 to 42 in.
BCT4	- South bank is 12 in. above the water's surface; north bank is 4 to 6 in. above the water's surface
	- Stream is meandering and has slow flow
	- Due to ponding at beaver dam, samples collected approximately 150 yd upstream of the planned location
	- Stream is 216 in. wide
	- Stream depth is 27 in.
BCT5	- Bank is 24 in. above the water's surface
	- Gravel bar extends 36 in. into the channel at the sample point
	- Stream has slight curves with high flow
	- Beaver dam near this location was blown out/destroyed, presumably from extreme flow from recent storm event



Table G.1. Field observations of Bear Creek during transect sampling (New table based on TDEC specific comment 3) (cont.)

Site identifier	Stream morphology
	- Stream is 348 in. wide - Stream depth is 18 in.
BCT6	<ul> <li>Bank is 24 in. above the water's surface</li> <li>Debris bank (gravel/rock/sediment, sticks/wood, or leaf litter) extends 96 in. into the channel at the sample point</li> </ul>
	- Debits bank (grave/rock/sediment, sticks/wood, of leaf inter) extends 90 m. into the channel at the sample point - Stream has slight curves with high flow
	- Beaver dam near this location was blown out/destroyed, presumably from extreme flow from recent storm event
	- Stream is 324 in. wide
D CTT	- Stream depth at middle of channel is unobtainable due to ponding/depth/safety (estimated to be at least 60 in. deep)
BCT7	- No true bank, only floodplain, due to ponding
	<ul><li>Stream is meandering and has slow flow</li><li>Site is directly below beaver dam</li></ul>
	- Stream is 396 in. wide
	- No defined bank, only floodplain, due to ponding
BCT8	- Edges of channel are 24 in. deep, and the deepest point is at least 60 in. (too deep to safely measure)
	- Channel is straight and has slow flow
	- Samples collected upstream of beaver dam at Haul Road intersection  - Stream is 456 in, wide
	- Stream is 436 in. wide - Edges of channel are 6 in. deep. Stream depth at middle of channel is unobtainable due to depth/safety concerns
ВСТ9	- No defined bank due to ponding
	- Channel is straight and has slow flow
	- Site is above beaver dam
	- Stream is 216 in. wide
	- Banks are 304 in. apart
BCT10	- Stream depth is 10 in.  Debria (convert/godingent, sticks/yyood, on loof litter) by ildyn is 82 in yyide
BC 110	<ul> <li>Debris (gravel/rock/sediment, sticks/wood, or leaf litter) buildup is 82 in. wide</li> <li>Banks are 30 in. above the water's surface</li> </ul>
	- Channel is straight and has high flow
	- Flow had decreased (post-storm event) and stream depth was 6 in. during a second sampling event



Table G.1. Field observations of Bear Creek during transect sampling (New table based on TDEC specific comment 3) (cont.)

Site identifier	Stream morphology
	- Stream is 180 in. wide - Stream depth at sample point is 8 in., but pools that reach 42 in. deep are present
BCT11	- Stream depth at sample point is 8 m., but pools that reach 42 m. deep are present  - Bank is 12 in. above the water's surface
БСТТ	- Debris (gravel/rock/sediment, sticks/wood, or leaf litter) buildup is 48 in. wide
	- Channel is curvy and has slow flow due to debris buildup
	- Stream is 60 in. wide
	- Banks are 132 in. apart
	- Stream depth of sample point is 6 in.; however, deeper pockets exist upstream and downstream that are 12 in. deep
BCT12A	- Gravel/sand bar is 72 in. wide
	- Bank is 50 in. above the water's surface
	- Channel is straight and has high flow
	- Flow had decreased (post-storm event) and water depth was 4 in. during a second sampling event
	- Stream is 96 in. wide
	- Banks are 168 in. apart
	- Stream depth at sample point is 10 in.
BCT12B	- Gravel/sand bar is 72 in. wide
	- Channel is straight and has high flow
	- Bank is 60 in. above the water's surface
	- Flow had decreased (post-storm event) and water depth was 6 in. during a second sampling event
	- Stream is 36 in. wide
	- Banks are 96 in. apart
BCT13	- Stream depth is 8 in.
	- Banks are 30 in. above the water's surface
	- Channel is curvy and has high flow
	- Flow had decreased (post-storm event) and water depth was 4 in. during a second sampling event  - Stream is 120 in. wide
	- Water is 16 in. deep at the sample point directly following a dropoff in the channel
BCT14	- Water is 70 iii. deep at the sample point directly following a droport in the channel  - Bank is 72 in. above the water's surface
	- Channel is straight and has moderate flow
	Chainer is straight and has inoderate now



#### Table G.1. Field observations of Bear Creek during transect sampling (New table based on TDEC specific comment 3) (cont.)

Site identifier	Stream morphology						
	- Stream is 36 in. wide - Stream depth is 4 in.						
BCT15	- Stream depth is 4 in Bank is 24 in. above the water's surface						
	- Stream is meandering with a moderate/average flow rate for that location						
HCTREF	- Stream is 372 in. wide						
(Hinds Creek	- Stream depth is 13 in.						
transect	- Bank is 24 in. above the water's surface						
reference	- Stream is straight and has fast flow						
site)	- Bottom of the channel is exposed bedrock with small pockets of gravel						

BCT = Bear Creek transect





# **ATTACHMENT 3**



Table 3.9. Results of mercury sequential extraction analyses from floodplain soil, creek bank soil, and channel sediment at eight BCT locations and at the Hinds Creek reference site (Revised table based on TDEC specific comment 5)

			BCT14			BCT12B			BCT11			ВСТ9		ВСТ8		
Chemical name	Units	Result	Percent of sum	Lab qual	Result	Percent of sum	Lab qual	Result	Percent of sum	Lab qual	Result	Percent of sum	Lab qual	Result	Percent of sum	Lab qual
					Floo	dplain soil										
Mercury (F0) – Volatile Elemental Mercury	μg/kg	0.18	0%	UJ	0.18	0%	UJ	0.17	0%	UJ	0.56	0%	J	0.22	0%	UJ
Mercury (F1) – Water Soluble Mercury	μg/kg	20	6%	J	54	3%	J	25	3%		13	6%		3.9	3%	J
Mercury (F2) – pH 2 Soluble Mercury	μg/kg	9.7	3%	J	40	3%	J	17	2%		3.3	2%	J	1.9	2%	J
Mercury (F3) – 1N Potassium Hydroxide Extractable Mercury	μg/kg	270	79%	J	850	53%	J	550	73%		160	73%		99	76%	
Mercury (F4) – 12N Nitric Acid Soluble Mercury	μg/kg	37	11%	J	560	35%	J	110	15%		31	14%		18	14%	J
Mercury (F5) – Aqua Regia Soluble Mercury	μg/kg	4.7	1%	J	53	3%	J	45	6%		11	5%		3.6	3%	J
Mercury (F6) – Mineral-bound Mercury	μg/kg	1.7	1%	J	2.6	0%	J	0.73	0%	J	1	1%	J	0.4	0%	J
Mercury (FS) <sup>a</sup> – Total Mercury by Summation	μg/kg	340	101%	J	1600	98%	J	750	100%		220	100%	J	130	98%	J
					В	ank soil										
Mercury (F0) – Volatile Elemental Mercury	μg/kg	0.17	0%	UJ	0.17	0%	UJ	0.18	0%	J	0.18	0%	UJ	0.2	0%	UJ
Mercury (F1) – Water Soluble Mercury	μg/kg	1.1	0%	UJ	26	3%	J	7.1	4%		8.8	4%		2.8	3%	J
Mercury (F2) – pH 2 Soluble Mercury	μg/kg	1.1	0%	UJ	20	2%	J	3.5	2%	J	4.9	2%		2.1	3%	J
Mercury (F3) – 1N Potassium Hydroxide Extractable Mercury	μg/kg	2.2	0%	UJ	390	46%	J	120	75%		200	80%		65	77%	
Mercury (F4) – 12N Nitric Acid Soluble Mercury	μg/kg	17	81%	J	150	18%	J	15	9%	J	25	10%		12	14%	J
Mercury (F5) – Aqua Regia Soluble Mercury	μg/kg	3.1	15%	J	250	30%	J	9.3	6%		6.8	3%		3	4%	J
Mercury (F6) – Mineral-bound Mercury	μg/kg	1.3	6%	J	6.8	1%	J	0.93	1%	J	0.82	0%	J	0.36	0%	UJ
Mercury (FS) <sup>a</sup> – Total Mercury by Summation	μg/kg	21	102%	J	840	100%	J	160	98%	J	250	99%	J	85	100%	J
					Chanı	nel sediment										
Mercury (F0) – Volatile Elemental Mercury	μg/kg	7.4	14%	J	0.19	0%	J	0.52	1%	J	0.18	0%	UJ	0.2	0%	UJ
Mercury (F1) – Water Soluble Mercury	μg/kg	1.3	0%	UJ	6.5	3%		5.3	6%		8.6	4%		3.4	4%	J
Mercury (F2) – pH 2 Soluble Mercury	μg/kg	1.3	0%	UJ	5.6	3%	J	4.3	4%	J	4.8	2%		1.3	2%	J
Mercury (F3) – 1N Potassium Hydroxide Extractable Mercury	μg/kg	2.7	0%	UJ	99	45%		37	38%		170	71%		70	80%	
Mercury (F4) – 12N Nitric Acid Soluble Mercury	μg/kg	32	58%	J	20	9%		8.3	9%	J	43	18%		9.6	11%	J
Mercury (F5) – Aqua Regia Soluble Mercury	μg/kg	23	42%	J	85	39%		40	41%		11	5%		2.7	3%	J
Mercury (F6) – Mineral-bound Mercury	μg/kg	0.46	1%	J	0.91	0%	J	1.9	2%	J	1.2	1%	J	0.71	1%	J
Mercury (FS) <sup>a</sup> – Total Mercury by Summation	μg/kg	55	101%	J	220	99%	J	97	100%	J	240	99%	J	88	100%	J



Table 3.9. Results of mercury sequential extraction analyses from floodplain soil, creek bank soil, and channel sediment at eight BCT locations and at the Hinds Creek reference site (cont.) (Revised table based on TDEC specific comment 5)

		BCT7				BCT6			BCT5			ek average	HCTREF		
Chemical name		Result	Percent of sum	Lab qual	Result	Percent of sum	Lab qual	Result	Percent of sum	Lab qual	Average result	Average percent	Result	Percent of sum	Lab qual
					F	loodplain soil	-			-					
Mercury (F0) – Volatile Elemental Mercury	μg/kg	0.2	0%	UJ	1.4	5%	J	0.17	0%	UJ	0.25	1%	0.22	1%	J
Mercury (F1) – Water Soluble Mercury	μg/kg	3.4	3%	J	2.5	8%	J	3.2	3%	J	15.6	4%	1.1	0%	UJ
Mercury (F2) – pH 2 Soluble Mercury	μg/kg	1.6	1%	J	1.2	0%	UJ	2	2%	J	9.4	2%	1.1	0%	UJ
Mercury (F3) – 1N Potassium Hydroxide Extractable Mercury	μg/kg	110	85%		26	84%		83	84%		269	76%	18	62%	
Mercury (F4) – 12N Nitric Acid Soluble Mercury	μg/kg	15	12%	J	5.9	0%	UJ	6.2	6%	J	97	13%	5.5	0%	UJ
Mercury (F5) – Aqua Regia Soluble Mercury	μg/kg	2.9	2%	J	2.5	8%	J	3.5	4%	J	16	4%	1.2	4%	J
Mercury (F6) – Mineral-bound Mercury	μg/kg	0.47	0%	J	0.42	1%	J	0.76	1%	J	1.0	1%	10	35%	J
Mercury (FS) <sup>a</sup> – Total Mercury by Summation	μg/kg	130	103%	J	31	102%	J	99	100%	J		101%	29	101%	J
						Bank soil									
Mercury (F0) – Volatile Elemental Mercury	μg/kg	0.19	0%	UJ	0.18	0%	UJ	0.93	1%	J	0.1	0%	0.37	1%	J
Mercury (F1) – Water Soluble Mercury	μg/kg	3.3	3%	J	1.5	6%	J	4.4	5%	J	6.7	4%	2.8	9%	J
Mercury (F2) – pH 2 Soluble Mercury	μg/kg	1.3	1%	J	1.2	0%	UJ	1.6	2%	J	4.2	2%	1.1	0%	UJ
Mercury (F3) – 1N Potassium Hydroxide Extractable Mercury	μg/kg	86	78%		19	76%		68	79%		118.5	64%	13	39%	
Mercury (F4) – 12N Nitric Acid Soluble Mercury	μg/kg	14	13%	J	5.9	0%	UJ	6.8	8%	J	30	19%	5.3	0%	UJ
Mercury (F5) – Aqua Regia Soluble Mercury	μg/kg	2.2	2%	J	4.5	18%		4.4	5%		35.4	10%	1.2	4%	J
Mercury (F6) – Mineral-bound Mercury	μg/kg	0.34	0%	UJ	0.48	2%	J	0.72	1%	J	1.4	1%	16	49%	J
Mercury (FS) <sup>a</sup> – Total Mercury by Summation	μg/kg	110	97%	J	25	102%	J	86	100%	J		100%	33	100%	J
					Ch	annel sedimen	ţ								
Mercury (F0) – Volatile Elemental Mercury	<u>μ</u> g/kg	0.21	0%	UJ	NA			NA			1.4	2%	NA		
Mercury (F1) – Water Soluble Mercury	μg/kg	1.6	3%	J	NA			NA			4.25	3%	NA		
Mercury (F2) – pH 2 Soluble Mercury	μg/kg	1.6	3%	J	NA			NA			2.9	2%	NA		
Mercury (F3) – 1N Potassium Hydroxide Extractable Mercury	μg/kg	45	88%		NA			NA			70.2	54%	NA		
Mercury (F4) – 12N Nitric Acid Soluble Mercury	$\mu g/kg$	6.6	0%	UJ	NA			NA			18.8	17%	NA		
Mercury (F5) – Aqua Regia Soluble Mercury	μg/kg	2.4	5%	J	NA			NA			27.4	22%	NA		
Mercury (F6) – Mineral-bound Mercury	μg/kg	0.38	0%	UJ	NA			NA			0.86	1%	NA		
Mercury (FS) <sup>a</sup> – Total Mercury by Summation	μg/kg	51	99%	J	NA			NA				100%	NA		

<sup>a</sup>Mercury (FS) sum does not include Mercury (F0) aliquot because volatile mercury is measured on a separate sample aliquot. Laboratory reporting at two-significant-figure level accounts for summed percentages greater than or less than 100%.

<sup>--</sup> Indicates no qualifier; detection, or not applicable.

BCT = Bear Creek transect

F = fraction

HCTREF = Hinds Creek transect reference site

J = estimated value

Lab = laboratory

N = normalityNA = not analyzed

qual = qualifier

UJ = not detected at estimated value






Figure 3.11. Results of Bear Creek floodplain soil sequential mercury extraction. (Revised figure based on TDEC specific comment 5)






Figure 3.12. Results of Bear Creek bank soil sequential mercury extraction. (Revised figure based on TDEC specific comment 5)






Figure 3.13. Results of Bear Creek channel sediment sequential mercury extraction. (Revised figure based on TDEC specific comment 5)






Figure 3.14. Sequential mercury extraction results at location BCT5. (Revised figure based on TDEC specific comment 5)






Figure 3.15. Sequential mercury extraction results at location BCT6. (Revised figure based on TDEC specific comment 5)






Figure 3.16. Sequential mercury extraction results at location BCT7. (Revised figure based on TDEC specific comment 5)






Figure 3.17. Sequential mercury extraction results at location BCT8. (Revised figure based on TDEC specific comment 5)






Figure 3.18. Sequential mercury extraction results at location BCT9. (Revised figure based on TDEC specific comment 5)






Figure 3.19. Sequential mercury extraction results at location BCT11. (Revised figure based on TDEC specific comment 5)






Figure 3.20. Sequential mercury extraction results at location BCT12B. (Revised figure based on TDEC specific comment 5)






Figure 3.21. Sequential mercury extraction results at location BCT14. (Revised figure based on TDEC specific comment 5)







Figure 3.22. Sequential mercury extraction results at the Hinds Creek reference site (HCTREF). (Revised figure based on TDEC specific comment 5)





Methylmercury flux comparison at SS-5 and BCK 11.54/11.54A. (Supplemental figure to address TDEC specific comment 6)